

A. I. Kydyrmanov, K. O. Karamendin

Scientific and Production Center of Microbiology and Virology, Almaty, Kazakhstan.

E-mail: aidyn.kydyrmanov@gmail.com, kobey.karamendin@gmail.com

VIRUSES OF MARINE MAMMALS AND METAGENOMIC MONITORING OF INFECTIOUS DISEASES

Abstracts. The review presents information on epizootic diseases in marine mammals caused by influenza viruses of genera A and B, morbilliviruses, adeno-, herpes-, calici- and coronaviruses. Virological and serological data about the spectrum of viral infections in marine mammals are given. Describes epizootic events in marine mammals induced by influenza A viruses and isolation of strains with antigenic formulas H1N1, H1N3, H3N3, H3N8, H4N5, H4N6, H7N7, H10N7, H13N2, H13N9. It is suggested that the seals may be infected with a wide range of influenza viruses by direct transmission without prior adaptation. It is emphasized that marine pinnipeds can serve as one of the reservoirs of human influenza B virus in nature. Data on the classification of pinniped and cetacean coronaviruses are presented. A variety of pinnipeds herpesviruses and the clinical significance of their morbidity indicators in the diagnosis of viral infections of seals are shown. It emphasizes the need for the development of proactive diagnostic tools for the timely detection of new viruses with zoonotic potential that can cause disease outbreaks. The negative effects of diseases with unknown etiologies on the population of Caspian seals are noted. The conclusion is made about the need for metagenomic monitoring of Caspian seal virome in order to establish the role of viral pathogens in their morbidity and regular mortality under conditions of intensive habitat development.

Key words: virus, marine mammals, Caspian seal, virom, metagenomic monitoring.

Health survey of marine mammal populations is one of the most important and significant scientific directions in ecology. Current knowledge about the diversity of pathogens of marine mammals is expanding rapidly [1, 2]. Viral diseases play a major role in regulating the dynamics of the population of wild animals, limiting their growth and enhancing selection at the genetic level. The impact of viruses becomes even more severe to the species endangered or fragmented at the result of anthropogenic pressure.

Viruses from marine mammals were isolated for the first time in 1972 from Californian sea lion (*Zalophus californianus*), on San Miguel Island during an outbreak of infection among these animals. The etiological agent attributed to caliciviruses and adopted as the San Miguel type I sea lion virus. They were isolated from 11 species of pinnipeds and cetaceans [3].

Paramyxovirus infection caused by members of the genus morbillivirus is widespread among marine mammals and causes serious threat to the population of these animals. To date, four species of morbilliviruses that infect marine mammals have been described: canine distemper virus in the Baikal (*Pusasibirica*) and Caspian seals (*Pusacaspica*); cetacean morbillivirus - in dolphins and whales; phocine distemper virus - in harbor seals (*Phocavitulina*); porpoise morbillivirus- in porpoises (*Phocoenaphocoena*).

Morbilliviruses in marine mammals were first discovered in the late 1980s during the mass mortality of harbor seals and gray seals (*Halichoerus grypus*) on the North European coasts in 1987-1988.

Phocine distemper virus (PDV) caused widespread epizootics among harbor seals in the coastal waters of Northern Europe in 1988, which killed more than 12 thousand animals [4]. In 2002, this virus returned to the indicated population, causing the death of about 1000 animals on the coasts of Scandinavia [5].

Morbillivirus epizootics (including canine distemper virus) occurred in populations of pinnipeds and cetaceans of the Northern Hemisphere [6]. The canine distemper virus caused the mass death of the Baikal seal (*Pusasibirica*) in 1987-1988 in Siberia, this virus was also involved in the mortality of crabeater seals

(*Lobodoncarcinophagus*) in 1955 in Antarctica. It was thought that the virus was transmitted to the population of pinnipeds through contact with domestic dogs [7].

Morbillivirus infections with high mortality have been registered among porpoises, striped dolphins (*Stenella coeruleoalba*) and bottlenose dolphins (*Tursiops truncatus*) [8, 9], harbor seals and harp seals (*Pagophilus groenlandicus*) [10]. The mass die-off of Caspian seals in the spring and summer of 2000 was caused by canine distemper virus [11].

An infection potential of influenza A virus for the seals was made known from the epizootic outbreak on the coast of New England in 1979-1980 resulted death of 600 animals [12]. The genetic characterization of the virus showed that all its RNA segments showed a close resemblance to those of avian strains. The second epizooty among the seals, occurring with pronounced clinical manifestations of pneumonia, occurred on the coast of New England in June 1982-August 1983 was associated with the influenza A virus (H4N5) [13].

Five strains of influenza A virus were isolated from seals that died from pneumonia on the Cape Cod Peninsula (Massachusetts, USA) from January 1991 - February 1992, two of which were identified as A (H4N6), and three as A H3N3. The HA genes of influenza A(H3N3) viruses were 99.7% identical to each other. Phylogenetic analysis demonstrated a close relationship between their sequences and those of the hemagglutinin H3 avian isolate. These data indicate that the influenza virus subtype H3 has been circulating for a long time in seal populations [14].

Identification of various types of influenza viruses in the seal populations indicates the possibility of their participation in genetic reassortment. Molecular-genetic studies established an avian origin of seal influenza isolates [13].

Pandemic H1N1 influenza A virus isolated from northern elephant seals (*Mirounga angustirostris*), near the California coast of the United States in 2010. Full genome sequencing revealed more than 99% homology with pandemic "swine" influenza virus A/California/04/2009 which circulated among people since 2009 [15].

In the fall of 2011 in New England (Massachusetts, USA), 162 seals died from pneumonia caused by the influenza A virus (H3N8). The pathogen was found to be similar to influenza viruses of waterfowl that circulated in North America since 2002 and had a mutation in the PB2 gene, characteristic of the highly pathogenic for humans variant H5N1, which indicated its potential for interspecies transmission and adapting to mammals [16].

Until recently, influenza A viruses did not isolate from the pinnipeds of the Palearctic. Previously, virological evidence of their involvement in outbreaks of infection among seals was obtained only in the North American continent.

Epizootics of marine mammals caused by influenza A viruses in other parts of the world, were first reported in 2014. The influenza A virus (H10N7) is isolated from found dead harbor seals (*Phocavulina*) on the North Sea coast in Sweden, Denmark, Germany [17-19] where over 1400 animals died.

Influenza A (H1N3) virus is isolated D.K. Lvov et al. [20] from a dwarf whale (*Balaenoptera acutorostrata*) in the South Pacific in 1975-1976. From toothed whales (long-finned pilot whale - *Globicephal melas*) V. Hinshaw et al. [21] identified the H13N9 and H13N2 influenza viruses genetically close to the H13 viruses circulating among gulls.

The data about isolation of influenza A virus (H7N7) from the materials collected during mass mortality of Caspian seals in April-June 2000-2002 was reported by Shestopalov A.M. et al. [22, 23] and Chuvakova Z.K. et al. [24, 25] in conference papers and review article. Despite of an extreme importance of those findings any other data about further characterization of genetic or pathobiological peculiarities of epizootic strain have not been published.

Later, influenza A (H4N6) viruses were isolated from seals in the Russian waters of the Caspian Sea in 2002 and 2012 [26, 27]. Phylogenetic analysis demonstrated that all genes of seal origin A(H4N6) were closely related to avian-derived influenza viruses of the classical Eurasian lineage circulating in wild birds.

In contrast to influenza A viruses, which have been isolated from many different species, influenza B viruses were human pathogens with no or unknown reservoirs in nature until 1999. However, further it was argued that common and gray seals can become infected with this virus [28]. Influenza virus B was isolated from juvenile harbor seal with symptoms of respiratory disease and *in vitro* infected cell culture of

phocine kidney cells. Since the moment of acknowledgment of marine mammals as new hosts, there have been several reports on the detection of antibodies to the influenza B virus in some species of *Otariid* and *Phocid* species [29, 30]. Based on these data, it was suggested that the seals may serve as a reservoir of the human influenza B virus.

Adenoviral infections with clinical signs of hepatitis and enteritis have been noted in sea lions [31]. Outbreaks of this infection have also been observed in fur seals, sei whales (*Balaenoptera borealis*), bowhead whales (*Balaena mysticetus*), beluga whales (*Delphinapterus leucas*), bottlenose dolphins [32-34].

Herpesviruses are isolated from seals with clinical signs of acute pneumonia and hepatitis in the Netherlands in 1985. [35]. They are subsequently isolated from bottlenose dolphins, killer whales (*Orcinus orca*), California sea lions and gray seals [36]. At present, seven species of herpes viruses (PhHV-1 - PhHV-7), belonging to the genera alpha and gamma herpes viruses, have been isolated from phocids [35-40]. In the infectious pathology of these animals, herpesvirus seals 1 [Phocid Herpesvirus type 1] (PhHV-1) of the genus α -herpesviruses are most significant pathogen. Infection caused by these viruses in the harbor seals and gray seals is characterized by respiratory manifestations with interstitial pneumonia and coagulative necrosis of the adrenal tissues and liver [35, 41].

The rest six herpesviruses (PhHV-2 - PhHV-7) belong to the genus γ - herpes viruses, of which PhHV-4 and PhHV-7 have diagnostic importance in inflammation and ulceration of oral cavity soft tissues of seals. PhHV-6 herpesvirus in seals is associated with eye disorders [39]. The role of the PhHV-2, PhHV-3 and PhHV-5 viruses in the infectious pathology of seals remains unknown.

Herpesvirus PhHV-1 was isolated from newborn harbor seals with clinical signs of acute pneumonia and hepatitis at a rehabilitation center in the Netherlands in 1985 [35]. In the Pacific harbor seal (*Phoca vitulina richardsii*); this virus is associated with focal coagulatory necrosis of the adrenal cortex and liver. In phylogenetic studies of PhHV-1 DNA polymerase (gB) gene, their close relationship with herpesviruses of dogs and cats has been established.

Serological studies have shown widespread infection of PhHV-1 or other closely related α -herpes viruses in the free-ranging seal populations of the North Sea, as well as in the waters of the Antarctic and the North Pacific [42-44].

The results of virological screening, molecular genetics and serological studies indicate the circulation of PhHV-1 among Caspian seals and its possible involvement in their infectious pathology and recurrent mortality [45].

Coronaviruses are common pathogens of the respiratory system, gastrointestinal tract and cause systemic infections of domestic animals and wildlife.

Coronaviruses in marine animals were first described during seal mortality investigation in an aquarium of Florida [46], as well as among free-ranging pinnipeds along the coast of central California [47]. In harbor seals (harbor seal coronavirus - HSCoV), the infection is manifested by hemorrhagic pneumonia and causes epizootics with high mortality. In cetaceans, coronaviruses were isolated from the beluga whales in captivity [47] and from the feces of three Indo-Pacific dolphins (*Tursiops aduncus*) [48].

HSCoV belongs to the genus of α -coronaviruses, closely related with coronaviruses of cats, dogs, ferrets and pigs [1]. Beluga whale coronavirus and dolphin coronavirus are γ -coronaviruses and represent a special type of cetacean coronavirus [50].

In addition to the listed viruses that infect marine mammals, there are reports of astro- and poxvirus zoonoses of cetaceans and pinnipeds [49-51].

Taken together, the available evidence suggests that the causative agents of these diseases are detected only after mass outbreaks of infections that have reached an epizootic scales. Therefore, it requires proactive diagnostic tools for the timely detection of new viruses with zoonotic potential capable of causing outbreaks of diseases, in order to develop an effective strategy to combat the devastating epizootics of vulnerable animal species.

Routinely, for the diagnosis of viral diseases, cell cultures are used to isolate pathogens, specific antibody based immunological tests, and polymerase chain reaction (PCR) for amplifying viral gene segments [52]. Unfortunately, all of them turned out to be ineffective in identifying previously unknown viral pathogens due to their unculture ability and lack of diagnostic antibodies to them. The possibilities of PCR diagnostics in such cases are also limited due to the need for specific oligonucleotide primers complementary to the conserved sequence of the pathogen genome fragment.

Unlike the above methods of identifying pathogens, viral metagenomics allows to identify new viruses without prior knowledge of their genome sequences. Viral metagenomics is used to directly identify new etiological agents from the affected animal tissues with pathological alterations [53, 54]. For example, in metagenomic studies of Ng et al. identified annelovirus (ZcAV) infection in two dead sea lions [55]. The authors showed that ZcAV is found only in the lungs and pleural cavity, absent in the tonsils, lymph nodes, liver and other organs. This indicates that metagenomics can be used to determine the tissue tropism of new viruses. Subsequent studies have shown that the prevalence of the virus was high during outbreaks among sea lions in captivity (100%), low in wild representatives (11%); all this indicates the possibility of using metagenomic studies to detect viruses in marine animals. In addition to ZcAV, seal anneloviruses and fur seal sacobuviruses have been discovered using metagenomic studies [56]. Bodewes et al. [40] using metagenomic studies identified herpesvirus seal type 7 as an etiologic agent for ulcerative gingivitis in seals. With a similar method of analysis, it is also shown that Enhydryalutis papilloma virus 1 is the etiological agent of oral cavity tumors in sea otters.

In the context of the above provided, the Caspian seal, as an endemic marine mammal, is one of the indicators of the Caspian Sea ecosystem. Diseases of unknown etiology [57-59] have a negative effect on its population, which indicates the importance of metagenomic studies of Caspian seal viromes to establish the role of viral pathogens in their morbidity and regular mortality in conditions of intensive habitat development.

А. И. Кыдырманов, К. О. Карамендин

Микробиологии және вирусология ғылыми өндірістік орталығы, Алматы, Қазақстан

ТЕҢІЗ СҮТҚОРЕКТІЛЕРІНІҢ ВИРУСТЫҚ ИНФЕКЦИЯЛАРЫ ЖӘНЕ МЕТАГЕНОМИКАЛЫҚ МОНИТОРИНГТЕ ЖАҢА ВИРУСТАРДЫ ТАБУ

Аннотация. Шолуда теңіз сүтқоректілеріндегі тұмау А және В вирустары, морбилливирустар, адено-, герпес-, калици- және короновирустар қоздырған індеттер туралы мағлұмат берілген. Теңіз сүтқоректілерінің вирустық инфекцияларының ауқымы жайында вирусологиялық және серологиялық деректер келтірілген. Теңіз сүтқоректілердегі тұмау А вирустарының індетімен олардан антигендік формулалары H1N1, H1N3, H3N3, H3N8, H4N5, H4N6, H7N7, H10N7, H13N2, H13N9 штамдарды оқшаулап алу жағдайлары сипатталады. Итбалықтардың алдын-ала бейімделмеген тұмау вирусының ауқымды спектрін тікелей жұқтыру мүмкіншілігі топшыланған. Теңіз ескекаяқтылары адамның В тұмауы вирусының табиғаттағы сақтаушы-сыдеген түсініке басымдық берілген. Ескекаяқтылар мен киттәрізділер короновирустарының классификациясы туралы мәліметтер ұсынылған. Итбалықтар герпесвирустарының түр алуандығы мен олар туғызатын ауруларды балауда клиникалық белгілерінің маңызы көрсетілген. Ауру тудыруы мүмкін зооноздық әлеуеті бар жаңа вирустарды дер кезінде анықтау үшін ұтқыр диагностикалық құралдарды әзірлеу қажеттілігіне баса назар аударды. Каспий итбалықтары популяциясына этиологиясы белгісіз аурулардың теріс әсерлері жөнінде айтылған. Шолуға тіршілік ортасын қарқынды игеру жағдайындағы каспий итбалығы ауруларымен тұрақты өлім-жітіміне вирустардың қатысын зерделеу үшін олардың виромына метагеномикалық мониторинг жүргізу қажет деген қорытынды жасалынған.

Түйін сөздер: вирус, теңіз сүтқоректілері, каспий итбалығы, вирус, метагеномдық мониторинг.

А. И. Кыдырманов, К. О. Карамендин

Научно производственный центр микробиологии и вирусологии, Алматы, Казахстан

ВИРУСНЫЕ ИНФЕКЦИИ МОРСКИХ МЛЕКОПИТАЮЩИХ И ОБНАРУЖЕНИЕ НОВЫХ ВИРУСОВ ПРИ МЕТАГЕНОМНОМ МОНИТОРИНГЕ

Аннотация. В обзоре представлены сведения об эпизоотиях среди морских млекопитающих, вызванных вирусами гриппа родов А и В, морбилливирусами, адено-, герпес-, калици-, короновирусами. Приводятся вирусологические и серологические данные о спектре вирусных инфекций морских млекопитающих. Описываются случаи эпизоотии морских млекопитающих вирусами гриппа А и изоляция от них штаммов с

антигенными формулами H1N1, H1N3, H3N3, H3N8, H4N5, H4N6, H7N7, H10N7, H13N2, H13N9. Высказывается предположение о возможности инфицирования тюленей широким спектром вирусов гриппа путем прямой трансмиссии без предварительной адаптации. Подчеркивается положение о том, что морские ластоногие могут служить одним из резервуаров вируса гриппа В человека в природе. Приведены данные о классификации коронавирусов ластоногих и китообразных. Показано разнообразие герпесвирусов ластоногих и клиническая значимость показателей их заболеваемости в диагностике вирусных инфекций тюленей. Подчеркивается потребность в разработках упреждающих диагностических инструментов для своевременного выявления новых вирусов с зоонозным потенциалом, способных вызывать вспышки болезней. Отмечаются отрицательные воздействия заболеваний с невыясненными этиологиями на популяцию каспийских тюленей. Делается заключение о необходимости проведения метагеномного мониторинга виroma каспийских тюленей для установления роли вирусных патогенов в их заболеваемости и регулярной смертности в условиях интенсивного освоения среды обитания.

Ключевые слова: вирус, морские млекопитающие, каспийский тюлень, виром, метагеномный мониторинг.

Information about authors:

Kydyrmanov Aidyn I., Doctor of Vet. Sci., LLP «SPC Microbiology and Virology», Almaty, Kazakhstan; aidyn.kydyrmanov@gmail.com; <https://orcid.org/0000-0002-8374-6128>

Karamendin Kobey O., Candidate of Vet. Sci., LLP «SPC Microbiology and Virology», Almaty, Kazakhstan; kobey.karamendin@gmail.com; <https://orcid.org/0000-0003-0829-3330>

REFERENCES

- [1] Nollens H., Wellehan J., Archer L., Lowenstine L., Gulland F. (2010) Detection of a respiratory coronavirus from tissues archived during a pneumonia epizootic in free ranging Pacific harbor seals *Phocavitularichardsii* // *Dis Aquat Organ.* 90:113-120. DOI: 10.3354/dao02190 (in Eng.).
- [2] Palacios G., Wellehan J., Raverty S., Bussetti A., Hui J., Savji N., Nollens H., Lambourn D., Celone C., Hutchison S., Calisher C., Nielsen O., Lipkin W. (2011) Discovery of an orthoreovirus in the aborted fetus of a Steller sea lion (*Eumetopiasjubatus*) // *J. Gen Virol.* 92:2549-57. DOI: 10.1099/vir.0.032649-0 (in Eng.).
- [3] Smith A.W., Skilling D.E., Cherry N., Mead J.H., Matson D.O. (1998) Calicivirus emergence from ocean reservoirs: zoonotic and interspecies movements // *Emerging Infect Dis.* 4:13. DOI: 10.3201/eid0401.980103 (in Eng.).
- [4] Osterhaus A.D., Vedder E.J. (1988) Identification of virus causing recent seal deaths // *Nature.* 335:20. DOI: 10.1038/335020a0 (in Eng.).
- [5] Jensen T., van de Bildt M., Dietz H.H., Andersen T.H., Hammer A.S., Kuiken T., Osterhaus A. (2002) Another phocine distemper outbreak in Europe // *Science.* 297:209. DOI: 10.1126/science.1075343 (in Eng.).
- [6] Kennedy S. (1998) Morbillivirus infections in marine mammals // *J Comp Pathol.* 119:201-225. PMID: 9807724 (in Eng.).
- [7] Bengston J.L., Boveng P., Franzen U., Have P., Heide-Jorgenson M.-P., Harkonen T.L. (1991) Antibodies to canine distemper virus in Antarctic seals // *Mar Mamm Sci.* 106:203-212. DOI: 10.1111/j.1748-7692.1991.tb00553.x (in Eng.).
- [8] Domingo M., Ferrer L., Pumarola M., Marco A., Plana J., Kennedy S., et al. (1990) Morbillivirus in dolphins // *Nature.* 348:21. DOI: 10.1038/348021a0 (in Eng.).
- [9] Lipscomb T.P., Kennedy S., Moffett P., Ford B.K. (1994) Morbilliviral disease in an Atlantic bottlenose dolphin (*Tursiops truncatus*) from the Gulf of Mexico // *J Wildl Dis.* 30:572-76. DOI: 10.7589/0090-3558-30.4.572 (in Eng.).
- [10] Markussen N.H., Have P. (1992) Phocine distemper virus infection in harp seals (*Phocagroenlandica*) // *Mar Mamm Sci.* 8:19-26. DOI: 10.1111/j.1748-7692.1992.tb00121x (in Eng.).
- [11] Kennedy S., Kuiken T., Jepson P., Deaville R., Forsyth M., Barrett T., van De Bildt M., Osterhaus A., Eybatov T., Duck C., Kydyrmanov A., Mitrofanov I., Wilson S. (2000) Mass die-off of Caspian seals caused by canine distemper virus // *Emerg Infect Dis.* 6:637-39. DOI: 10.3201/eid0606.000613 (in Eng.).
- [12] Geraci J., St Aubin D., Barker I., Webster R., Hinshaw V., Bean W., Ruhnke H., Prescott J., Early G., Baker A., Madoff S., Schooley R. (1982) Mass mortality of harbor seals: pneumonia associated with influenza A virus // *Science.* 215:1129-31. PMID: 7063847 (in Eng.).
- [13] Hinshaw V.S., Bean W.J., Webster R.G., et al. (1984) Are seals frequently infected with avian influenza viruses? // *J Virol.* 51:863-865. PMID: 6255856 (in Eng.).
- [14] Callan R.G., Early C., Rida H., Hinshaw V.S. (1995) The appearance of H3 influenza viruses in seals // *J Gen Virol.* 76(1):199-203. DOI: 10.1099/0022-1317-76-1-199 (in Eng.).
- [15] Goldstein T., Mena I., Anthony S.J., et al. (2013) Pandemic H1N1 Influenza Isolated from Free-Ranging Northern Elephant Seals in 2010 off the Central California Coast // *PLoS ONE* 8(5): e62259. DOI: 10.1371/journal.pone.0062259 (in Eng.).
- [16] Anthony S.J., St Leger J.A., Puglianes K., Ip H.S., Chan J.M., Carpenter Z.W., Navarrete-Macias I., Sanchez-Leon M., Saliki J.T., Pedersen J., Karesh W., Daszak P., Rabadan R., Rowles T., Lipkin W.I. (2012) Emergence of fatal avian influenza in New England harbor seals // *mBio.* 3(4):e00166-12. DOI: 10.1128/mBio.00166-12 (in Eng.).

- [17]Zohari S., Neimanis A., Härkönen T., Moraes C., Valarcher J.F. (2014) Avian influenza A(H10N7) virus involvement in mass mortality of harbour seals (*Phocavitulina*) in Sweden, March through October 2014. *Euro Surveill.* 19(46):pii=20967.PMID: 25425511 (in Eng.).
- [18]Bodewes R., Bestebroer T.M., van der Vries E., Verhagen J.H., Herfst S., Koopmans M.P., Fouchier R.A., Pfanckuche V.M., Wohlsein P., Siebert U., Baumgärtner W., Osterhaus A.D. (2015) Avian Influenza A(H10N7) virus-associated mass deaths among harbor seals // *Emerg Infect Dis.* 21(4):720-22. DOI: 10.3201/eid2104.141675 (in Eng.).
- [19]Krog J.S., M Hansen M.S., Holm E., Hjulsgager C.K., Chriél M., Pedersen K., Andresen L.O., Abildstrøm M., Jensen T.H., Larsen L.E. (2015) Influenza A(H10N7) virus in dead harbor seals // Denmark. *Emerg Infect Dis.* 21(4):684-87. DOI: 10.3201/eid2104.141484 (in Eng.).
- [20] Lvov D.K., Zhdanov V.M., Sazonov A.A., et al. (1978) Comparison of influenza viruses isolated from man and from whales // *Bull WHO.* 56:923-930. PMID:310734 (in Eng.).
- [21]Hinshaw V.S., Bean W.J., Geraci J.R. (1986) Characterization of two influenza A viruses from a pilot whale // *J Virol.* 58:655-656. PMID:3701925 (in Eng.).
- [22] Shestopalov A.M., Beklemishev A.B., Khuraskin L.S., Ternovoy V.A., Blinov V.M., Netesov S.V. (2002) Para- and orthomyxoviruses in Caspian seals. *Marine Mammals of Holarctic. Abstracts of the conference presentations.* M.: KMK. P. 287-288 (in Eng.).
- [23] Durymanova A.A., Belikov S.I., Zolotykh S.I., Tumanov Yu.V., Kuznetsov V.N., Khuraskin L.S., Dimov S.K., Shestopalov A.M. (2004) Monitoring of infectious diseases in Caspian seals (*Phocacaspica*) // *Marine Mammals of Holarctic. Collection of Scientific Papers.* M.: KMK. P. 200-202. ISBN 5-87317-171-8 (in Eng.).
- [24] Chuvakova Z.K., Ikranbegiyn R., Glebova T.I. (2001) Influenza in seals // *News of the National academy of sciences of the Republic of Kazakhstan. Series of biological and medical.* 3:47-54 (in Rus.).
- [25] Bekmagambetova A.Zh., Ikranbegijn R., Glebova T.I. (2001) Immunofermentnaja diagnostika binarnoj mikrovirusnoj infekcii vovremja massovoj gibeli tjulenej na Severnom Kaspii letom 2000 g. # *Mezhdunar. nauch. konf. Molodyh uchenyh i studentov, posvjashhennaja 10-letiju Nezavisimosti Respubliki Kazakhstan "Aktual'nye voprosy sovremennoj biologii i biotekhnologii"* 25-27 aprelja, 2001. Almaty. P. 159-161 (in Rus.).
- [26] Alekseev A.Y., Gulyaeva M.A., Sivay M.V., Sharshov K.A., Kuznetsov V.A., Shipulin S.V., Shestopalov A.M. (2015) Isolation of influenza A subtype H4N6 in Caspian seals (*Phocacaspica*) // *Marine Mammals of the Holarctic. Collection of Scientific Papers.* 1:27-28. Moscow. ISBN 978-5-9904294-3-7.
- [27]Gulyaeva M., Sobolev I., Sharshov K., Kurskaya O., Alekseev A., Shestopalova L., Kovner A., Bi Y., Shi W., Shchekkanov M., Shestopalov A. (2018) Characterization of Avian-like Influenza A (H4N6) Virus Isolated from Caspian Seal in 2012 // *Virol Sin.* 33(5):449-452. DOI: 10.1007/s12250-018-0053-y (in Eng.).
- [28] Osterhaus A.D.M.E., Rimmelzwaan G.F., Martina, B.E.E. et al. (2000) Influenza B virus in seals // *Science.* 288:1051-53. PMID:10807575 (in Eng.).
- [29]Ohishi K., Ninomiya A., Kida H. et al. (2002) Serological evidence of transmission of human influenza A and B viruses to Caspian seals (*Phocacaspica*) // *Microbiol Immunol.* 46(9):639-44. PMID:12437032 (in Eng.).
- [30] Blanc A., Ruchansky D., Clara M., Achaval F., Le Bas A., Arbiza J. (2009) Serologic evidence of influenza A and B viruses in South American fur seals (*Arctocephalus australis*) // *J Wildl Dis.* 45:519-21. DOI: 10.7589/0090-3558-45.2.519 (in Eng.).
- [31] Goldstein T., Colegrove K.M., Hanson M., Gulland F.M. (2011) Isolation of a novel adenovirus from California sea lions *Zalophus californianus* // *Dis Aquat Organ.* 94:243-248. DOI: 10.3354/dao02321 (in Eng.).
- [32]Inoshima Y., Murakami T., Ishigur, N., Hasegawa K., Kasamatsu M. (2013) An outbreak of lethal adenovirus infection among different otariid species // *Vet Microbiol.* 165:455-459. DOI: 10.1016/j.vetmic.2013.04.013 (in Eng.).
- [33] Smith A.W., Skilling D.E., Benirschke K., Albert T.F., Barlough J.E. (1987) Serology and virology of the bowhead whale (*Balaenamysticetus L.*) // *J Wildl Dis.* 23:92-98. PMID: 3820430 (in Eng.).
- [34] Rubio-Guerri C., García-Párraga D., Nieto-Pelegrín E., Melero M., Álvaro T., Valls M., Crespo J.L., Sánchez-Vizcaino J.M. (2015) Novel adenovirus detected in captive bottlenose dolphins (*Tursiops truncatus*) suffering from self-limiting gastroenteritis // *BMC Vet Res* 11:1. DOI: 10.1186/s12917-015-0367-z (in Eng.).
- [35] Osterhaus A.D., Yang H., Spijkers H.E., Groen J., Teppema J.S., van Steenis, G. (1985) The isolation and partial characterization of a highly pathogenic herpesvirus from the harbor seal (*Phocavitulina*) // *Arch Virol.* 86:239-51. PMID: 4062560 (in Eng.).
- [36] Harder T.C., Harder M., Vos H., Kulonen K., Kennedy-Stoskopf S., Liess B., Appel M.J., Osterhaus A.D. (1996) Characterization of phocid herpesvirus-1 and-2 as putative alpha- and gammaherpesviruses of North American and European pinnipeds // *J Gen Virol.* 77:27-35. DOI: 10.1099/0022-1317-77-1-27 (in Eng.).
- [37] Goldstein T., Gulland F.M.D., Braun R.C., Antonelis G.A., Kashinsky L., Rowles T.K., Mazet J.A.K., Dalton L.M., Aldridge B.M., Stott J.L. (2006) Molecular identification of a novel gamma herpesvirus in the endangered Hawaiian monk seal (*Monachus schauinslandi*) // *Mar. Mamm. Sci.* 22(2):465-471. DOI: 10.1111/j.1748-7692.2006.00025.x (in Eng.).
- [38] Maness H.T.D., Nollens H.H., Jensen E.D., et al. (2011) Phylogenetic analysis of marine mammal herpesviruses // *Vet Microbiol.* 149:23-29. DOI: 10.1016/j.vetmic.2010.09.035 (in Eng.).
- [39] Wright E.P., Waugh L.F., Goldstein T., Freeman K.S., Kelly T.R., Wheeler E.A., Smith B.R., Gulland F.M. Evaluation of viruses and their association with ocular lesions in pinnipeds in rehabilitation // (2015) *Vet Ophthalmol.* Jan; 18 Suppl 1:148-59. DOI: 10.1111/vop.12235. Epub 2014 Nov 17. DOI: 10.1111/vop.12235 (in Eng.).
- [40] Bodewes R., Contreras G.J., Garcia A.R., Hapsari R., van de Bildt, M.W., Kuiken T., Osterhaus AD. (2015) Identification of DNA sequences that imply a novel gammaherpesvirus in seals // *J Gen Virol.* 96:1109-14. DOI: 10.1099/vir.0.000029 (in Eng.).

- [41] Gulland F.M.D., Lowenstine L.J., Lapointe J.M., Spraker T., King D.P. (1997) Herpesvirus infection in stranded Pacific harbor seals of coastal California // *J Wildl Dis.* 33(3):450-458. DOI: 10.7589/0090-3558-33.3.450 (in Eng.).
- [42] Have P., Nielsen J., Botner A. (1991) The seal death in Danish waters 1988. II. Virological studies // *Acta Vet Scand.* 32:211–219. PMID: 1803934 (in Eng.).
- [43] Harder T.C., Plötz P., Liess B. (1991) Detection of antibodies against European seal herpesviruses in sera of Antarctic seals // *Polar Biol.* 34:43-47. DOI: 10.1007/BF00233087 (in Eng.).
- [44] Zarnke R.L., Harder T.C., Vos H.W., VerHoef J.M., Osterhaus A.D. (1997) Serologic survey for phocid herpesvirus-1 and -2 in marine mammals from Alaska and Russia // *J Wildl Dis.* 33:459-465. DOI: 10.7589/0090-3558-33.3.459
- [45] Kydyrmanov A.I., Karamendin K.O., Kasymbekov E.T., Goodman S. (2018) Circulation of Phocine herpesvirus PhHV-1 in the population of Caspian seals (*Pusa caspica*) // Abstracts of the X International Conference "Marine mammals of the Holarctic" dedicated to the memory of A.V. Yablokov, 29 October-2 November 2018, Arkhangel'sk, 216 p. P. 171-172 (in Eng.).
- [46] Bossart G.D., Schwartz J.C. (1990) Acute necrotizing enteritis associated with suspected coronavirus infection in three harbor seals (*Phocavitulina*) // *J Zoo Wildl Med.* 21:84-87 (in Eng.).
- [47] Mihindukulasuriya K.A., G. Wu. J. St. Leger, Nordhausen R.W., Wang D. (2008) Identification of a novel coronavirus from a beluga whale by using a panviral microarray // *J Virol.* 82:5084-88. DOI: 10.1128/JVI.02722-07 (in Eng.).
- [48] Woo P.C., Lau S.K., Lam C.S., et al. (2014) Discovery of a novel bottlenose dolphin coronavirus reveals a distinct species of marine mammal coronavirus in Gammacoronavirus // *J Virol.* 88:1318-1331. DOI: 10.1128/JVI.02351-13 (in Eng.).
- [49] Van Bresse M.F., Van Waerebeek K., Reyes J.C., Dekegel P., Pastoret P.P. (1993) Evidence of poxvirus in dusky dolphin (*Lagenorhynchus obscurus*) and Burmeister's porpoise (*Phocoenaspini pinnis*) from coastal Peru // *J Wildl Dis.* 29:109-13. DOI: 10.7589/0090-3558-29.1.109 (in Eng.).
- [50] Bracht A.J., Brudek, R.L., Ewing R.Y., Manire, C.A., Burek K.A., Rosa C., Beckmen K.B., Maruniak J.E., Romero C.H. (2006) Genetic identification of novel poxviruses of cetaceans and pinnipeds // *Arch Virol.* 151:423-38. DOI: 10.1007/s00705-005-0679-6 (in Eng.).
- [51] Rivera R., Nollens H.H., Venn-Watson S., Gulland F.M., Wellehan J.F.Jr. (2010) Characterization of phylogenetically diverse astroviruses of marine mammals // *J Gen Virol.* 91:166-73. DOI: 10.1099/vir.0.015222-0 (in Eng.).
- [52] Leland D.S., Ginocchio C.C. (2007) Role of cell culture for virus detection in the age of technology // *Clin Microbiol Rev.* 20:49-78. DOI: 10.1128/CMR.00002-06 (in Eng.).
- [53] Yongfeng H., Fan Y., Jie P. et al. (2011) Direct pathogen detection from swab samples using a new high-throughput sequencing technology // *Clin Microbiol Infect.* 17, 241-44. DOI: 10.1111/j.1469-0691.2010.03246.x (in Eng.).
- [54] Karamendin K.O., Kydyrmanov A.I., Kasymbekov E.T., Daulbaeva K.D., Sayatov M.Kh., Fereidouni S. (2018) Application of massive parallel sequencing for the investigation of wild birds viruses // *Bulletin of National academy of sciences of The Republic of Kazakhstan.* 4(374): 13-17. ISSN 2518-1467 (Online). ISSN 1991-3494 (Print) (in Eng.).
- [55] Ng T.F., Suedmeyer W.K., Wheeler E., Gulland F., Breitbart M. (2009) Novel anellovirus discovered from a mortality event of captive California sea lions // *J Gen Virol.* 90:1256-61. DOI: 10.1099/vir.0.008987-0 (in Eng.).
- [56] Ng T.F., Wheeler E., Greig P., Waltzek T.B., Gulland F., Breitbart M. (2011) Metagenomic identification of a novel anellovirus in Pacific harbor seal (*Phocavitulina richardsii*) lung samples and its detection in samples from multiple years // *J Gen Virol.* 92:1318-23. DOI: 10.1099/vir.0.029678-0 (in Eng.).
- [57] Kydyrmanov A.I. (2014) Influenza viruses circulating in marine mammals populations of the Holarctic. – *Biotechnology. Theory and practice.* N 2, June. P. 11-6. DOI: 10.11134/btp.2.2014.2 (in Rus.).
- [58] Kydyrmanov A.I., Karamendin K.O., Kasymbekov E.T., Sayatov M.Kh., Goodman S. (2015) surveillance for circulation of orthomyxo and morbilliviruses among seals in the Kazakh part of Northern Caspian (2007–2014) // *News of the National academy of sciences of the Republic of Kazakhstan Series of biological and medical.* 5(311):41-47 (ISSN 2224-5308) (in Rus.).
- [59] Zhumatov K.Kh., Kydyrmanov A.I., Karamendin K.O., Daulbaeva K.D., Khan Ye.Ya., Kassymbekov Ye.T., Sayatov M.Kh., Fereidouni S. (2018) Phylogenetic analysis of genes of surface proteins of highly pathogenic influenza A subtype H5N1 virus, isolated from gull in Kazakhstan // *Bulletin of National academy of sciences of the Republic of Kazakhstan.* 5(375):6-11. ISSN 2518-1467 (Online), ISSN 1991-3494 (Print) <https://doi.org/10.32014/2018.2518-1467.1> (in Eng.).