

**ISSN 2518-1467 (Online),
ISSN 1991-3494 (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ФЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Ш Ы С Ы

ВЕСТНИК

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

THE BULLETIN

THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944

3

MAY – JUNE 2020

ALMATY, NAS RK

NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources CitationIndex, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.

Б а с р е д а к т о р ы

х.ғ.д., проф., КР ҰҒА академигі
М.Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

Абиев Р.Ш. проф. (Ресей)
Абишев М.Е. проф., корр.-мүшесі (Қазақстан)
Аврамов К.В. проф. (Украина)
Аппель Юрген проф. (Германия)
Баймуқанов Д.А. проф., корр.-мүшесі (Қазақстан)
Байтулин И.О. проф., академик (Қазақстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Қазақстан)
Велесько С. проф. (Германия)
Велихов Е.П. проф., РГА академигі (Ресей)
Гашимзаде Ф. проф., академик (Әзірбайжан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., корр.-мүшесі (Қазақстан)
Джрабашян Р.Т. проф., академик (Армения)
Қалимолдаев М.Н. проф., академик (Қазақстан), бас ред. орынбасары
Лаверов Н.П. проф., академик РАН (Россия)
Лупашку Ф. проф., корр.-мүшесі (Молдова)
Моҳд Ҳасан Селамат проф. (Малайзия)
Мырхалықов Ж.У. проф., академик (Қазақстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., корр.-мүшесі (Қазақстан)
Полещук О.Х. проф. (Ресей)
Поняев А.И. проф. (Ресей)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Қазақстан)
Таткеева Г.Г. проф., корр.-мүшесі (Қазақстан)
Үмбетаев И. проф., академик (Қазақстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., РГА академигі (Ресей)
Якубова М.М. проф., академик (Тәжікстан)

«Қазақстан Республикасы Үлттық ғылым академиясының Хабаршысы».

ISSN 2518-1467 (Online),
ISSN 1991-3494 (Print)

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы»РКБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрагат комитетінде 01.06.2006 ж. берілген №5551-Ж мерзімдік басылым тіркеуіне қойылу туралы қуәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://www.bulletin-science.kz/index.php/en/>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Г л а в н ы й р е д а к т о р

д.х.н., проф. академик НАН РК

М.Ж. Журинов

Р е д а к ц и о н на я кол л е г и я:

Абиев Р.Ш. проф. (Россия)
Абишев М.Е. проф., чл.-корр. (Казахстан)
Аврамов К.В. проф. (Украина)
Аппель Юрген проф. (Германия)
Баймukanов Д.А. проф., чл.-корр. (Казахстан)
Байтулин И.О. проф., академик (Казахстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Казахстан)
Велесько С. проф. (Германия)
Велихов Е.П. проф., академик РАН (Россия)
Гашимзаде Ф. проф., академик (Азербайджан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., чл.-корр. (Казахстан)
Джрабашян Р.Т. проф., академик (Армения)
Калимолдаев М.Н. академик (Казахстан), зам. гл. ред.
Лаверов Н.П. проф., академик РАН (Россия)
Лупашку Ф. проф., чл.-корр. (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Мырхалыков Ж.У. проф., академик (Казахстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., чл.-корр. (Казахстан)
Полещук О.Х. проф. (Россия)
Поняев А.И. проф. (Россия)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Казахстан)
Таткеева Г.Г. проф., чл.-корр. (Казахстан)
Умбетаев И. проф., академик (Казахстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., академик РАН (Россия)
Якубова М.М. проф., академик (Таджикистан)

«Вестник Национальной академии наук Республики Казахстан».

ISSN 2518-1467 (Online),
ISSN 1991-3494 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5551-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 2000 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.
<http://www.bulletin-science.kz/index.php/en/>

© Национальная академия наук Республики Казахстан, 2020

Адрес типографии: «NurNazGRACE», г. Алматы, ул. Рыскулова, 103.

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

M.Zh. Zhurinov

E d i t o r i a l b o a r d:

Abiyev R.Sh. prof. (Russia)

Abishev M.Ye. prof., corr. member (Kazakhstan)

Avramov K.V. prof. (Ukraine)

Appel Jurgen, prof. (Germany)

Baimukanov D.A. prof., corr. member (Kazakhstan)

Baitullin I.O. prof., academician (Kazakhstan)

Joseph Banas, prof. (Poland)

Bersimbayev R.I. prof., academician (Kazakhstan)

Velesco S., prof. (Germany)

Velikhov Ye.P. prof., academician of RAS (Russia)

Gashimzade F. prof., academician (Azerbaijan)

Goncharuk V.V. prof., academician (Ukraine)

Davletov A.Ye. prof., corr. member (Kazakhstan)

Dzhrbashian R.T. prof., academician (Armenia)

Kalimoldayev M.N. prof., academician (Kazakhstan), deputy editor in chief

Laverov N.P. prof., academician of RAS (Russia)

Lupashku F. prof., corr. member (Moldova)

Mohd Hassan Selamat, prof. (Malaysia)

Myrkhalykov Zh.U. prof., academician (Kazakhstan)

Nowak Isabella, prof. (Poland)

Ogar N.P. prof., corr. member (Kazakhstan)

Poleshchuk O.Kh. prof. (Russia)

Ponyaev A.I. prof. (Russia)

Sagyan A.S. prof., academician (Armenia)

Satubaldin S.S. prof., academician (Kazakhstan)

Tatkeyeva G.G. prof., corr. member (Kazakhstan)

Umbetayev I. prof., academician (Kazakhstan)

Khripunov G.S. prof. (Ukraine)

Yuldashbayev Y.A. prof., academician of RAS (Russia)

Yakubova M.M. prof., academician (Tadzhikistan)

Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5551-Ж, issued 01.06.2006.

Periodicity: 6 times a year.

Circulation: 2000 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://www.bulletin-science.kz/index.php/en/>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

**BULLETIN OF NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN**

ISSN 1991-3494

Volume 3, Number 385 (2020), 36 – 42

<https://doi.org/10.32014/2020.2518-1467.67>

UDC 633.913.322

K. R. Uteulin, G. T. Bari, A. Zheksenbai

Republican state enterprise «Institute of Plant Biology and Biotechnology, Committee of Sciences,
Ministry of Education and Science of the Republic of Kazakhstan, Almaty, Kazakhstan.
E-mail: gen_uteulink@mail.ru

**DANDELION KOK-SAGHZ (TARAXACUM KOK-SAGHZ
L. RODIN) AS A ONE-YEAR CULTURE DEVELOPED
UNDER CONDITIONS OF SOUTHEAST KAZAKHSTAN**

Abstract. Dandelion kok-saghyz (*Taraxacum kok-saghyz* L.Rodin) is a herbaceous plant, a source of high-quality rubber. At present, *Taraxacum kok-saghyz* (Tks) is introduced into agriculture in Kazakhstan.

"Saryzhaz" is Kazakh variety of Tks tested as a one-year culture on the experimental plots of the South-Eastern Kazakhstan (N43.1471, E77.2977). The vegetation period of Tks "Saryzhaz" from the preparation of seedlings in the greenhouse (early March) until achene development maturity in the open field takes five months. The studies of kok-saghyz "Saryzhaz" as a one-year culture were repeated three times.

From the 200 m² plots, the raw root harvest was 0.35 (2017), 0.32 (2018), and 0.37 centers (2019), with a density of 10 plants per 1 m². The rubber content is 8 ± 3% per the dry weight of the root. By extrapolating the results of research to one hectare (at a plant density of 100,000), one can predict the harvest of a raw root from 16.0 to 18.5 centers per hectare.

This article demonstrates the possibility of Tks "Saryzhaz" cultivating as a one-year culture in the South-East of Kazakhstan. Tks "Saryzhaz" passes the vegetation cycle before flowering, achene development\maturity in a short spring and early summer period of 5 months.

The combination of the following conditions makes it possible to use Tks as a one-year-old culture in the South-East of Kazakhstan: the use of seedlings, sufficient moisture supply (rains, watering), increased (optimal) temperatures of the spring months for the accelerated plant development.

The increase of the root crop of Tks as a one-year-old crop can be achieved through optimization of farming practices (fertilizers, stocking density) and selection work.

Key words: Kok-saghyz, one-year culture, Kazakhstan.

Introduction. Dandelion kok-saghyz (*Taraxacum kok-saghyz* L.Rodin) is a perennial herbaceous plant, with the habitat in the valleys of the mountains of the South-Eastern Kazakhstan [1].

Taraxacum kok-saghyz (Tks) was introduced into agriculture in the last century, as a technical culture with the roots synthesizing natural rubber in the quality not inferior to the natural rubber of the tropical tree of hevea [2].

Natural Rubber (NR) is a strategically important material, used in production of a wide range of different products: tires (automobile, airplane), medical gloves and other medical products.

At present time, there is a real threat of destruction by the diseases of the largest plantations of the traditional source of rubber tree - hevea (*Hevea brasiliensis*) in the countries of South-East Asia - the main world producers of NK, and therefore a threat to the NK market.

In the light of these developments, EU [3], the US, Canada [4] (PENRA, 2018), and Kazakhstan are taking measures to develop research and introduction of Tks into agriculture as an alternative source of natural rubber.

In Kazakhstan, from 1932 to 1935, on the fields of the Turkestan district, South Kazakhstan, the Tks was cultivated as a one-year plant and harvest about 1 ton of roots per hectare, with the output of pure natural rubber up to 8%.

By the beginning of 1934, 5 871 hectares had been developed for sowing of natural rubber plants. But, as explained in the documents of the Archive of the Republic of Kazakhstan, "due to the financing cut in 1936, the development of industrial natural rubber production in Kazakhstan was suspended" [5].

Tks studies in Kazakhstan have been resumed since 2005 by the Institute of Plant Biology and Biotechnology (IBBR), Almaty [2].

There is a need for Tks genotypes with high root yield, high rubber concentration, and high seed vigor [6].

The purpose of this study was to test the first Kazakh variety Tks "Saryzhaz" as a one-year culture on the experimental plots of the South-East of Kazakhstan [7].

Materials and methods. Tests of Tks "Saryzhaz" were carried out in the open ground in the territory of the agricultural farm "Mirnoe": village "Ostemir": Talgar district, Almaty region, Kazakhstan (N43°37'49", E77°15'41"). The presence of artesian wells provides low-cost water supply, irrigation of farming areas "Ostemir".

The experimental plot of 200 m² combined eight small sections of 25 m² each. The soil of the site is sandy loam, with manure fertilization applied (400 kg per 200 m² plot) to a depth of 15 cm.

The seedling method of cultivation of Tks was used. In a greenhouse, seed sowing was carried out on peat pills, the seedlings developed for two months (March, April) under 25°C [8].

The planting of the Tks "Saryzhaz" seedlings was carried out in the open ground of the experimental plots "Ostemir". Density of planting is 10 plants per square meter. Care of the plantings consisted of the following measures: fertilization (manure), watering, loosening of rows, weeding.

The climate in "Ostemir" is steppe, continental with hot summers and cold winters. The spring and autumn rainy, average air temperatures in April are +18.2° C, in May + 23.9°C, that is optimal for the development of plants Tks.

The hottest months are July, August. Weather with the temperature close to +30°C is observed on average in 36 days a year (Weather in Ostemir, 2017, 2018, 2019) [9].

The collected roots were washed and dried under 80°C till the constant weight was achieved. The method of alkaline extraction of rubber from dry roots was used, followed by gravimetric determination of the rubber content by Koyalovich method [10]. The results were statistically processed [11].

Results and discussion. The life cycle of Tks "Saryzhaz" was documented across the testing period of nine months (March-November) 2017, 2018 and 2019. That is, the studies of kok-saghyz "Saryzhaz" as a one-year culture were repeated three times. The chronology and duration of various stages of Tks "Saryzhaz" growth under the conditions of the village of "Ostemir", Southeast Kazakhstan are shown in figure 1.

Growth stage	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV
Preparation of seedlings. Development seedling.									
Planting seedlings. Adaptation of seedlings to the open soil. Leaf development/rosette growth.									
Inflorescence emerge									
Flowering, achene development\maturity. Peak.									
Summer resting									
Autumn flowering									
Winter resting									

Figure 1 – Chronology and duration of various stages of growth of Tks "Saryzhaz" under the conditions of "Ostemir"

In early May, seedlings with 6-8 real leaves were planted in the open ground of the experimental site (figure 2).



Figure 2 – Experimental plots "Ostemir".
Planting Tks "Saryzhaz" seedlings in April (a, b), the plots in June (c) and July (d)

The period of adapting Tks "Saryzhaz" seedlings to open soil conditions took two weeks - the leaves of the seedlings died off and then new leaves formed, a new rosette.

In early June, the beginning of flowering Tks "Saryzhaz" was marked.

The period of active flowering and achene development\maturity was mainly in July and early August. In early August, the number of leaves in the rosette reached 40-56, the length of the leaf plate 10 cm, the leaf width 1.5 cm.

The height of the aerial part with the peduncle was 22 ± 5 cm, of the total number of plants, non-flowering forms amounted to 49%, flowering forms had from 8 to 26 peduncles. Mature seeds appeared about 8-12 days later. On one peduncle there are 42-56 seeds.

The leaves of Tks "Saryzhaz" are green, in shape a leaf is entire, incised or dissected. The shares of the cut or incised leaf are wide and round, without small burrs. The leaves of the basket wrapper have carob-shaped appendages.

In August, in "Ostemir" even under the conditions of watering, the leaves dry up as Tks "Saryzhaz" has the "summer resting", which lasts until the end of September.

With a decrease in temperature in October (+ 16 °C) and due to rainfalls, Tks "Saryzhaz" leaves come out from the "summer resting".

New leaves grow, an outlet is formed, and then an autumn flowering is observed. After the period of "summer break" in October, the regenerated plants did not have time to recover up to the mass of the first fruiting period before "summer resting", and also 1/3 of the number of plants died.

The growth stages of the Tks "Saryzhaz" are shown in figure 3.



Figure 3 – Growth stages of Tks "Saryzhaz" as one-year culture. 60 -days-old seedlings (a), adaptation of seedlings to open ground (b), rosette growth (c), the beginning of flowering and maturity (d), the peak of flowering and of achene development\maturity (e), summer resting (f), plant regeneration (g), autumn flowering (h)

In the beginning of August, during the period of the fruiting Tks "Saryzhaz" root harvesting was carried out. Illustrations of the samples of the harvest of the roots and rubber of Tks "Saryzhaz" are shown in figure 4.

Tks "Saryzhaz" has a fibrous root system (which does not have the main root). The sternal root was less common (18-22% of the total number of roots).



Figure 4 – Samples of the harvest of Tks "Saryzhaz". Fibrous root types (a), raw root samples (b), dry root samples (c), and rubber samples (d)

Similar results were obtained in the last century in the South of Kazakhstan by Mynbayev K.

When manure is applied (40 tons per ha) to a small depth of up to 15 cm, about 80 percent of the total number of plants form branched roots; when applied the same amount of manure deeper up to 20-25 cm, the branching percentage drops down to 58%.

The more favorable the conditions in the surface layers of the soil, the more branched roots and the higher the yield of the root mass [12].

From the 200 m² plots, the raw root harvest were 0.35 (2017) 0.32 (2018) and 0.37 centers (2019), with a density of 10 plants per 1 m².

The average weight of one raw root of Tks "Saryzhaz" is 16 ± 5 grams, with the rubber content of 8 ± 3% per the dry weight of the root.

By extrapolating the results of research to one hectare (at a plant density of 100,000), one can predict the harvest of the raw root from 16.0 to 18.5 centers per a hectare.

Thus, this article demonstrates the possibility of Tks "Saryzhaz" cultivating as a one-year culture in the South-East of Kazakhstan. Tks "Saryzhaz" passes the vegetation cycle before flowering, achene development\maturity in a short spring and early summer period of 5 months.

The combination of the following conditions makes it possible to use Tks as a one-year-old culture in the South-East of Kazakhstan: the use of seedlings, sufficient moisture supply (rains, watering), increased, optimal temperatures of the spring months for the accelerated plant development.

The increase of the root crop Tks as a one-year-old crop can be achieved through optimization of farming practices (fertilizers, stocking density) and selection work.

Acknowledgments. The author expresses his gratitude for the help in carrying out the investigations to Mr. Murat Akhmetzhanov, the director of the farm "Akhmetzhanov", in the village of Ostemir, Almaty oblast, Kazakhstan. These studies were funded by the Ministry of Science and Education of Kazakhstan and the World Bank.

К. Р. Утеулин, Г. Т. Бари, А. Жексенбай

«Биология және өсімдіктер биотехнологиясы институты» Республикалық мемлекеттік кәсіпорыны, Казақстан Республикасының Білім және ғылым министрлігі, Алматы, Қазақстан

ОНГҮСТІК-ШЫҒЫС ҚАЗАҚСТАН ЖАҒДАЙЫНДАҒЫ КӨК-САҒЫЗ (TARAXACUM KOK-SAGHYZ L. RODIN): БІРЖЫЛДЫҚ ЕКПЕ ӨСІМДІГІ РЕТИНДЕ

Аннотация. Көк-сағыз (*Taraxacum kok-saghyz* L.Rodin) шебі – жоғары сапалы каучук көзі болатын өсімдік. Қазіргі кезде көк-сағыз Қазақстанда егістікке енгізіліп келеді.

Оңтүстік-Шығыс Қазақстан тәжірибе алқаптарында (N43.1471, E77.2977) бір жылдық екпе өсімдік ретінде бірінші қазақстандық «Сарыжаз» атты сұрыптың сынақтары өткен.

«Сарыжаз» көк-сағызының вегетациялық мерзімі жылышайда өскіндерді дайындаудан бастап (наурыз айының басында), тұқым беруге дейін ашық жерде бес айды қамтиды. Көк-сағызды «Сарыжаз» жыл сайынғы мәдениет ретінде зерттеу үш рет қайталанды.

1 м²-де 10 өсімдік тығыздығының 200 м² алқабында дымқос тамырдан 0,35 (2017), 0,32 (2018) және 0,37 центнер (2019) алынған.

Тамырдың орташа құрғақ салмағы 8±3 % каучук мөлшеріне бір тал «Сарыжаз» көк-сағызының дымқос тамырлының 16 ±5 граммына тең.

Бір гектарда (100 000 өсімдік тығыздығында) зерттеу нәтижелерін экспроприациялап, 16,0-дан 18,5 центнерге дейін дымқос тамыр өнімін бір гектардан алушы болжакуға болады.

Сол себепті бұл мақала Оңтүстік-Шығыс Қазақстан аумағында «Сарыжаз» ТКС-ың бір жылдық екпе өсімдік ретінде өсіру мүмкіншілігін көрсетеді. «Сарыжаз» ТКС-ы 5 айлық көктемдегі қысқа және ерте жаздық мерзімде гүлденуге, жетіліп пісуге дейін вегетациялық кезеңінен өтіп жатыр.

Келесі жағдайлардың үйлесуі ТКС-ның бір жылдық екпе өсімдік ретінде қолданылуы: көшеттерді қолдану, жеткілікті ылғалдық (жауын, суару), өсімдіктердің тез дамуы үшін көктем айларының оптимальды температурасы мүмкіндік береді.

Бір жылдық ретіндегі өнімінде ТКС тамырының ұлғаоы, ауыл шаруашылық (тыңайтқыштар, отырғызу тығыздығы) пен селекциялық жұмыстар жүргізу әдістерін үйлесімділеуде жүзеге асады.

Түйін сөздер: көк-сағыз, бір жылдық екпе өсімдік, Оңтүстік-Шығыс Қазақстан.

К. Р. Утеулин, Г. Т. Бари, А. Жексенбай

РГП «Институт биологии и биотехнологии растений» КН МОН РК,
Алматы, Казахстан

ОДУВАНЧИК КОК-САГЫЗ (*TARAXACUM KOK-SAGHYZ L. RODIN*) КАК ОДНОЛЕТНЯЯ КУЛЬТУРА В УСЛОВИЯХ ЮГО-ВОСТОЧНОГО КАЗАХСТАНА

Аннотация. Одуванчик кок-сагыз (*Taraxacum kok-saghyz L.Rodin*) – травянистое растение, источник каучука высокого качества. В настоящее время кок-сагыз вводится в культуру в Казахстане.

На экспериментальных участках Юго-Восточного Казахстана (N43.1471, E77.2977) были проведены испытания первого казахстанского сорта кок-сагыза «Сарыжаз» как одногодичной культуры.

Вегетационный период кок-сагыза "Сарыжаз" от подготовки рассады в оранжерее (начало марта) до плодоношения в открытом грунте занимает пять месяцев. Исследования кок-сагыза «Сарыжаз» как однолетней культуры повторялись трижды.

С участков по 200 м² получены урожаи сырого корня 0,35 (2017) 0,32 (2018) и 0,37 центнеров (2019) при плотности 10 растений на 1 м².

Средний вес одного сырого корня кок-сагыза «Сарыжаз» равен 16±5 грамм с содержанием каучука 8±3 % на сухой вес корня.

Экстраполируя результаты исследований на один гектар (при плотности растений 100 000), можно прогнозировать урожай сырого корня от 16,0 до 18,5 центнеров с одного гектара.

Таким образом, в настоящей работе продемонстрирована возможность возделывания кок-сагыза на территории Юго-Восточного Казахстана как одногодичной культуры. Кок-сагыз «Сарыжаз» проходит вегетационный цикл до цветения, плодоношения за короткий весенний и ранний летний период – 5 месяцев.

Сочетание следующих условий позволяет использовать кок-сагыз как одногодичную культуру на юге Казахстана: использование рассады, достаточное влагообеспечение (дожди, полив), повышенные (оптимальные) температуры весенних месяцев для ускоренного развития растений.

Увеличение урожая корня однолетнего кок-сагыза может быть достигнуто через оптимизацию агротехнических приемов (удобрения, плотность посадки) и селекционную работу.

Ключевые слова: кок-сагыз, однолетняя культура, Казахстан.

Information about authors:

Uteulin Kairat, Head of the department, Associate professor, Institute of Plant Biology and Biotechnology, Committee of Sciences, Ministry of Education and Science of the Republic of Kazakhstan, Almaty, Kazakhstan; gen_uteulink@mail.ru; <https://orcid.org/0000-0002-5459-0902>. First co-author for correspondence.

Bari Gabit, Researcher, Institute of Plant Biology and Biotechnology, Committee of Sciences, Ministry of Education and Science of the Republic of Kazakhstan, Almaty, Kazakhstan; baracuda.co@mail.ru; <https://orcid.org/0000-0003-1987-7315>

Zheksenbay Aslan, Researcher, Institute of Plant Biology and Biotechnology, Committee of Sciences, Ministry of Education and Science of the Republic of Kazakhstan, Almaty, Kazakhstan; aslan.zhexenbay@mail.ru; <https://orcid.org/0000-0002-1101-7153>

REFERENCES

- [1] Bykov B.A. (1981) Red date book of Kazakh SSR. Rare and endangered species of animals and plants. Part 2. Plants. Publishing House "Nauka" of Kazakhstan SSR. Alma-Ata. 1981. 211 p. (in Russ.).
- [2] Volis S., Uteulin K., Mills D. (2009) Russian dandelion (*Taraxacum kok-saghyz*): one more example of over collecting in the past. The Journal of Applied Botany and Food Quality 83: 60–63 (in Eng.).
- [3] Drive4eu (2018). Website: <http://www.drive4eu.eu>
- [4] PENRA (2018). Website: <https://u.osu.edu/penra>
- [5] Central State Archive of the Republic of Kazakhstan [Utesheva Alipa Ves'ma rentabel'noe, legko kul'tiviruemoe... Kazahstanskaja pravda. 2017. 29 June (N 123). Fond arhiva]. Website: vesma-rentabelnoe-legko-kultiviruemoe. (in Russ.).
- [6] Kreuzberger Marie, Hahn Thomas, Zibek Susanne, Schiemann Joachim, Thiele Katja (2016) Seasonal pattern of biomass and rubber and inulin of wild Russian dandelion (*Taraxacum kok-saghyz* L. Rodin) under experimental field conditions // European Journal of Agronomy. 80. 66–77 journal homepage: www.elsevier.com/locate/eja. <https://doi.org/10.1016/j.eja.2016.06.011> (in Eng.).
- [7] Order of Ministry of Agriculture of the Republic of Kazakhstan, number 22 "On Approval of State Register of Selective Achievements, Recommended for Use in RK, and Perspective Agricultural Varieties." Article 5. Technical. Paragraph 5.3 Kok-saghyz *Taraxacum kok-saghyz*, Index Number 717, dated 20.01.2017.[Prikaz Ministerstva sel'skogo hozjajstva Respubliki Kazahstan № 22 «Ob utverzhdenii Gosudarstvennogo reestra selekcionnyh dostizhenij, rekomenduemyh k primeneniju v RK, i perspektivnyh sel'skohozjajstvennyh sortov». Stat'ja 5. Tehnicheskaja. Paragraf 5.3 Kok-saghyz *Taraxacum kok-saghyz*, indeks № 717 ot 20.01.2017] (in Russ.).
- [8] Uteulin K.R., Mukhambetzhanov S.K. (2014) "Method of seed cultivation of *Taraxacum kok-saghyz* Rodin". Number of the innovative patent: 28878. [Uteulin K.R., Muhamabetzhanov S.K. (2014) Sposob semennogo vyrashhivaniya Taraxacum kok-saghyz Rodin. Nomer innovacionnogo patenta: 28878] (in Russ.).
- [9] Pogoda v Ostemir. Website: <https://www.meteonova.ru/frc/-4189> (in Russ.).
- [10] Koyalovich N., Alpatieva M. (1939) Method of simultaneous qualitative and quantitative determination of rubber in the roots of kok-saghyz. Rubber and gum elastic. 7: 205-210. [Kojalovich N, Alpat'eva M. Metod odnovremennogo kachestvennogo i kolichestvennogo opredelenija kauchuka v kornej kok-sagyza. Nauchnyj otchet VIR. M.: OGIZ-Sel'hozgiz; 1945. P. 205-10] (in Russ.).
- [11] Dospechov B.A. (1979) Methodology of field experience Moscow: Kolos: 416. [Dospehov B. A. (1985) Metodika polevogo opyta (s osnovami statisticheskoy obrabotki rezul'tatov issledovanij). M.: Agropromizdat, 351 p.] (in Russ.).
- [12] Mynbayev K. (1946) Kok-saghyz: Biological specifics of the development and new methods of selection. Alma-Ata. 44 p. [Mynbaev K. (1946). Kok-Sagyz. Biologicheskie osobennosti razvitiya i novye metody selekcii. Alma-Ata : KazOGIZ, 148 p.] (in Russ.).

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka–nanrk.kz

ISSN 2518–1467 (Online), ISSN 1991–3494 (Print)

<http://www.bulletin-science.kz/index.php/en/>

Редакторы *M. С. Ахметова, Т. А. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. А. Абдрахимовой*

Подписано в печать 12.06.2020.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
19,1 п.л. Тираж 500. Заказ 3.