

ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Д.В. Сокольский атындағы «Жанармай,
катализ және электрохимия институты» АҚ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
АО «Институт топлива, катализа и
электрохимии им. Д.В. Сокольского»

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
JSC «D.V. Sokolsky institute of fuel, catalysis
and electrochemistry»

SERIES
CHEMISTRY AND TECHNOLOGY

3 (441)

MAY – JUNE 2020

PUBLISHED SINCE JANUARY 1947

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Б а с р е д а к т о р ы
х.ғ.д., проф., ҚР ҰҒА академигі
М.Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

Ағабеков В.Е. проф., академик (Белорус)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Ресей)
Газалиев А.М. проф., академик (Қазақстан)
Ергожин Е.Е. проф., академик (Қазақстан)
Жармағамбетова А.К. проф. (Қазақстан), бас ред. орынбасары
Жоробекова Ш.Ж. проф., академик (Қырғыстан)
Иткулова Ш.С. проф. (Қазақстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Қазақстан)
Баешов А.Б. проф., академик (Қазақстан)
Бүркітбаев М.М. проф., академик (Қазақстан)
Джусипбеков У.Ж. проф., корр.-мүшесі (Қазақстан)
Молдахметов М.З. проф., академик (Қазақстан)
Мансуров З.А. проф. (Қазақстан)
Наурызбаев М.К. проф. (Қазақстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф., академик (Қазақстан)
Стрельцов Е. проф. (Белорус)
Тәшімов Л.Т. проф., академик (Қазақстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Тәжікстан)
Фарзалиев В. проф., академик (Әзірбайжан)

«ҚР ҰҒА Хабарлары. Химия және технология сериясы».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Редакцияның мекенжайы: 050100, Алматы қ., Қонаев к-сі, 142, «Д. В. Сокольский атындағы отын, катализ және электрохимия институты» АҚ, каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор
д.х.н., проф., академик НАН РК
М.Ж. Журинов

Редакционная коллегия:

Агабеков В.Е. проф., академик (Беларусь)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Россия)
Газалиев А.М. проф., академик (Казахстан)
Ергожин Е.Е. проф., академик (Казахстан)
Жармагамбетова А.К. проф. (Казахстан), зам. гл. ред.
Жоробекова Ш.Ж. проф., академик (Кыргызстан)
Иткулова Ш.С. проф. (Казахстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Казахстан)
Баешов А.Б. проф., академик (Казахстан)
Буркитбаев М.М. проф., академик (Казахстан)
Джусипбеков У.Ж. проф., чл.-корр. (Казахстан)
Мулдахметов М.З. проф., академик (Казахстан)
Мансуров З.А. проф. (Казахстан)
Наурызбаев М.К. проф. (Казахстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф., академик (Казахстан)
Стрельцов Е. проф. (Беларусь)
Ташимов Л.Т. проф., академик (Казахстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Таджикистан)
Фарзалиев В. проф., академик (Азербайджан)

«Известия НАН РК. Серия химии и технологии».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел. 272-13-19; 272-13-18,

<http://chemistry-technology.kz/index.php/en/arhiv>

© Национальная академия наук Республики Казахстан, 2020

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142, АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского», каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief

doctor of chemistry, professor, academician of NAS RK

M.Zh. Zhurinov

Editorial board:

Agabekov V.Ye. prof., academician (Belarus)
Volkov S.V. prof., academician (Ukraine)
Vorotyntsev M.A. prof., academician (Russia)
Gazaliyev A.M. prof., academician (Kazakhstan)
Yergozhin Ye.Ye. prof., academician (Kazakhstan)
Zharmagambetova A.K. prof. (Kazakhstan), deputy editor in chief
Zhorobekova Sh.Zh. prof., academician (Kyrgyzstan)
Itkulova Sh.S. prof. (Kazakhstan)
Mantashyan A.A. prof., academician (Armenia)
Praliyev K.D. prof., academician (Kazakhstan)
Bayeshov A.B. prof., academician (Kazakhstan)
Burkitbayev M.M. prof., academician (Kazakhstan)
Dzhusipbekov U.Zh. prof., corr. member (Kazakhstan)
Muldakhmetov M.Z. prof., academician (Kazakhstan)
Mansurov Z.A. prof. (Kazakhstan)
Nauryzbayev M.K. prof. (Kazakhstan)
Rudik V. prof., academician (Moldova)
Rakhimov K.D. prof., academician (Kazakhstan)
Streltsov Ye. prof. (Belarus)
Tashimov L.T. prof., academician (Kazakhstan)
Toderash I. prof., academician (Moldova)
Khalikov D.Kh. prof., academician (Tadjikistan)
Farzaliyev V. prof., academician (Azerbaijan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,

<http://chemistry-technology.kz/index.php/en/arhiv>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Editorial address: JSC «D.V. Sokolsky institute of fuel, catalysis and electrochemistry», 142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22, e-mail: orgcat@nursat.kz

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES CHEMISTRY AND TECHNOLOGY

ISSN 2224-5286

<https://doi.org/10.32014/2020.2518-1491.42>

Volume 3, Number 441 (2020), 44 – 51

UDC 547.99

IRSTI 31.23.23

Y.S. Ikhsanov, A.K. Umbetova, A. T. Amangazieva, G.Sh. Burasheva

Al-Farabi Kazakh National University Research Institute

of New Technologies and Materials Republic of Kazakhstan, Almaty.

E-mail: erbol.ih@gmail.com, alma_0875@mail.ru, amangaziyevaa@gmail.com, gauharbur@mail.ru

**BIOLOGICALLY ACTIVE SUBSTANCES OF COMPOSITIONS
BASED ON PLANTS OF THE GENUS HAPLOPHYLLUM**

Abstract. The article presents the results of research of biologically active substances of compositions based on the plant of the genus *Haplophyllum (Tuberculatum)*, which grows in the Bakanas district of Almaty region, *Bidens (Froncosa)*, collected in the Enbekshikazakh district and *Polygonum (Aviculare)* from the territory of Tashtykara district of Almaty region. The mineral composition was studied, and the article also provides data on vitamin, amino and fatty acid compositions.

Medicinal raw materials contain many compounds that are potentially harmful to the human body, these can be both mineral components, for example, salts of heavy metals, and various organic substances, both being natural metabolites of the plant and trapped in raw materials from the environment (pesticides used in agriculture or industrial emissions).

The study of mineral composition was carried out using atomic absorption analysis on the basis of the center of physico-chemical methods of research and analysis (cfhma). As a result, information was obtained about the micro- and macronutrient composition of compositions based on a plant of the genus *Haplophyllum (Tuberculatum)*.

The content of fatty acids was determined by the GVC method and using the following chromatography conditions: carrier gas-helium, flame ionization detector, carrier gas speed 30ml / min, detector temperature 188°C, furnace temperature 230°C, analysis time 1 hour, steel column 0.4×3mm filled with polyethylene glycoladipinate (20%) on cellulite-545.

Studies of amino acids were carried out on the amino acid analyzer "Carlo Erba", using the gzhx method, chromatography conditions: carrier gas-helium, flame ionization detector 300°C, evaporator temperature 250°C, on the WAW chromosome.

The content of vitamins a (retinol) and (tocopherol) was determined by the fluorimetric method on the spectrofluorimeter device. Vitamin C in biological samples was determined by titrimetric method.

Keywords: *Haplophyllum (Tuberculatum)*, *Bidens (Froncosa)*, *Polygonum (aviculare)*, mineral composition, amino acids, fatty acids.

Introduction. Minerals are of great importance for the normal existence of organisms. They are classified as macro- and microelements. Macronutrients include potassium, sodium, calcium, magnesium, phosphorus, chlorine, and sulfur. Comparatively, they come in large quantities.

Trace elements are a group of chemical elements that are contained in the human body and animals in very small amounts, within the range of 10⁻³-10⁻¹² mg %, these include Nickel, cadmium, lead, chromium, mercury and other d - elements [1].

In ecologically unfavorable areas, a very large number of heavy metals, such as lead, Nickel, chromium, mercury, and any changes in the optimal ratios of trace elements in their composition can lead to unforeseen consequences [2].

In an acidic environment, the movement of molybdenum decreases, but the movement of copper, manganese, zinc, and cobalt increases. Trace elements such as chalk, fluorine, and iodine also move in acidic and alkaline environments. Some trace elements, for example, form compounds that are soluble with organic matter, while others (iodine and copper) are fixed and become inaccessible to plants. The

lack or excess of trace elements in the soil leads to their deficiency or excess in the plant and animal body. At the same time, changes in the nature of accumulation (deposition), weakening or strengthening the synthesis of biologically active substances, repeated violation of the processes of interline exchange, creating a new adaptation or violations that lead to endemic diseases of humans and animals develop [3].

Thus, the mineral composition of plants is directly affected by natural and anthropogenic factors. All this should be taken into account when preparing vegetable raw materials.

Currently, there are 14 trace elements necessary for life: iron, copper, manganese, zinc, cobalt, iodine, fluorine, molybdenum, vanadium, Nickel, strontium, silicon and selenium. They increase the activity of enzymes, catalyze biochemical procedures, and promote the synthesis of carbohydrates, proteins, and vitamins. It also participates in metabolism. Trace elements are part of plant preparations and influence their activity[4].

Most studies of *Haplophyllum tuberculatum* evaluated medicinal and phytochemical properties, analysis of some of which produced two alkaloids called alkaloids, lignans, glycosides and flavonoids, etc. in the phytochemical study of *Haplophyllum tuberculatum*, alkaloids such as acutifolium, halilofitin were obtained [5]. The chemical composition of *Polygonum aviculare* is rich and varied. It is a natural storehouse of vegetable protein (17 %), biologically active substances (44%), vegetable fiber (27%), ash compounds (8.9%), simple and complex sugars (2.5%), resin, wax, and tannins. Useful trace elements such as calcium, phosphorus, zinc, and silicon were also found in it [6]. *Bidens frondosa* contains organic acids, essential oil, tannins, polysaccharides, coumarins (umbelliferon, skoletin), triterpenoids, vitamin C (up to 0.9%), carotenoids (0.05 %), carotenes, a large amount of vitamin C (up to 1000 mg%), manganese salts, polyacetylenes, and aromatic acids containing derivatives and thiophenes, flavonoids - (glycosides luteolin, Butin, sulfuretin, Auron) [7-12].

Fatty acids are the basis of metabolic processes in the body, an integral part of the biological synthesis of the body's processes along with the components of lipids. Several fatty acids that have essential properties have vitamin-like effectiveness: olein, arachidone, linol, linolene. Essential fatty acids have cardioprotective properties, improving blood circulation and the cardiovascular system.

Amino acids are organic compounds whose molecules simultaneously contain carboxyl and amine groups. These are substances of primary synthesis, they are present in all organs of all plants. Amino acids are divided into α , β , γ , σ , and others. amino acids, depending on the location of the amino and carboxyl groups. Of these, the most common are α , β and γ . α - l-configuration amino acids are the most important components of peptides and proteins. Also, plants may contain monobasic diamino-and dibasic monoaminoacids.

Vitamin A and E occur in pairs. They improve the condition of the skin, vision, and bones, helping to assimilate each other, increase resistance to infections, and protect the mucous membrane. Retinol is one of the vital vitamins necessary and vital for our body, which belong to the fat-soluble group. The usefulness of vitamin A for health is invaluable: it participates in redox processes, affects protein synthesis, and cell membranes. Retinol interacts well with another fat-soluble vitamin-tocopherol (vitamin E). if the body lacks vitamin E, the absorption of retinol is absorbed, so it is optimal to take these vitamins together.

Ascorbic acid is an organic compound similar to glucose, the sour taste is found in the form of a white crystalline powder. It performs the biological functions of a coenzyme and a reducing agent of certain metabolic processes, and is an antioxidant. Vitamin C strengthens the immune system and protects it from viruses and bacteria, accelerates the healing process, promotes the synthesis of several hormones, regulates the processes of the circulatory system and normalizes capillary permeability, participates in the synthesis of collagen protein. This is necessary for the growth of tissue cells, bones and cartilage of the body, regulates metabolism, improves bile secretion, and restores the external function of the pancreas and thyroid gland.

Materials and methods. *Determination of mineral composition.* The objects of the study were plants of the genus *Haplophyllum (Tuberculatum)* growing in the Bakanas district of Almaty region, *Bidens (Frondosa)* collected in the Enbekshikazakh district and *Polygonum (aviculare)* from the territory of Tashtykara district of Almaty region, harvesting was carried out in compliance with sanitary requirements and the requirements Of the state Pharmacopoeia of the Republic of Kazakhstan. The quantitative content of micro-and macronutrients was determined from ash residues obtained by the following method.

About 1 g of the drug or 3-5 g of crushed medicinal plant raw materials (exact weight) is placed in a pre-calcined and precisely weighted porcelain, quartz or platinum crucible, evenly distributing the substance along the bottom of the crucible.

Then the crucible is carefully heated, allowing the substance to burn or evaporate at the lowest possible temperature. The burning of the remaining coal particles must also be carried out at a lower temperature as possible; after the coal has burned almost completely, the flame is increased. In case of incomplete combustion of coal particles, the residue is cooled, moistened with water or a saturated solution of ammonium nitrate, evaporated in a water bath and the residue is calcined. If necessary, repeat this operation several times.

Calcination is carried out at low red heat (about 500°C) to a constant mass, avoiding fusing the ash and sintering with the walls of the crucible. At the end of calcination, the crucible is cooled in a desiccator and then the resulting ash is burned again at 600°C until a uniform gray color is obtained.

If the result is not achieved, the remainder is dissolved in concentrated nitric acid, after which it is heated on a tile removing the nitric acid and then in a muffle at 400°C for 30 minutes.

Finally, the precipitate is dissolved in 5 ml of HNO₃ (1:1) when heated. The resulting solution must be heated on the tile to wet salts. The result is dissolved in 10-15 ml of 1N HCl or 1N HNO₃ (the Second option is preferable) and transferred to a 25ml volumetric flask, bringing the volume to the label.

In parallel, a single experiment is carried out, which consists in preparing a solution of the same concentration from the same acid using the same utensils.

Then the finished samples were transferred to the center for physical and chemical methods of research and analysis to determine the mineral composition by atomic adsorption spectroscopy on the ASSIN device of the Karl Zeiss company.

The results obtained are presented below [13,14].

It was revealed that the amount of heavy metals does not exceed the permissible norms of their presence in medicinal raw materials.

Study of the fatty acid composition of the obtained extracts by the GLC method. The content of fatty acids was determined using the GLC method using the following chromatography conditions: carrier gas-helium; flame ionization detector; carrier gas speed 30ml / min; detector temperature 188°C; furnace temperature 230°C; analysis time 1H; steel column 0.4 m × 3mm filled with polyethylene glycoladipinate (20%) on cellulite-545.

Sample preparation: 10ml of methanol, 2-3 drops of acetyl chloride are added To the chloroform extracts of the test samples, and then methylation is performed at 60-70°C in a special system for 30 minutes. Methanol is removed using a rotary evaporator, and samples are extracted with 5ml of hexane and analyzed in a Carlo-Erbo-4200 gas chromatograph[15].

Study of the amino acid composition of the obtained extracts by the GLC method. 1 g of the analyte is hydrolyzed in 5ml of 6N hydrochloric acid at 105°C for 24 hours, in ampoules sealed under a jet of argon. The resulting hydrolysate is evaporated three times to dry on a rotary evaporator at a temperature of 40-50°C. The resulting precipitate is dissolved in 5ml of sulfosalicylic acid. After centrifugation for 5 minutes, the supernatant is passed through a column with an ion-exchange resin Daux 50, with a speed of 1 drop per second. After that, the resin is washed to a neutral pH.

For elution of amino acids from the column, 3ml of 6N NH₄OH co solution is passed through it at a rate of 2 drops per second. The eluate is collected in a round-bottomed flask with distilled water, which is used to wash the column to a neutral pH. Then the contents of the flask are evaporated dry on a rotary evaporator at a pressure of 1 atm. and a temperature of 40-50°C.

After adding to this flask, 1 drop of freshly prepared 1.5 % SnCl₂ solution, 1 drop of 2,2-dimethoxypropane and 1-2ml of hydrochloric acid-saturated propanol, ee is heated to 110 °C, maintaining this temperature, for 20 minutes, and then the contents are again evaporated from the flask on a rotary evaporator.

In the next step, 1 ml of freshly prepared acetylizing reagent (1 volume of acetic anhydride, 2 volumes of triethylamine, 5 volumes of acetone) is injected into the flask and heated at a temperature of 60 °C for 1.5-2min. Then the sample is again evaporated on a rotary evaporator to dry and 2ml of ethyl acetate and 1ml of saturated NaCl solution is added to the flask. The contents of the flask are thoroughly mixed and, as 2 layers of liquids are clearly formed, the upper one (ethyl acetate) is taken for gas chromatographic analysis, which was performed on the gas-liquid chromatograph "Carlo-Erba-4200" (Italy-USA) [16].

Study of vitamins A, E, and C. 0.2 ml of the sample, 1ml of alcohol and 1ml of distilled water are placed in centrifuge tubes, tightly closed with lids and mixed with careful shaking. Add 5ml of hexane and shake again. The contents are centrifuged for 10 minutes at 1500rpm. The Separated hexane layer (3 ml) is used for measurements at wavelengths of 292-310 nm (for tocopherol) and 335-430 nm (for retinol), respectively [17-22].

The content of vitamin C in biological samples was determined by titrimetric method. A sample of at least 0.3 ml is taken in a centrifuge tube, the walls of which are covered with sodium citrate powder. After centrifuging the sample for 30 minutes at 3000 rpm, it is transferred to another test tube and an equal amount of bidistilled water and double the amount of freshly prepared 5% metaphosphoric acid solution is added. The protein precipitate is stirred with a stick and centrifuged for 10 minutes at 3000 rpm. The nadosadochnuyu liquid in an amount (0.1-0.5 ml) is introduced into porcelain titration cuvettes (2 parallel samples) and titrated 0.001 n-0.0005 n with a solution of the sodium salt of 2.6 dichlorophenolindophenol from a special micropipette with a capacity of 0.1 ml.

Results and discussion. To create complexes, first of all, the composite correspondence of plant raw materials to each other was studied and the ratio was chosen. Depending on which ones show up, they were obtained in three types. The percentage of components in the complexes is theoretically justified by taking into account the contribution of each component to the last therapeutic effect and evaluating the activity of plant components. This takes into account the phenomenon of interaction of plant components (interference), since some types of plant raw materials show the greatest activity in an individual or a mixture, and when it changes, there is a decrease in the influence of the participating components. The plant complex was obtained in different ratios. The results are presented in table 1.

Table 1 - plant ratio

| № | complex |
|---|---------|
| 1 | 1:1:3 |
| 2 | 2:1:1 |
| 3 | 2:3:1 |

Based on the data from table 1, the mineral composition of plant complexes was studied in 3 different ratios.

Data on the mineral composition were obtained by atomic absorption analysis conducted at the material and technical base of the center for physical and chemical analysis methods. The results are presented in tables 2 and 3.

Table 2 - Quantitative content of trace elements

| Element | Zn | Cu | Pb | Cd | Fe | Ni | Mn |
|--------------------------|-----------|-----------|------------|------------|---------|------------|----------|
| 1:1:3 | | | | | | | |
| Content in the sample, % | 0.0004892 | 0.0002358 | 0.00009821 | 0.00001683 | 0.01304 | 0.00006447 | 0.001113 |
| 2:1:1 | | | | | | | |
| Content in the sample, % | 0.0006301 | 0.0002323 | 0.00006677 | 0.00001795 | 0.01558 | 0.00006005 | 0.001059 |
| 2:3:1 | | | | | | | |
| Content in the sample, % | 0.0007055 | 0.0004068 | 0.0001103 | 0.00002102 | 0.01162 | 0.00008746 | 0.001415 |

Table 3 - Quantitative content of macronutrients

| Element | K | Na | Ca | Mg |
|--------------------------|--------|---------|--------|---------|
| 1:1:3 | | | | |
| Content in the sample, % | 0.6040 | 0.01851 | 0.4576 | 0.07897 |
| 2:1:1 | | | | |
| Content in the sample, % | 0.5823 | 0.02051 | 0.3448 | 0.08191 |
| 2:3:1 | | | | |
| Content in the sample, % | 0.6166 | 0.02578 | 0.3786 | 0.06799 |

Comparing the data given in table 2, we can conclude that the quantitative content of iron dominates all samples of plant complexes in 3 different ratios. Iron is a part of herbal preparations and affects their activity. It is also known that iron increases the activity of enzymes and catalyzes biochemical processes. Promotes the synthesis of carbohydrates, proteins and vitamins.

Among the macro elements presented in table 3 and figure 2, the largest number of elements are K, Ca. They are part of the cell nucleus. Calcium plays an important role in the growth and action of cells. It has a significant impact on the exchange process and contributes to the full digestion of food substances. By strengthening the body's protective function, it increases resistance to external adverse conditions, especially infections.

The content of fatty acids was determined by the GZHC method. The results are presented in table 4.

Table 4 - Quantitative analysis of fatty acids

| № | Fatty acids | | % |
|---|-------------------|---|------|
| 1 | C _{14:0} | Miristin (C ₁₄ H ₂₈ O ₂) | 0,8 |
| 2 | C _{15:0} | Pentadecan (C ₁₆ H ₃₀ O ₂) | 1,2 |
| 3 | C _{16:0} | Palmitin (C ₁₆ H ₃₂ O ₂) | 4,6 |
| 4 | C _{16:1} | Palmitalein (C ₁₆ H ₃₀ O ₂) | 0,4 |
| 5 | C _{18:0} | Stearin (C ₁₈ H ₃₆ O ₂) | 2,3 |
| 6 | C _{18:1} | Olein (C ₁₈ H ₃₄ O ₂) | 36,5 |
| 7 | C _{18:2} | Linol (C ₁₈ H ₃₂ O ₂) | 53,9 |
| 8 | C _{18:3} | Linolene (C ₁₈ H ₃₀ O ₂) | 0,3 |

From the data obtained, it follows that 8 fatty acids were detected. In terms of quantitative content, the dominant fatty acids were oleic and linoleic acids, relatively palmitic and stearic acids. Linolenic acid, which showed the lowest index, plays a large role in the body. Fatty acid, most necessary for the functioning of cell and subcellular membranes.

Amino acids were identified from the plant complex. Studies of amino acids were carried out on the amino acid analyzer "Carlo Erba", using the gzhx method. The data is presented in table 5.

Table 5 - Quantitative analysis of the amino acid composition of aqueous extracts of compositions

| | Name of amino acids | | Amino acids, mg / 100 g |
|----|---------------------|---------------|-------------------------|
| 1 | Ala | Alanine | 612 |
| 2 | Gly | Glycine | 234 |
| 3 | Val | Valine. | 195 |
| 4 | Leu | Leucine. | 370 |
| 5 | Ile | Isoleucine | 334 |
| 6 | Thr | Threonine | 177 |
| 7 | Ser | Serine | 317 |
| 8 | Pro | Proline | 415 |
| 9 | Met | Methionine | 51 |
| 10 | Asp | Asparate | 1360 |
| 11 | Cys | Cystine | 22 |
| 12 | O-Prp | Oxyproline | 1 |
| 13 | Phe | Phenylalanine | 264 |
| 14 | Glu | Glutamate | 2546 |
| 15 | Orn | Ornithine | 1 |
| 16 | Tyr | Tyrosine | 288 |
| 17 | His | Arginine | 370 |
| 18 | Arg | Lysine | 318 |
| 19 | Lys | Histidine | 189 |
| 20 | Trp | Tryptophan | 72 |

The analysis revealed their qualitative identity and insignificant quantitative relationships, and also found that the dominant amino acids are aspartate, glutamate, alanine: 1360 mg / 100g, 2546 mg/100g, 612 mg / 100g.

The content of vitamins a (retinol) and (tocopherol) was determined by the fluorimetric method on the spectrofluorimeter device . Vitamin C in biological samples was determined by titrimetric method. The data is presented in table 6.

Table 6 - Quantitative content of vitamins A, E and C

| Vitamins | mg/100g |
|----------|---------|
| A | 0,09 |
| E | 2,8 |
| C | 22 |

Comparing the data shown in table 6 and figure 2, we can conclude that the quantitative content of vitamin C dominates.

Conclusion

1. For the first time, biologically active substances were studied, including mineral composition, vitamins, amino and fatty acids of compositions based on a plant of the genus *Haplophyllum* (*Tuberculatum*) growing in the Bakanas district of Almaty region, *Bidens* (*Froncosa*) collected in the Enbekshikazakh district and *Polygonum* (*aviculare*) from the territory of the Tashtykara district of Almaty region.

2. It was found that the level of heavy metal content in the studied samples does not exceed the maximum permissible norm.

Е.С. Ихсанов, А.К. Умбетова, А.Т. Амангазиева, Г.Ш. Бурашева

әл-Фараби атындағы Қазақ ұлттық университеті,

Жаңа технологиялар мен материалдар ғылыми-зерттеу институты, Қазақстан, Алматы

НАРЛОПХУЛЛУМ ТЕКТЕС ӨСІМДІКТЕР НЕГІЗІНДЕГІ КОМПОЗИЦИЯЛАРДЫҢ БИОЛОГИЯЛЫҚ БЕЛСЕНДІ ЗАТТАРЫ

Аннотация. Мақалада Алматы облысының Бақанас ауданында өсетін *Haplophyllum* (*Tuberculatum*) тұясының өсімдігі және Еңбекшіқазақ ауданында жиналған *Bidens* (*Froncosa*) және Алматы облысының Таштықара ауданының аумағынан *Polygonum* (*aviculare*) өсімдітер негізінде композициялардың биологиялық белсенді заттарын зерттеу нәтижелері келтіріледі. Минералды құрам зерттелді, мақалада витаминдік, амин және майлы қышқылдық құрам туралы мәліметтер келтірілген.

Дәрілік шикізат құрамында адам ағзасына зиян келтіретін көптеген қосылыстар бар, олар минералды компоненттер болуы мүмкін, мысалы, ауыр металдардың тұздары, сондай-ақ әр түрлі органикалық заттар, олар өсімдіктердің табиғи метаболиттері болып табылады және қоршаған ортадан шикізатқа түседі (қолданылатын пестицидтер) ауыл шаруашылығында немесе өнеркәсіптік шығарындыларда).

Минералдық құрамды зерттеу физика - химиялық зерттеу және талдау әдістері орталығы (ЦФХМА) базасында атомдық - абсорбциялық талдау әдісімен жүргізілді. Нәтижесінде *haplophyllum* (*Tuberculatum*) тұяқымының негізіндегі композициялардың микро - және макроэлементтік құрамы туралы мәліметтер алынды.

Май қышқылдарының құрамын ГЖХ әдісімен және хроматографиялаудың келесі шарттарын қолдану арқылы анықтадық: газ-тасығыш – гелий, жалынды-иондаушы детектор, газ тасығыштың жылдамдығы 30мл/мин, детектордың температурасы 188°С, пештің температурасы 230°С, талдау уақыты 1с, Болат бағанасы 0.4×3мм, целлит-545 полиэтиленгликольадипинатпен толтырылған (20%).

Аминқышқылдарын зерттеу "Карло Эрба" аминқышқылды талдағышта ГЖХ әдісімен жүргізілді, хроматографиялау шарттары: газ-тасығыш – гелий, 3000с жалынды-иондау детекторы, булау температурасы 250°С, WAW хромосорбасында жүргізілді.

А (ретинол) және Е (токоферол) витаминдері спектрофлуориметр аспабында флуориметрлік әдіспен анықталған. С витамині биологиялық үлгілерде титриметриялық әдіспен анықталды.

Түйін сөздер: *Haplophyllum* (*Tuberculatum*), *Bidens* (*Froncosa*), *Polygonum* (*aviculare*), минералды құрамы, амин қышқылдары, майлы қышқылдар.

Е.С. Ихсанов, А.К. Умбетова, А.Т. Амангазиева, Г.Ш. Бурашева

Казахский национальный университет имени аль-Фараби,
Научно-исследовательский институт новых технологий и материалов
Республика Казахстан, Алматы

БИОЛОГИЧЕСКИ АКТИВНЫЕ ВЕЩЕСТВА КОМПОЗИЦИЙ НА ОСНОВЕ РАСТЕНИЯ РОДА HAPLOPHYLLUM

Аннотация. В статье приводятся результаты исследования биологически активных веществ композиций на основе растения рода *Haplophyllum* (*Tuberculatum*), произрастающего в Баканасском районе Алматинской области, *Bidens* (*Froncosa*), собранный в Енбекшиказахском районе и *Polygonum* (*aviculare*) из территории Таштыкараского района Алматинской области. Были исследованы минеральный состав, в статье также приведены данные по витаминному, аминок- и жирнокислотному составам.

В лекарственном сырье содержится множество соединений, потенциально способных нанести вред организму человека, это могут быть как минеральные компоненты, например, соли тяжёлых металлов, так и различные органические вещества, как являющиеся естественными метаболитами растения, так и попавшие в сырьё из окружающей среды (пестициды, применяемые в сельском хозяйстве, или выбросы промышленных предприятий).

Исследования минерального состава проводились методом атомно- абсорбционного анализа на базе центра физико- химических методов исследования и анализа (ЦФХМА). В результате получены сведения о микро- и макроэлементном составе композиций на основе растения рода *Haplophyllum* (*Tuberculatum*).

Содержание жирных кислот определяли методом ГЖХ и использованием следующих условий хроматографирования: газ-носитель – гелий, пламенно-ионизационный детектор, скорость газо-носителя 30мл/мин, температура детектора 188⁰С, температура печи 230⁰С, время анализа 1ч, стальная колонка 0.4×3мм, заполненная полиэтиленгликольадипинатом (20%) на целлите-545.

Исследования аминокислот проводили на аминокислотном анализаторе «Карло Эрба» методом ГЖХ, условия хроматографирования: газ-носитель – гелий, пламенно-ионизационный детектор 300⁰С, температура испарителя 250⁰С, на хромосорбе WAW.

Содержание витаминов А (ретинол) и (токоферол) определили флуориметрическим методом на приборе спектрофлуориметр. Витамин С в биологических образцах определили титрометрическим методом.

Ключевые слова: *Haplophyllum* (*Tuberculatum*), *Bidens* (*Froncosa*), *Polygonum* (*aviculare*), минеральный состав, аминокислоты, жирные кислоты.

Information about authors:

Ikhsanov Yerbol Saginovich, PhD Department of Chemistry and Chemical Technology of the Al-Farabi Kazakh National University, e-mail: erbol.ih@gmail.com, <https://orcid.org/0000-0003-4640-9584>;

Umbetova Almagul Kengebaeva, Candidate of chemical Sciences, faculty of chemistry and chemical technology, al-Farabi Kazakh national University, alma_0875@mail.ru, <https://orcid.org/0000-0001-9879-5398>;

Amangazieva Asem Talgatovna, Al-Farabi Kazakh National University, Faculty of Chemistry and Chemical Technology, MSc student of 2-nd course, amangaziyevaa@gail.com, <https://orcid.org/0000-0002-0448-9849>;

Burasheva Gauhar Shahmanovna, Doctor of Science professor, Department of Chemistry and Chemical Technology of the Al-Farabi Kazakh National University, gauharbur@mail.ru, <https://orcid.org/0000-0003-2935-3531>

REFERENCES

[1] Zengin G., Sarikurkcü C., Aktumsek A., Ceylan R., Ceylan, O. (2014). A comprehensive study on phytochemical characterization of *Haplophyllum myrtifolium* Boiss. endemic to Turkey and its inhibitory potential against key enzymes involved in Alzheimer, skin diseases and type II diabetes. *Industrial Crops and Products*, 53, 244–251. doi:10.1016/j.indcrop.2013.12.043

[2] Oscanov B. S., Ikhsanov Y. S., Litvinenko Yu. A., Adekenov S. M., Burasheva G. Sh. Biologically active substances from plant *Suaeda vera* and their anestizing activity. *News of the national academy of sciences of the republic of kazakhstan series of biological and medical* ISSN 2224-5308. Volume 5, Number 329 (2018), 63–66. <https://doi.org/10.32014/2018.2518-1629.8>

[3] Aitkulova R.E., Abubakirova A.A., Kudasova D.E., Kaldybekova G.M. (2016) Role of medicinal plants from south-kazakhstan region for addition into livestock's fodder. *News of the national academy of sciences of the republic of kazakhstan series of biological and medical* 314: 155–158. <https://doi.org/10.32014/2018.2518-1629> (in Eng).

[4] Aitkulova R.E., Abubakirova A.A., Kudasova D.E., Kaldybekova G.M. (2016) Role of medicinal plants from south-kazakhstan region for addition into livestock's fodder. *News of the national academy of sciences of the republic of kazakhstan series of biological and medical* 314: 155 – 158. <https://doi.org/10.32014/2018.2518-1629> (in Eng).

[5] Hamdi A., Majouli K., Vander Heyden Y., Flamini G., & Marzouk Z. (2017). Phytotoxic activities of essential oils and hydrosols of *Haplophyllum tuberculatum*. *Industrial Crops and Products*, 97, 440–447. doi:10.1016/j.indcrop.2016.12.053

- [6] Al-Rehaily A.J., Alqasoumi S.I., Yusufoglu H.S., Al-Yahya M.A., Demirci B., Tabanca N., Baser K. H. C. (2014). Chemical Composition and Biological Activity of *Haplophyllum tuberculatum* Juss. Essential Oil. *Journal of Essential Oil Bearing Plants*, 17(3), 452–459. doi:10.1080/0972060x.2014.895211
- [7] Eissa T.F., González-Burgos E., Carretero M.E., Gómez-Serranillos M.P. (2013). Biological activity of HPLC-characterized ethanol extract from the aerial parts of *Haplophyllum tuberculatum*. *Pharmaceutical Biology*, 52(2), 151–156. doi:10.3109/13880209.2013.819517
- [8] Tekin M., Eruygur N. (2016). The structural studies on the medicinal plant *Haplophyllum telephioides*. *Revista Brasileira de Farmacognosia*, 26(5), 544–552. doi:10.1016/j.bjp.2016.05.007
- [9] Debouba M., Khemakhem B., Zouari S., Meskine A., Gouia H. (2014). Chemical and Biological Activities of *Haplophyllum tuberculatum* Organic Extracts and Essential Oil. *Journal of Essential Oil Bearing Plants*, 17(5), 787–796. doi:10.1080/0972060x.2014.958545
- [10] Hamdi A., Viane J., Mahjoub M. A., Majouli K., Gad M. H. H., Kharbach M., Heyden Y.V. (2018). Polyphenolic contents, antioxidant activities and UPLC–ESI–MS analysis of *Haplophyllum tuberculatum* A. Juss leaves extracts. *International Journal of Biological Macromolecules*, 106, 1071–1079. doi:10.1016/j.ijbiomac.2017.08.107
- [11] Rasulova K. A., Kodirova D. R., Bobakulov K. M., Abdullaev N. D. (2015). Griffithine, a New Furanquinolone Alkaloid from *Haplophyllum griffithianum*. *Chemistry of Natural Compounds*, 51(4), 743–745. doi:10.1007/s10600-015-1398-1
- [12] Azadi B., Khaef S., Ziarati P. (2014). Chemical composition of *Haplophyllum villosum* (M. B.) G. Don Essential Oil. *Journal of Essential Oil Bearing Plants*, 17(6), 1161–1164. doi:10.1080/0972060x.2014.958556
- [13] Ulukuş D., Tugay O., Celep F. (2016). Morphology, Anatomy and Palynology of Turkish endemic species *Haplophyllum myrtifolium*, *H. vulcanicum* & *H. megalanthum* (Rutaceae) and their systematics implications. *Phytotaxa*, 247(3), 197. doi:10.11646/phytotaxa.247.3.3
- [14] Cao J., Pang X., Guo S., Wang Y., Geng Z., Sang Y., Du S. (2019). Pinene-rich essential oils from *Haplophyllum dauricum* (L.) G. Don display anti-insect activity on two stored-product insects. *International Biodeterioration & Biodegradation*, 140, 1–8. doi:10.1016/j.ibiod.2019.03.007
- [15] Hamdi A., Majouli K., Abdelhamid A., Marzouk B., Belghith H., Chraief I., Heyden Y. V. (2018). Pharmacological activities of the organic extracts and fatty acid composition of the petroleum ether extract from *Haplophyllum tuberculatum* leaves. *Journal of Ethnopharmacology*, 216, 97–103. doi:10.1016/j.jep.2018.01.012
- [16] Azadi B., & Khaef S. (2015). Volatile constituents of *Haplophyllum buhsei* Boiss. flowering aerial parts. *Bulletin of the Chemical Society of Ethiopia*, 29(2), 327. doi:10.4314/bcse.v29i2.15
- [17] Aberrane S., Djouahri A., Djerrad Z., Saka B., Benseradj F., Aitmoussa S., Boudarene L. (2018). Changes in essential oil composition of *Haplophyllum tuberculatum* (Forssk.) A. Juss. aerial parts according to the developmental stage of growth and incidence on the biological activities. *Journal of Essential Oil Research*, 1–21. doi:10.1080/10412905.2018.1511483
- [18] Rasulova K. A., Bobakulov K. M., Eshonov M. A., Abdullaev N. D. (2019). Pedicine, a New Alkaloid from *Haplophyllum pedicellatum*. *Chemistry of Natural Compounds*. doi:10.1007/s10600-019-02784-7
- [19] Kodirova D. R., Rasulova K. A., Sagdullaev S. S., & Aisa H. A. (2018). *Haplophyllum griffithianum* as a Source of Quinoline Alkaloids. *Chemistry of Natural Compounds*, 54(1), 213–214. doi:10.1007/s10600-018-2303-5
- [20] Abdelgaleil S. A. M., Saad M. M. G., Ariefa N. R., Shiono Y. (2020). Antimicrobial and phytotoxic activities of secondary metabolites from *Haplophyllum tuberculatum* and *Chrysanthemum coronarium*. *South African Journal of Botany*, 128, 35–41. doi:10.1016/j.sajb.2019.10.005
- [21] Khan B., Ahmad H., Alam J., Nadeem M. S., Ud-Din A., Ullah I., Muhammad K. (2019). DNA barcoding of *Haplophyllum gilesii* (Hemsel.) C. C. Townsend from the Gilgit-Baltistan Province of Pakistan. *Brazilian Journal of Botany*. doi:10.1007/s40415-019-00568-5
- [22] Mamarozikov U. B., Bobakulov K. M., Turaeva S. M., Zakirova R. P., Rakhmatov K. A., Abdullaev N. D., Khidyrova, N. K. (2019). Constituent Composition of the Hexane Fraction of the Extract of *Haplophyllum perforatum* and Its Insecticidal Activity. *Chemistry of Natural Compounds*. doi:10.1007/s10600-019-02746-z

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации
в журнале смотреть на сайте:

www.nauka-nanrk.kz

<http://chemistry-technology.kz/index.php/en/arhiv>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 08.06.2020.

Формат 60x88¹/₈. Бумага офсетная. Печать – ризограф.
10,8 п.л. Тираж 300. Заказ 3.