

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Әль-фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Республикасының
Ғылым Академиясының
Әль-Фараби атындағы
Қазақ ұлттық университетінің

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-farabi kazakh
national university

SERIES
PHYSICO-MATHEMATICAL

5 (327)

SEPTEMBER-OCTOBER 2019

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.
Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жангаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф., чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d :

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.60>

Volume 5, Number 327 (2019), 70 – 77

УДК 517.9: 515.16

МРНТИ 27.31.21

A.A. Zhadyranova

Eurasian International Center for Theoretical Physics and Department
of General & Theoretical Physics, Eurasian National University, Astana 010008, Kazakhstan.
a.a.zhadyranova@gmail.com

**HIERARCHY OF WDVV ASSOCIATIVITY EQUATIONS
FOR $n = 3$ AND $N = 2$ CASE WHEN $V_0 = 0$ WITH NEW SYSTEM a_t, b_t, c_t**

Abstract. We investigate solutions of Witten-Dijkgraaf-E.Verlinde-H.Verlinde (WDVV) equations. The article discusses nonlinear equations of the third order for a function $f = f(x,t)$ of two independent variables x,t . The equations of associativity reduce to the nonlinear equations of the third order for a function $f = f(x,t)$ when prepotential F dependet of the metric η . In this work we consider the WDVV equation for $n = 3$ case with an antidiagonal metric η . The solution of some cases of hierarchy equations of associativity illustrated. Lax pairs for the system of three equations, that contains the equation of associativity are written to find the hierarchy of associativity equation. Using the compatibility condition are found the relations between the matrices U, V_2, V_1 . The elements of matrix V_2 are found with the expression of z_{ij} and independent and dependent variables for the matrix V_2 . Also solving elements of matrix V_1 expressed through y_{ij} and independent and dependent variables for the matrix V_1 . We accepted that elements of matrix V_0 are zero. In the physical setting the solutions of WDVV describe moduli space of topological conformal field theories [1, 2]. Let us introduce new variables a, b, c . In the above variables the nonlinear equations of the third order for a function $f = f(x,t)$ we rewritten as a new system of three equations. Expressed are variables a_t, b_t, c_t of three equations are written with the help of matrix elements z_{ij}, y_{ij} .

Key words: equations of Witten-Dijkgraaf-E.Verlinde-H.Verlinde, the equations of associativity, nonlinear equations of the third order, antidiagonal metric, the Lax pair, the compatibility condition, independent elements, dependent variables, system with equations.

Introduction. The WDVV equations, in general, have the following form [1, 2, 3]:

$$\frac{\partial^3 F}{\partial t^i \partial t^j \partial t^p} \eta^{pq} \frac{\partial^3 F}{\partial t^q \partial t^k \partial t^r} = \frac{\partial^3 F}{\partial t^j \partial t^k \partial t^p} \eta^{pq} \frac{\partial^3 F}{\partial t^i \partial t^q \partial t^r}, \quad \forall i, j, k, r \in \{1, \dots, n\},$$

where F is a prepotential, η is a metric. The coordinates t^i can be linearly rearranged so that the metric, η , is antidiagonal [1], i.e.

$$\eta = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

In this work we consider the WDVV equation for $n = 3$ case with an antidiagonal metric η [2]. In this case, two types of dependence of the function F on the fixed variable t^1 were found by Dubrovin [2, 3, 4] which are

$$F = \frac{1}{2}(t^1)^2 t^3 + \frac{1}{2} t^1 (t^2)^2 + f(t^2, t^3) \tag{1}$$

and

$$F = \frac{1}{6}(t^1)^3 + t^1 t^2 t^3 + f(t^2, t^3).$$

For these cases the equations of associativity reduce to the following two nonlinear equations of the third order for a function $f = f(x, t)$ of two independent variables ($x = t^2, t = t^3$):

$$f_{ttt} = f_{xxt}^2 - f_{xxx} f_{xtt} \quad (2)$$

and

$$f_{xxx} f_{ttt} - f_{xxt} f_{xtt} = 1,$$

correspondingly.

The function F in equation (1) has the form from the law of multiplication in the three-dimensional algebra A_1 with the basis $e_1 = 1, e_2, e_3$ [3]. Every basis is a complete uniformly minimal system [2].

In this work, we consider the solution (1). Let us introduce new variables a, b, c as follows [2, 3]:

$$a = f_{xxx}, \quad b = f_{xxt}, \quad c = f_{xtt}.$$

In the above variables the equation (2) can be rewritten as a system of three equations as follows:

$$\begin{aligned} a_t &= b_x, \\ b_t &= c_x, \\ c_t &= (b^2 - ac)_x. \end{aligned} \quad (3)$$

The Lax pair for the system (3) is given by [8]

$$\begin{aligned} \Psi_x &= \lambda U \Psi, \\ \Psi_t &= \lambda V \Psi, \end{aligned} \quad (4)$$

where U is given by

$$U = \begin{pmatrix} 0 & 1 & 0 \\ b & a & 1 \\ c & b & 0 \end{pmatrix}$$

and V is given by

$$V = \begin{pmatrix} 0 & 0 & 1 \\ c & b & 0 \\ (b^2 - ac) & c & 0 \end{pmatrix}.$$

The compatibility condition for the system (4) is given by

$$\begin{aligned} U_t &= V_x, \\ [U, V] &= 0. \end{aligned}$$

In the following sections we work with the new system (3).

Methods. The solution to a hierarchy for $N = 1$ case corresponds to the system of equations (3). Hierarchy for $N = 2$ case when $V_0 \neq 0$ is given in the work [2]

In this section we consider a hierarchy for $N = 2$ case when $V_0 = 0$ and the following system

$$\begin{aligned} a_t &= \varepsilon_1 b_x + \varepsilon_2 F_x, \\ b_t &= \varepsilon_1 c_x + \varepsilon_2 H_x, \\ c_t &= \varepsilon_1 (b^2 - ac)_x + \varepsilon_2 G_x. \end{aligned} \quad (5)$$

The Lax representation of the above system is same as before in the work [13].

In particular, for $N = 2$ case when $V_0 = 0$ we have

$$\begin{aligned} \Psi_x &= \lambda U \Psi, \\ \Psi_t &= (\lambda^2 V_2 + \lambda V_1) \Psi = V \Psi \end{aligned}$$

The compatibility condition of (4) is given by

$$\lambda U_t - V_x + \lambda[U, V] = 0.$$

The compatibility condition of the Lax representation is given by the system

$$[U, V_2] = 0, \quad (6)$$

$$U_t = V_{1x}, \quad (7)$$

$$V_{2x} = [U, V_1] \quad (8)$$

Statement of problem. We first consider the second equation of the system and let V_1 to be given by

$$V_1 = \begin{pmatrix} y_{11} & y_{12} & y_{13} \\ y_{21} & y_{22} & y_{23} \\ y_{31} & y_{32} & y_{33} \end{pmatrix}.$$

From the above system it follows that $y_{11}, y_{12}, y_{13}, y_{23}, y_{33}$ are constants w.r.t. x . Writing a system with equations for a_t, b_t, c_t only yields

$$\begin{aligned} a_t &= y_{22x}, \\ b_t &= y_{21x}, \\ b_t &= y_{32x}, \\ c_t &= y_{31x}. \end{aligned} \quad (9)$$

Now we equate similar terms in the systems (5) and (9), i.e. we have a system

$$\begin{aligned} a_t &= y_{22x} = \varepsilon_1 b_x + \varepsilon_2 F_x, \\ b_t &= y_{21x} = \varepsilon_1 c_x + \varepsilon_2 H_x, \\ b_t &= y_{32x} = \varepsilon_1 c_x + \varepsilon_2 H_x, \\ c_t &= y_{31x} = \varepsilon_1 (b^2 - ac)_x + \varepsilon_2 G_x. \end{aligned} \quad (10)$$

Scheme of the method and reduction to equivalent problem. From the above system (10) we find the following

$$\begin{aligned}
 y_{22} &= \varepsilon_1 b + \varepsilon_2 F, \\
 y_{21} &= \varepsilon_1 c + \varepsilon_2 H, \\
 y_{32} &= \varepsilon_1 c + \varepsilon_2 H, \\
 y_{31} &= \varepsilon_1 (b^2 - ac) + \varepsilon_2 G.
 \end{aligned}$$

Thus the matrix V_1 has the form

$$V_1 = \begin{pmatrix} y_{11} & y_{12} & y_{13} \\ \varepsilon_1 c + \varepsilon_2 H & \varepsilon_1 b + \varepsilon_2 F & y_{23} \\ \varepsilon_1 (b^2 - ac) + \varepsilon_2 G & \varepsilon_1 c + \varepsilon_2 H & y_{33} \end{pmatrix}. \quad (11)$$

Now we solve the equation (6). Denote V_2 as follows:

$$V_2 = \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ z_{21} & z_{22} & z_{23} \\ z_{31} & z_{32} & z_{33} \end{pmatrix},$$

Plugging U , V_2 into (6) we obtain the following relations:

$$\begin{aligned}
 z_{23} &= z_{12}, \\
 z_{32} &= z_{21}, \\
 z_{33} &= z_{11}.
 \end{aligned}$$

Hence, we are left with the equations

$$\begin{aligned}
 z_{21} &= bz_{12} + cz_{13}, \\
 z_{22} &= z_{11} + az_{12} + bz_{13}, \\
 z_{31} &= cz_{12} + (b^2 - ac)z_{13}.
 \end{aligned}$$

Thus the matrix V_2 has the form

$$V_2 = \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ bz_{12} + cz_{13} & z_{11} + az_{12} + bz_{13} & z_{12} \\ cz_{12} + (b^2 - ac)z_{13} & bz_{12} + cz_{13} & z_{11} \end{pmatrix}.$$

Hence, only z_{11}, z_{12}, z_{13} are independent elements of V_2 , and the other elements can be written in terms of them.

Now let us find the elements of V_1 in (11). To do so we use the equation (8). First we evaluate $[U, V_1]$.

We have elementwise yields the following system:

$$\begin{aligned}
 11: \quad z_{11x} &= \varepsilon_1 c + \varepsilon_2 H - by_{12} - cy_{13}, \\
 12: \quad z_{12x} &= \varepsilon_1 b + \varepsilon_2 F - y_{11} - ay_{12} - by_{13}, \\
 13: \quad z_{13x} &= y_{23} - y_{12}, \\
 21: \quad b_x z_{12} + bz_{12x} + c_x z_{13} + cz_{13x} &= by_{11} + a(\varepsilon_1 c + \varepsilon_2 H) + (\varepsilon_1(b^2 - ac) + \varepsilon_2 G) - b(\varepsilon_1 b + \varepsilon_2 F) - cy_{23}, \\
 22: \quad z_{11x} + a_x z_{12} + az_{12x} + b_x z_{13} + bz_{13x} &= by_{12} - by_{23}, \\
 23: \quad z_{12x} &= by_{13} + ay_{23} + y_{33} - (\varepsilon_1 b + \varepsilon_2 F), \\
 31: \quad c_x z_{12} + cz_{12x} + (b^2 - ac)_x z_{13} + (b^2 - ac)z_{13x} &= cy_{11} - cy_{33}, \\
 32: \quad b_x z_{12} + bz_{12x} + c_x z_{13} + cz_{13x} &= cy_{12} + b(\varepsilon_1 b + \varepsilon_2 F) - (\varepsilon_1(b^2 - ac) + \varepsilon_2 G) - a(\varepsilon_1 c + \varepsilon_2 H) - by_{33}, \\
 33: \quad z_{11x} &= cy_{13} + by_{23} - (\varepsilon_1 c + \varepsilon_2 H).
 \end{aligned}$$

Now let us express $\varepsilon_1 c + \varepsilon_2 H$, $\varepsilon_1 b + \varepsilon_2 F$, y_{23} in the element 11, 12, 13 of the above system.

$$\begin{aligned}
 \varepsilon_1 c + \varepsilon_2 H &= z_{11x} + by_{12} + cy_{13}, \\
 \varepsilon_1 b + \varepsilon_2 F &= z_{12x} + y_{11} + ay_{12} + by_{13}, \\
 y_{23} &= z_{13x} + y_{12}.
 \end{aligned}$$

Now let us express $\varepsilon_1(b^2 - ac) + \varepsilon_2 G$ in the element 21 and substitute the values for $\varepsilon_1 c + \varepsilon_2 H$, $\varepsilon_1 b + \varepsilon_2 F$, y_{23}

$$\varepsilon_1(b^2 - ac) + \varepsilon_2 G = b_x z_{12} + bz_{12x} + c_x z_{13} + 2cz_{13x} - az_{11x} + bz_{12x} + (b^2 - ac)y_{13} + cy_{12}$$

Now let us express y_{33} in the element 23 and substitute the values for $\varepsilon_1 b + \varepsilon_2 F$, y_{23}

$$y_{33} = 2z_{12x} - az_{13x} + y_{11}$$

Hence, dependent elements of V_1 are given by:

$$\begin{aligned}
 \varepsilon_1(b^2 - ac) + \varepsilon_2 G &= b_x z_{12} + bz_{12x} + c_x z_{13} + 2cz_{13x} - az_{11x} + bz_{12x} + (b^2 - ac)y_{13} + cy_{12}, \\
 \varepsilon_1 c + \varepsilon_2 H &= z_{11x} + by_{12} + cy_{13}, \\
 \varepsilon_1 b + \varepsilon_2 F &= z_{12x} + y_{11} + ay_{12} + by_{13}, \\
 y_{23} &= z_{13x} + y_{12}, \\
 y_{33} &= 2z_{12x} - az_{13x} + y_{11}.
 \end{aligned} \tag{12}$$

Now let us rewrite the element 22 by substituting the values for y_{23} . So we have

$$z_{11x} + 2bz_{13x} + a_x z_{12} + az_{12x} + b_x z_{13} = 0$$

Now let us rewrite the element 31 by substituting the values for y_{33} . So we have

$$c_x z_{12} + 3cz_{12x} + (b^2 - ac)_x z_{13} + (b^2 - 2ac)z_{13x} = 0$$

Now let us rewrite the element 32 by substituting the values for $\varepsilon_1 b + \varepsilon_2 F$, $\varepsilon_1(b^2 - ac) + \varepsilon_2 G$, $\varepsilon_1 c + \varepsilon_2 H$, y_{33} . So we have

$$2b_x z_{12} + 4bz_{12x} + 2c_x z_{13} + (3c - ab)z_{13x} = 0$$

Now let us rewrite the element 33 by substituting the values for y_{23} , $\varepsilon_1 c + \varepsilon_2 H$

$$2z_{11x} - bz_{13x} = 0$$

Also, the independent variables z_{11}, z_{12}, z_{13} of the matrix V_2 have to satisfy the following system of equations:

$$\begin{aligned} z_{11x} + 2bz_{13x} + a_x z_{12} + az_{12x} + b_x z_{13} &= 0, \\ c_x z_{12} + 3cz_{12x} + (b^2 - ac)_x z_{13} + (b^2 - 2ac)z_{13x} &= 0, \\ 2b_x z_{12} + 4bz_{12x} + 2c_x z_{13} + (3c - ab)z_{13x} &= 0, \\ 2z_{11x} - bz_{13x} &= 0. \end{aligned} \tag{13}$$

From the above system (13) it follows that

$$z_{13x} = \left(\frac{4a_x b - 2ab_x}{3ac - a^2 b - 10b^2} \right) z_{12} + \left(\frac{4bb_x - 2ac_x}{3ac - a^2 b - 10b^2} \right) z_{13} \tag{14}$$

$$\begin{aligned} z_{12x} &= \left(-\frac{c_x}{3c} - \frac{b^2 - 2ac}{3c} \cdot \frac{4a_x b - 2ab_x}{3ac - a^2 b - 10b^2} \right) z_{12} + \\ &+ \left(-\frac{b^2 - 2ac}{3c} \cdot \frac{4bb_x - 2ac_x}{3ac - a^2 b - 10b^2} - \frac{(b^2 - ac)_x}{3c} \right) z_{13} \end{aligned} \tag{15}$$

Results. Using necessary terms in the system (12) in (10), we obtain

$$\begin{aligned} a_t &= \frac{a_x z_{13x}}{2} + a_x y_{12} + b_x y_{13}, \\ b_t &= \frac{b_x z_{13x}}{2} + b_x y_{12} + c_x y_{13}, \\ c_t &= b_{xx} z_{12} + 3b_x z_{12x} + c_{xx} z_{13} + \left(a_x b + 3c_x - \frac{ab_x}{2} \right) z_{13x} - a_x z_{11x} + (b^2 - ac)_x y_{13} + c_x y_{12} \end{aligned} \tag{16}$$

We plug $z_{11x}, z_{12x}, z_{13x}$ in (13), (15), (14) into (16) and obtain the following equation

$$\begin{aligned} a_t &= \left(\frac{2ba_x^2 - aa_x b_x}{3ac - a^2 b - 10b^2} \right) z_{12} + \left(\frac{2ba_x b_x - aa_x c_x}{3ac - a^2 b - 10b^2} \right) z_{13} + a_x y_{12} + b_x y_{13}, \\ b_t &= \left(\frac{2ba_x b_x - ab_x^2}{3ac - a^2 b - 10b^2} \right) z_{12} + \left(\frac{2bb_x^2 - ab_x c_x}{3ac - a^2 b - 10b^2} \right) z_{13} + b_x y_{12} + c_x y_{13}, \\ c_t &= \left(b_{xx} - \frac{b_x c_x}{c} + \frac{5abcab_x - 4b^3 a_x b_x + 2ab^2 b_x^2 - 3a^2 c b_x^2 + 2b^2 c a_x^2 + 12bcac_x - 6ac b_x c_x}{3ac^2 - a^2 bc - 10b^2 c} \right) z_{12} \\ &+ \left(c_{xx} - \frac{b_x (b^2 - ac)_x}{c} + \frac{6abc b_x^2 - 4b^3 b_x^2 + 2ab^2 b_x c_x - 3a^2 c b_x c_x + 2b^2 c a_x b_x + 12bc b_x c_x - abc a_x c_x - 6ac c_x^2}{3ac^2 - a^2 bc - 10b^2 c} \right) z_{13} \\ &+ (b^2 - ac)_x y_{13} + c_x y_{12} \end{aligned} \tag{17}$$

Conclusion. The solution to a hierarchy for $N = 2$ case when system is given by (5) corresponds to the system of equations (17).

So, we considered of some cases of hierarchy of WDVV associativity equations. Lax pairs for the system of three equations, that contained the equation of associativity written to find the hierarchy of associativity equation. Using the compatibility condition are found the relations between the matrices U , V_2 , V_1 . Thus, we obtained the elements of the matrices V_2 , V_1 for case $N = 2$ when $V_0=0$ and the above system a_t, b_t, c_t . It was found, that only z_{11}, z_{12}, z_{13} are independent elements of V_2 , and the other elements can be written in terms of them. From the above system it follows that $y_{11}, y_{12}, y_{13}, y_{23}, y_{33}$ are constants w.r.t. X . It is found, that y_{11}, y_{12}, y_{13} are independent elements of V_1 , and the other elements can be written in terms of them and z_{11}, z_{12}, z_{13} . Expressed are variables a_t, b_t, c_t of three equations are written with the help of matrix elements z_{ij}, y_{ij} .

Acknowledgments. I express gratitude to Professor R. Myrzakulov for useful discussions and advices. The work is performed under the financial support of the scientific and technical program BR05236277 "Investigation of some problems of astrophysics and cosmology in the framework of the Einstein and non-Einstein theories of gravity", 2018.

УДК 517.9: 515.16
МРНТИ 27.31.21

А.А. Жадыранова

Кафедра общей и теоретической физики Евразийского
национального университета имени Л.Н.Гумилева, Астана, Казахстан

**ИЕРАРХИЯ УРАВНЕНИЙ АССОЦИАТИВНОСТИ WDVV
ДЛЯ СЛУЧАЯ $n = 3$ И $N = 2$ ПРИ $V_0 = 0$ С НОВОЙ СИСТЕМОЙ a_t, b_t, c_t**

Аннотация. В данной статье исследуются уравнения Виттена-Диджкграфа-Е.Верлинде-Г.Верлинде (ВДВВ). В работе обсуждаются нелинейные уравнения третьего порядка для функции $f = f(x,t)$ двух независимых переменных x, t . Уравнения ассоциативности сводятся к нелинейным уравнениям третьего порядка для функции $f = f(x,t)$, когда потенциал функции F связан с метрикой η . В этой работе рассматривается уравнение WDVV для случая $n = 3$ с антидиагональной метрикой η . Описано решение некоторых случаев иерархии уравнений ассоциативности. Для нахождения иерархии уравнений ассоциативности были записаны пары Лакса для системы из трех уравнений, которая содержит уравнения ассоциативности. С применением условия совместности найдены соотношения между матрицами U , V_2 , V_1 . Были вычислены элементы матрицы V_2 , выраженные через z_{ij} , независимые и зависимые переменные матрицы V_2 . Также были найдены элементы матрицы V_1 , выраженные через y_{ij} , независимые и зависимые переменные матрицы V_1 . Элементы матрицы V_0 равны 0. В физическом приложении решение уравнения ассоциативности WDVV описывает пространство модулей топологических конформных теорий поля. Введены новые переменные a, b, c . В новых переменных нелинейные уравнения третьего порядка для функции $f = f(x,t)$ записаны через новую систему трёх уравнений. Выраженные переменные a_t, b_t, c_t системы из трех уравнений были записаны через матричные элементы z_{ij}, y_{ij} .

Ключевые слова: уравнения Виттена-Диджкграфа-Е.Верлинде-Г.Верлинде, уравнения ассоциативности, нелинейные уравнения третьего порядка, антидиагональная метрика, пары Лакса, условие совместности, независимые элементы, зависимые переменные, система с уравнениями.

УДК 517.9: 515.16
МРНТИ 27.31.21

А.А. Жадыранова

Л.Н.Гумилев атындағы Еуразия ұлттық университетінің
жалпы және теориялық физика кафедрасы, Астана, Қазақстан

**$n = 3$ ЖӘНЕ $N = 2$ ЖАҒДАЙЛАРЫ ҮШІН ЕНГІЗГІЗІЛГЕН ЖАҢА ЖҮЙЕ a_t, b_t, c_t $V_0 = 0$ БОЛҒАНДАҒЫ
WDVV АССОЦИАТИВТІЛІК ТЕНДЕУІНІҢ ИЕРАРХИЯСЫ**

Аннотация. Берілген мақалада Виттен – Диджкграф - Е.Верлинде - Г.Верлинде (ВДВВ) теңдеулері зерттеледі. Бұл жұмыста x, t тәуелсіз айнымалыларынан тұратын $f = f(x,t)$ функциясы үшін үшінші ретті сызықты емес теңдеулер талқыланады. Тәуелсіз x, t айнымалыларынан тұратын $f = f(x,t)$ функциясы үшін үшінші ретті сызықты емес теңдеулер F потенциалы η метрикасымен байланысты болғанда келтіріледі. Сонымен қатар ассоциативтілік теңдеулер иерархиясының бірнеше шешімдері сипатталады. Ассоциативтілік теңдеулерінің иерархиясын табу мақсатында ассоциативтілік теңдеулерінен құралған теңдеулер жүйесі үшін Лакс жұптары жазылды. Сәйкестік шартының қолдану арқылы U , V_2 , V_1 матрицалары арасындағы қатынастар анықталды. z_{ij} арқылы өрнектелген V_2 матрицасының элементтері мен V_2 матрицасының тәуелді және тәуелсіз айнымалылары

есептелінді. u_{ij} арқылы өрнектелген V_1 матрицасының элементтері мен V_1 матрицасының тәуелді және тәуелсіз айнымалылары табылды. Сонымен қатар V_0 матрицасының элементтері нөлге тең деп алынды. Физикалық қолданылуда WDVV ассоциативтілік теңдеуінің шешімі өрістің топологиялық конформдық теориясының модульдерінің кеңістігін сипаттайды. Жаңа айнымалылар енгізілген. Жаңа айнымалыларда $f = f(x,t)$ функциясы үшін үшінші ретті сызықты емес теңдеулер жаңа жүйе арқылы жазылған. Теңдеулер жүйесінен тұратын a_i, b_i, c_i айнымалылары z_{ij}, u_{ij} матрицалық элементтері арқылы өрнектеліп жазылды.

Түйін сөздер: Виттен-Диджграф-Е.Верлинде-Г.Верлинде теңдеулері, ассоциативтілік теңдеуі, үшінші ретті сызықты емес теңдеулер, антидиагональ метрика, Лакс жұптары, үйлесімділік шарты, тәуелсіз элементтер, тәуелді айнымалылар, теңдеулер жүйесі.

Information about authors:

Zhadyranova A.A. - PhD student of the department of general and theoretical physics, L.N. Gumilyov Eurasian National University, Satpayev str., Astana, Kazakhstan. E-mail: a.a.zhadyranova@gmail.com

REFERENCES

- [1] Kontsevich M., Manin Yu., Gromov-Witten classes, quantum cohomology and enumerative geometry, Mirror symmetry, II, 607-653, AMS/IP Stud. Adv. Math., 1, Amer. Math. Soc., Providence, RI, (1997).
- [2] Kontsevich M., Manin Yu., Relations between the correlators of the topological sigma-model coupled to gravity, Comm. Math. Phys., 196 (1998), no. 2, 385-398.
- [3] Dubrovin B.A. Geometry of 2D topological field theories, Springer Lecture Notes in Math. 1620, 120-348, 1996. [arXiv:hep-th/9407018]
- [4] Dijkgraaf R., Verlinde E., Verlinde H., Notes on topological string theory and 2D quantum gravity, Nucl. Phys. B 352 (1991) 59.
- [5] Witten E., On the structure of the topological phase of two-dimensional gravity, Nucl. Phys. B 340 (1990) 281-332.
- [6] Hertling C., Frobenius manifolds and moduli spaces for singularities, Cambridge University Press, Cambridge (UK), 2002.
- [7] Hertling C., Manin Y., Weak Frobenius manifolds, Internat. Math. Res. Notices, no. 6 (1999), 277-286.
- [8] Mokhov O.I., Ferapontov Y.V. Equations of Associativity in Two-Dimensional Topological Field Theory as Integrable Hamiltonian Nondiagonalizable Systems of Hydrodynamic Type, Functional analysis and its applications 30(3), 1995. [arXiv:hep-th/9505180]
- [9] Dubrovin B.A., Novikov S.P., Hydrodynamics of weakly deformed soliton lattices. Differential geometry and Hamiltonian theory, Uspekhi Mat.Nauk. 44 (1989), 29-98. English translation in Russ. Math. Surveys 44 (1989), 35-124.
- [10] Dubrovin B.A., On almost duality for Frobenius manifolds, Amer. Math. Soc. Transl. 212 (2004) 75-132.
- [11] A.Sh. Shaldanbayev, A.A. Shaldanbayeva, B.A. Shaldanbay (2019) On projectional orthogonal basis of a linear non-self -adjoint operator. News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series. Volume 2, Number 324 (2019), PP, 79-89. ISSN 2518-1726 (Online), ISSN 1991-346X (Print). <https://doi.org/10.32014/2019.2518-1726.15>
- [12] Mokhov O.I. Symplectic and poisson geometry on loop spaces of manifolds and nonlinear equations, Translations of the American Mathematical Society-Series 2 170, 121-152, 1995. [arXiv:hep-th/9503076]
- [13] Ferapontov E.V., Mokhov O.I., Nonlocal Hamiltonian operators of hydrodynamic type that are connected with metrics of constant curvature, Russ. Math. Surv. 45 (1990), no. 3, 218-219.
- [14] Zhadyranova A.A., Myrzakul Zh.R., Anuarbekova Y.Ye. Hierarchy of WDVV associativity equations for $n = 3$ case and $N = 2$ when $V_0 \neq 0$ [Ierarchiya WDVV uravneniya dlya $n = 3$ i $N = 2$ sluchaya, kogda $V_0 \neq 0$] Bulletin of L.N. Gumilyov Eurasian National University [Vestnik Evrazijskogo nacional'nogo universiteta imeni L.N. Gumileva]. 4(125), 60-66, 2018.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 10.10.2019.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
9,6 п.л. Тираж 300. Заказ 5.