

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
 PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2019.2518-1726.38>

Volume 4, Number 326 (2019), 5 – 13

**D. M. Zazulin^{1,2}, N. Burtebayev^{1,2}, R. J. Peterson³, S. V. Artemov⁴,
 S. Igamov⁴, Zh. K. Kerimkulov¹, D. K. Alimov¹, E. S. Mukhamejanov^{1,2},
 Maulen Nassurlla^{1,2}, A. Sabidolda¹, Marzhan Nassurlla^{1,2}, R. Khojaye^{1,2}**

¹Institute of Nuclear Physics, Almaty, Kazakhstan;²Al-Farabi Kazakh National University, Almaty, Kazakhstan;³University of Colorado, Boulder, Colorado, USA;⁴Institute of Nuclear Physics, Tashkent, Uzbekistan

nburtebayev@yandex.ru; denis_zazulin@mail.ru; nburtebayev@yandex.ru; jerry.peterson@colorado.edu;
artemov1943@gmail.com; igamov@inp.uz; zhambul-k@yandex.ru; diliyo@mail.ru;
craftinho@mail.ru; morzhic@mail.ru; nespad@mail.ru; asabidolda@mail.ru; ramazan_inp@mail.ru

NEW RESULTS FOR THE P-¹²C RADIATIVE CAPTURE AT LOW ENERGIES

Abstract. New measurements of the differential cross sections of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ radiative capture reaction at the angle 0° for the transition to the ground state in ^{13}N have been made at the energies of incident protons from 1088 to 1390 keV (uncertainties of about 12%). Based on the obtained differential cross sections and on the assumption that the angular distributions are isotropic in this energy region, the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factors have been determined for the transition to the ground state in ^{13}N with an uncertainties of about 16%. Within the limits of uncertainties, our experimental results are consistent with the previous data. Analysis of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factor at low energies has been carried out within the modified R-matrix approach by using previously measured asymptotic normalization coefficient of the overlap integral of the wave functions of ^{12}C and ^{13}N nuclei bound states to minimize the uncertainties due to calculation of the direct capture part of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factor at extremely low energies. For the energies of 0.25 and 50 keV in center of mass frame the calculated values of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction S-factors for the transition to the ground state in ^{13}N are presented. In the temperature range from 0 to 10^{10} K the rates of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ fusion reaction have been obtained. The results of our calculations are compared with the experimental and calculated results of previous works.

Keywords: differential cross sections, total cross sections, astrophysical S-factor, asymptotic normalization coefficient, reaction rates.

Introduction

It is well known that, in addition to the hydrogen pp chain, in stars more massive than the Sun, hydrogen can burn in the reactions of the carbon-nitrogen cycle (CNO cycle) [1]. The sequence of the cold CNO cycle consists of the following reactions: $^{12}\text{C}(p,\gamma)^{13}\text{N}(\rightarrow ^{13}\text{C} + e^+ + \nu_e)$, $^{13}\text{C}(p,\gamma)^{14}\text{N}(p,\gamma)^{15}\text{O}(\rightarrow ^{15}\text{N} + e^+ + \nu_e)$, $^{15}\text{N}(p,\alpha)^{12}\text{C}$. In this sequence, four protons are transformed into α particle ($4p \rightarrow \alpha$) and as a result the energy of 26.73 MeV is released. Approximately 1.7 MeV of this energy is carried away by the neutrinos. As calculations show, the rate of energy release in the CNO cycle with increasing temperature ($T \sim 10^7$ K) increases much faster than the rate of energy release in the pp-chain.

The $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction is the first reaction of the hydrogen-burning CNO cycle and plays an important role as a source of generation of both nuclear energy [1] and low-energy neutrinos [2-5] in massive stars. In the low-energy region, the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ radiative capture reaction with formation of ^{13}N nucleus in the ground state mainly goes via both direct and resonant ($E^* = 2.365$ MeV, $J^\pi = 1/2^+$ and $E^* = 3.502$ MeV, $J^\pi = 3/2^-$) captures. Therefore, calculations of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factor, which is based on the analysis of experimental data, should take into account the contributions of the abovementioned two resonant and direct radiative captures, as well as their interference, in the energy

region $E_{c.m.} < 2.5$ MeV (c.m. here is the center of mass frame). Unfortunately, in the energy region between these two resonances, the available experimental data have large uncertainties [1].

Therefore, in this region $E_{p, lab.} = 1100-1400$ keV (lab. here is the laboratory frame), we have made new experimental measurements of the astrophysical S-factor for the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ radiative capture reaction with an uncertainty of about 16%. The calculation of the astrophysical S-factor, including the analysis of our new experimental data, has been made within the modified R-matrix method proposed previously in [6-9].

Experimental method and results of the measurements

The experimental part of our work was done on the electrostatic tandem accelerator UKP-2-1 of the Institute of Nuclear Physics ME RK in Almaty [10]. Protons were accelerated to energies $E_{p, lab.} = 340-1400$ keV. Calibration of proton energies in the beam was made with uncertainties of ± 1 keV according to the $^{19}\text{F}(p,\alpha\gamma)^{16}\text{O}$ and $^{27}\text{Al}(p,\gamma)^{28}\text{Si}$ reactions with many well-separated resonances in the region of $E_{p, lab.} = 340-1400$ keV [11, 12].

In our experiments, a specially made reaction chamber [6] with indium vacuum seals, a water-cooled target holder, and a quartz glass for obtaining a luminous image of the beam shape in front of the target were used. By an external handle, the quartz glass could be placed in front of the target for alignment. The γ -ray registration system was realized by using high-pure Germanium (HPGe) γ -detector with a Ge-crystal of volume of 111 cm^3 . To reduce the room and cosmic ray background the γ -detector was surrounded by 6 cm thick lead shield. The resolution of the γ -detector was about 5 keV at $E_\gamma = 2200 - 3250$ keV. The target was produced by sputtering natural carbon onto a 2 mm thick Cu substrate (thickness ≈ 2 mm, length ≈ 30 mm, and width ≈ 15 mm). The thickness of the carbon film sputtered onto the substrate was $110 \pm 8.8\ \mu\text{g}/\text{cm}^2$. The detailed description of the reaction chamber, the target production technology, and the target thickness determination method can be found in Refs. [6, 13, 14].

The absolute detector γ -ray efficiency for $E_\gamma = 2200 - 3250$ keV was determined by using γ -lines ($E_\gamma = 2034.92, 2598.58$ and 3253.6 keV) of a calibrated ^{56}Co source, with γ -lines intensities known to better than 4.5% [15].

When measuring the absolute efficiency, the detector and the ^{56}Co source were located precisely in the geometry of the experiment. At the same time, the statistical uncertainty in determining the number of counts for each γ -line was no more than 2%, and the electronics dead time did not exceed 1%.

The experimental differential cross sections of radiative capture were obtained at the measurement complex of INP [6], which allow to study the yields of the nuclear reactions on the extracted beams of the cyclotrons of the institute at the low and ultra low energies for the astrophysical and thermonuclear applications, see papers [6, 13, 14, 16].

During the measurements, γ -detector was about 8 cm away from the beam spot on the target. The γ -detector and ^{56}Co source were located with an uncertainty of about 1 mm. The dependence of the γ -ray registration rate on the source-detector distance was determined using ^{56}Co source. It was determined that at a distance of 8 cm, a deviation of ± 1 mm leads to a change in the γ -ray registration rate by $\pm 3\%$. Thus, uncertainties in the source and detector positions, dead time, γ -lines intensities, and counting statistics lead to an overall uncertainty of 6% for the detector efficiency over the entire energy interval of registered γ -rays (i.e. from 2200 to 3250 keV).

The differential cross sections of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for the transition to the ground state were determined at $E_{p, lab.} = 1100, 1150, 1250$ and 1400 keV and at $\theta_{\gamma, lab.} = 0^\circ$.

Beam currents ranged from 5 to 8 μA . The energy spread of the beam was determined by the width of the front of the $^{27}\text{Al}(p, \gamma)^{28}\text{Si}$ reaction yield curve near the resonance at $E_{p, lab.} = 992$ keV (resonance width < 0.1 keV) and did not exceed 1.5 keV. The accumulated charges on the target with an uncertainty of 3% were 0.28, 1.26, 1.51 and 1.4 Coulomb for $E_{p, lab.} = 1100, 1150, 1250$ and 1400 keV, respectively. Dead-time effects were kept below 1.5% at all beam energies.

Figure 1 shows the γ -ray spectrum obtained at $E_{p, lab.} = 1100$ keV and $\theta_{\gamma, lab.} = 0^\circ$. The energy calibration of the spectrometer was determined using well-known γ -lines of the ^{56}Co source and the room background γ -lines at 1461 keV (^{40}K) and at 2614 keV (Th).

The number of counts in the spectral peak with preliminarily subtracted background (trapezium shaped) divided by the calibrated integrator counter value was taken as the yield of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ capture

reaction. Statistical uncertainties in the determination of the yields (including uncertainties introduced by backgrounds subtracted) were about 15% for the measurement at $E_{p, \text{lab.}} = 1100$ keV and about 6% for the measurements at all other energies.

During each measurement, we computed the number of registered γ -rays of the transition to the ground state in ^{13}N (N_γ) over the integrator counter (N_p) as the yield per proton. For each of the energies presented in the work, the dependence of N_γ on N_p represented a straight line within the current statistical uncertainty of determining N_γ , which indicated the stability of the target and the stability of the beam position on it during the whole exposure.




Figure 1 - γ -ray spectrum of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for the radiative capture transition to the ground state in ^{13}N as obtained at the laboratory proton energy of 1100 keV ($\theta_{\gamma, \text{lab.}} = 0^0$) by our HPGe γ -detector of volume 111 cm³, located 8 cm. from the reaction region

Since, in the region $E_{p, \text{lab.}} = 1000\text{-}1400$ keV the differential cross sections change only slightly with energy, as can be seen, for example, from previous works [6, 17], the effective laboratory energies were found using the expression $E_{p, \text{eff}} = E_{p, \text{lab.}} - 0.5\Delta_{\text{lab.}}(E_{p, \text{lab.}})$, where $\Delta_{\text{lab.}}$ is the energy loss of protons in the target.

Because the γ -detector energy resolution and the proton beam energy spread are significantly less than the energy losses of protons in the target, the upper part of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction γ -line, for example, shown in Figure 1, repeats the course of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction yield curve in the corresponding energy region (the spectrum in Figure 1 has too big statistical uncertainties), and the width of this γ -line is largely due to the target thickness. This circumstance allowed us to determine the target thickness also by analyzing the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction γ -lines shapes (i.e. as a second independent method) and to confirm the value obtained by the first method within the uncertainties. Moreover, the second method allowed us to check the uniformity of the target thickness for all the spectra obtained.

The differential cross sections of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for radiative capture to the ground state of ^{13}N at $\theta_{\gamma, \text{lab.}} = 0^0$ were determined by using the relation

$$\frac{d\sigma}{d\Omega}(E_{p, \text{eff.}}, 0^0) = \frac{N_\gamma}{N_p N_{\text{C}^{12}} \varepsilon(E_{\gamma, \text{eff.}})}$$

where N_γ is the number of counts observed for the capture transition, $\varepsilon(E_{\gamma, \text{eff.}})$ is the absolute detector γ -ray efficiency, $E_{\gamma, \text{eff.}} = E_{p, \text{lab.}} \frac{12}{13} + 1941 - 0.5 \Delta_{\text{lab.}}(E_{p, \text{lab.}})$, N_p is the number of incident protons, and $N_{\text{C}^{12}}$ is the areal density of ^{12}C atoms.

As an additional test of the method for obtaining the absolute values of the differential cross sections, we have measured the yields over the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ resonance at $E_{p, \text{lab.}} = 457$ keV (the differential cross sections in this energy range are known, for example, from [6, 17] with uncertainties of no more than 10%). Due to the energy losses of protons in the target in this energy region match with the resonance width (which is about 40 keV) the yields in the resonance region were determined by the relation

$$N_{\gamma} = N_p N_{C^{12}} \int_{E_{p, \text{exit}}}^{E_{p, \text{lab.}}} \frac{d\sigma}{d\Omega}(\mathbf{E}, 0^0) \varepsilon(\mathbf{E}) \frac{d\Delta}{dE}(\mathbf{E}) dE$$

where $E_{p, \text{lab.}}$ is the proton energy at the target entrance ($E_{p, \text{lab.}} = 440, 450, 460, 470, 480$ keV), $E_{p, \text{exit}}$ is the corresponding proton energy at the target exit, $\frac{d\sigma}{d\Omega}(\mathbf{E}, 0^0)$ is the differential cross section of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for the capture to the ground state of ^{13}N at $\theta_{\gamma, \text{lab.}} = 0^0$ (the experimental data were taken from [6, 17]), $\frac{d\Delta}{dE}(\mathbf{E})$ is the stopping power of protons in carbon (calculated by the LISE++ program [18]). Such measurements were carried out before and after the main experiments, and the calculated and measured yields matched within the uncertainties.

The differential cross sections obtained in present work are given in the second row of Table 1. Assuming isotropy in the angular distributions of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction in the energy region of incident protons from 400 to 1390 keV (which is confirmed with uncertainties of 10% in the works [6, 17]) in the present work, the total sections are calculated according to the formula:

$$\sigma(E_{p, \text{eff.}}) = 4\pi \frac{d\sigma}{d\Omega}(E_{p, \text{eff.}}, 0^0)$$




Figure 2 - The astrophysical S-factor for the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction. The experimental data: filled circles are the result of the present work, filled triangles are the data from [6] and open circles are the data from [17]. The solid line is our fit, dotted line is our calculated contribution for the direct radiative capture, dashed (dashed-dotted) line shows our calculated contribution for the first (second) resonance and dashed-dotted-dotted line presents our calculated contribution for the third ($E^* = 10.250$ MeV, $J^\pi = 1/2^+$) γ -resonance tail

Further, according to the relation:

$$S(\text{keV b}) = \sigma(b) E(\text{keV}) \exp\left(\frac{180.29}{\sqrt{E(\text{keV})}}\right)$$

the astrophysical S-factors were calculated, which are given in the third row of Table 1 and are shown in Figure 2 in comparison with the results of previous works. From Figure 2, it is clear that the experimental data of our work are in good agreement with the results obtained in [6, 17].

Table 1 + Experimental differential cross sections and experimental astrophysical S - factors of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for radiative capture to the ground state of ^{13}N

$E_{p,\text{eff.}}(\text{keV})$	1088	1138.5	1239.3	1390
$\frac{d\sigma}{d\Omega}^{\text{exp.}}(E,0^0) - \left(\frac{\mu\text{b}}{\text{sr}}\right)$	0.027±0.0049	0.029±0.0035	0.048±0.0058	0.064±0.0077
$S^{\text{exp.}}(E) - (\text{MeV b})$	0.1±0.02	0.098±0.015	0.14±0.022	0.16±0.025

Results of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factor calculation by the modified R-matrix analysis method

The detailed description of the modified R-matrix method that was used in the calculations can be found in [6-9, 19, 20]. The input parameters required for calculating the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factor for transition to the ground state of ^{13}N (channel radius, proton width and radiative widths of the first and second resonance states) were taken from [6]. The experimental results of this work and the experimental results of [6, 17] were used as experimental data. The asymptotic normalization coefficient (ANC) value was taken equal to $1.43 \pm 0.09 \text{ fm}^{-1/2}$ [21, 22] as in [6].

The results of our calculations of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factor taking into account the analysis of new experimental data completely repeat our earlier results [6] and are shown in Figure 2, where the contributions of direct radiative capture (dotted line), first resonance (dashed line) and second resonance (dashed-dotted line) are given separately. The inclusion of the third resonant state ($E^* = 10.250 \text{ MeV}$, $J^\pi = 1/2^+$, the proton width $\Gamma_3^p = 280 \text{ keV}$ [23] and the radiation width $\Gamma_3^\gamma = 6000 \text{ eV}$ [6]) tail part contribution in the calculations significantly improved the theoretical description of the experimental data (dashed-dotted-dotted line). The values of the resonant parameters used in present work are listed in Table 2 of [6].

The results of our calculations of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factors for the transition to the ground state of ^{13}N at the most important energies for astrophysics $E = 0; 25$ and 50 keV are $S(0 \text{ keV}) = 1.62 \pm 0.20 \text{ keVb}$, $S(25 \text{ keV}) = 1.75 \pm 0.22 \text{ keVb}$ and $S(50 \text{ keV}) = 1.88 \pm 0.24 \text{ keVb}$, respectively. The uncertainties quoted for these astrophysical S-factors are due to those of the parameters of proton and γ widths and ANC given earlier [6]. Our central value for $S(25 \text{ keV})$ within the specified uncertainty is consistent with the values of $S(25 \text{ keV}) = 1.54 \pm 0.08 \text{ keVb}$ obtained in [24] and $S(25 \text{ keV}) = 1.45 \pm 0.20 \text{ keVb}$ obtained in [17], and in satisfactory agreement with the value of $S(25 \text{ keV}) = 1.33 \pm 0.15 \text{ keVb}$ obtained in [25, 26]. However, our result for $S(0 \text{ keV})$ is noticeably larger than that of $S(0 \text{ keV}) = 1.0$ and 1.3 keVb obtained in [27] using the Minnesota and V2 forms of the NN potential, respectively, as well as the value of $S(0 \text{ keV}) = 1.4 \text{ keVb}$ recommended in [28]. It should be emphasized that a value of $S(25 \text{ keV}) = 1.54 \pm 0.08 \text{ keVb}$ in [24] has been also obtained within the R-matrix approach [7, 9]. In contrast to our work in [24], the ANC, which is responsible for the contribution of direct radiative capture, was a fitting parameter in order to better describe the experimental data at first resonance region. As a result, this artificial overestimation of the ANC value led to resonance ($\Gamma_1^\gamma = 0.50 \pm 0.05 \text{ eV}$) decrease in the total amplitude of the radiative capture process, which led to an underestimated value of $S(25 \text{ keV})$. In our present work, we used the fixed ANC value obtained independently from the analysis of the peripheral proton transfer reaction [21, 22], which allowed us earlier in [6] to carry out fitting of resonant width parameters (for example, $\Gamma_1^\gamma = 0.65 \pm 0.07 \text{ eV}$) in a correct way.

$^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction rate

The calculated astrophysical S-factors, as well as the data of [6, 17], were used for calculating the rate of ^{13}N nucleus formation in the stellar interior as a function of stellar temperature T_9 , where $T_9 = T \times 10^9$ K. The Maxwellian-averaged reaction rates $N_A\langle\sigma v\rangle$ as a function of temperature are defined by

$$N_A\langle\sigma v\rangle = N_A \left(\frac{8}{\pi\mu} \right)^{1/2} (k_B T)^{-3/2} \int_0^\infty \sigma(E) \exp(-E/k_B T) E dE,$$

where N_A is Avogadro's number, k_B is the Boltzmann constant, and $\mathcal{G} = \sqrt{2E/\mu}$ is the relative velocity of the colliding particles. The calculation performed in the present work matches with the results we obtained in [6]. Figure 3(a) shows the reaction rates of our calculation (solid line) and its comparison with the experimental data of [29]. It is seen that the result of our calculation is in good agreement with those recommended in [29]. That work used a very different method, with independent systematic uncertainties, and cited uncertainties equal to the estimated values for unobserved energies. The ratio of our calculation of reaction rates $N_A\langle\sigma v\rangle$ to the result recommended in [30] (solid line) is also given in Figure 3(b). As is seen from Figure 3(b) there is a noticeable difference (up to $\approx 20\%$) between our recommended results and those given in [30] within a wide interval of stellar temperatures. The probable reason for the observed difference is that in [30] the calculation of the reaction rates included all the experimental astrophysical S-factors obtained in [17, 25, 26, 30], some of which have uncertainties up to 40%, by a smooth spline fit. Moreover, it was assumed in [30] that the spectroscopic factor for the ground state of the ^{13}N nucleus in ($^{12}\text{C} + p$) - configuration is 1. However, as can be seen from [31], this assumption is not justified and, in fact, the empirical value of the spectroscopic factor is 0.55 ± 0.18 . It can be considered that the result in [30] is model-dependent, while our calculation of the reaction rate does not contain free parameters and in this sense it can be regarded as more reliable.




Figure 3 - (a) The $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction rate calculated in present work (solid line) and points taken from [29].
 (b) The ratio of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction rates $N_A\langle\sigma v\rangle$ of present work to those from [30] (solid line)

Conclusion

In this work new experimental data on the differential cross sections of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for the transition to the ground state of ^{13}N have been obtained at four energies from 1088 to 1390 keV in the laboratory system at an angle of 0° with uncertainties of about 12%. The astrophysical S-factors for radiative capture of a proton to the ground state of ^{13}N (the only bound state of this nucleus) have been determined with uncertainties of about 16% at these energies. Our experimental data are in good agreement with those previously obtained in [6, 17].

We have analyzed the new experimental data on $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction astrophysical S-factors for the transition to the ground state of ^{13}N at extremely low energies within the one-level R-matrix approach where the direct part of the amplitude is expressed in terms of the ANC for ^{13}N in the $(p + ^{12}\text{C})$ channel. Such a parametrization allowed us to calculate the direct capture part of the amplitude in a correct way using the indirectly measured value of ANC found previously in [21, 22] from the analysis of the peripheral $^{12}\text{C}(^3\text{He},d)^{13}\text{N}$ reaction.

It is shown that using information about ANC value provides good fitting of the experimental astrophysical S-factor of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for the transition to the ground state of ^{13}N and reduces to a minimum the model dependence of the calculated direct capture part of the astrophysical S-factor on the parameters of the R-matrix approach. In general, the results of the new analysis of the experimental astrophysical S-factors match with the results of our previous work [6].

We have also calculated the rates of the $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction for the astrophysically important transition to the ground state of ^{13}N in the low energy region. It has been shown that these present reaction rates (as well as the rates of [6]) are in good agreement with those recommended in [29] using a very different technique from that described in present work, whereas a notable difference (up to $\approx 20\%$) occurs between our result (as well as the result of [6]) and that given in [30] within a wide interval of stellar temperatures.

Acknowledgments

The authors would like to thank the UKP-2-1 staff for assistance with the experiment. This work was supported by program targeted funding titled “Development of the complex scientific research activities in nuclear and radiation physics based on the Kazakhstan’s accelerator facilities” # 0118PK01174.

Д.М. Зазулин^{1,2}, Н. Буртебаев^{1,2}, Р.Ж. Петерсон³, С.В. Артемов⁴,
С. Игамов⁴, Ж.К. Керимкулов¹, Д.К. Алимов¹, Е.С. Мухамеджанов^{1,2},
Маулен Насурлла^{1,2}, А. Сабидолла¹, Маржан Насурлла^{1,2}, Р. Ходжаев^{1,2}

¹Ядролық физика институты, Алматы, Қазақстан;

²аль-Фараби атындағы ҚазҰУ, Алматы, Қазақстан;

³Колорадо Университеті, Болдер, Колорадо, АҚШ;

⁴Ядролық физика институты, Ташкент, Узбекистон

ТӨМЕН ЭНЕРГИЯЛАРДАҒЫ P-¹²C РАДИАЦИЯЛЫҚ ҚАРПУЫНЫҢ ЖАҢА НӘТИЖЕЛЕРІ

Аннотация. Жұмыста үдетілінген протондардың 1088 – 1390 кэВ энергияларында 0^0 бұрышта $^{12}\text{C}(p,\gamma)^{13}\text{N}$ радиациялық қарпуы реакциясының ^{13}N ядросының негізгі күйіне көшуінің дифференциалдық қимасының жаңа өлшеулер нәтижелері (12% дәлдікпен) ұсынылды. Алынған дифференциалдық қималар және осы энергия аймағындағы бұрыштық таралулардың изотроптық қасиетке ие болатындығына негізделі отырып, 16% дәлдікпен $^{12}\text{C}(p,\gamma)^{13}\text{N}$ реакциясы үшін ^{13}N негізгі күйге ауысуының астрофизикалық S факторы анықталды. Қателер шегінде, осы жұмыстың эксперименттік нәтижелері бұрынғы жұмыстардың деректерімен сәйкес келеді. Модификацияланған R - матрицалық әдісті пайдалана отырып, астрофизикалық S-фактор бойынша эксперименттік мәліметтер талданды. Аса төменгі энергияларда ^{12}C ядросымен протондардың тікелей қарпылуына байланысты есептеу қателіктерін азайту мақсатында талдау барысында ^{12}C және ^{13}N ядроларының байланысқан күйлерінің толқындық функцияларының қабаттасу интегралының өлшенілген асимптотикалық нормалау коэффициентінің мәні пайдаланылды. Массалар орталығы жүйесінде $E = 0,25$ және 50 кэВ энергияларда $^{12}\text{C}(p,\gamma)^{13}\text{N}$ реакциясынан ^{13}N ядросының негізгі күйге ауысуы үшін S-фактордың есептік мәндері келтірілген. $0 - 10^{10}$ К температура аймағында $^{12}\text{C}(p,\gamma)^{13}\text{N}$ термоядролық реакциясының жылдамдықтары анықталды. Бұл жұмыстың есептеу нәтижелері бұрынғы жұмыстардың эксперименттік және есептік мәліметтерімен салыстырылды.

Түйін сөздер: дифференциалдық қималар, толық қима, астрофизикалық S-фактор, асимптотикалық нормалау коэффициенті, реакцияның жылдамдықтары.

Д.М. Зазулин^{1,2}, Н. Буртебаев^{1,2}, Р.Ж. Петерсон³, С.В. Артемов⁴,
С. Игамов⁴, Ж.К. Керимкулов¹, Д.К. Алимов¹, Е.С. Мухамеджанов^{1,2},
Маулен Насурлла^{1,2}, А. Сабидолда¹, Маржан Насурлла^{1,2}, Р. Ходжаев^{1,2}

¹Институт ядерной физики, Алматы, Казахстан;

²КазНУ им. аль-Фараби, Алматы, Казахстана;

³Университет Колорадо, Болдер, Колорадо, США;

⁴Институт ядерной физики, Ташкент, Узбекистан

НОВЫЕ РЕЗУЛЬТАТЫ ДЛЯ РАДИАЦИОННОГО ЗАХВАТА p -¹²C ПРИ НИЗКИХ ЭНЕРГИЯХ

Аннотация. Представлены результаты новых измерений дифференциальных сечений реакции $^{12}\text{C}(p,\gamma)^{13}\text{N}$ радиационного захвата на основное состояние ^{13}N для угла θ^0 и при энергиях налетающих протонов от 1088 до 1390 кэВ (точность около 12%). На основе полученных дифференциальных сечений и в предположении об изотропном характере угловых распределений в данной области энергий с точностью около 16% определены астрофизические S – факторы реакции $^{12}\text{C}(p,\gamma)^{13}\text{N}$ для перехода на основное состояние ^{13}N . В пределах погрешностей экспериментальные результаты настоящей работы согласуются с данными более ранних работ. С использованием модифицированного R – матричного метода проведен анализ экспериментальных данных по астрофизическому S – фактору. В целях минимизации вычислительной неопределенности, связанной с прямым захватом протона ядром ^{12}C для самых низких энергий, при анализе использовалось значение измеренного ранее асимптотического нормировочного коэффициента интеграла перекрытия волновых функций связанных состояний ядер ^{12}C и ^{13}N . Для энергий $E = 0,25$ и 50 кэВ в системе центра масс приведены вычисленные значения S – фактора реакции $^{12}\text{C}(p,\gamma)^{13}\text{N}$ для перехода на основное состояние ^{13}N . В области температур от 0 до 10^{10} К получены скорости термоядерной реакции $^{12}\text{C}(p,\gamma)^{13}\text{N}$. Результаты расчетов настоящей работы сравниваются с экспериментальными и расчетными данными предыдущих работ.

Ключевые слова: дифференциальные сечения, полные сечения, астрофизический S -фактор, асимптотический нормировочный коэффициент, скорости реакции.

Information about authors:

Zazulin Denis Michailovich - Institute of Nuclear Physics, Almaty, Kazakhstan, Senior Researcher, candidate of physical and mathematical sciences, denis_zazulin@mail.ru, experimental data processing and analyzing of data;

Burtebayev Nassurlla - Institute of Nuclear Physics, Almaty, Kazakhstan, Head of laboratory, Doctor of physical and mathematical sciences, Professor of Physics, nburtebayev@yandex.ru, setting and planning tasks;

Peterson Roy Jerome – University of Colorado, Boulder, Colorado, USA, Professor of Physics, jerry.peterson@colorado.edu, setting and planning tasks;

Artemov Sergey Viktorovich - Institute of Nuclear Physics, Tashkent, Uzbekistan, Doctor of physical and mathematical sciences, professor, artemov1943@gmail.com, setting and planning tasks;

Igamov Sayram - Institute of Nuclear Physics, Tashkent, Uzbekistan, candidate of physical and mathematical sciences, professor, igamov@inp.uz, theoretical calculations;

Kerimkulov Zhambul Kuanyshbekovich - Institute of Nuclear Physics, Almaty, Kazakhstan, Senior Researcher, candidate of physical and mathematical sciences, zhambul-k@yandex.ru, experimental data processing and analyzing of data;

Alimov Dilshod Kamalovich - Institute of Nuclear Physics, Almaty, Kazakhstan, Senior Researcher, PhD, diliyo@mail.ru, analysis of experimental data;

Mukhamejanov Yerzhan Serikovich - Institute of Nuclear Physics, Almaty, Kazakhstan, Senior Researcher, PhD, craftinho@mail.ru, analysis of experimental data;

Marzhan Nassurlla - Institute of Nuclear Physics, Almaty, Kazakhstan, Senior Researcher, PhD, morzhic@mail.ru, analysis of experimental data;

Nassurlla Maulen - al-Farabi Kazakh National University, Almaty, Kazakhstan, PhD student, nepad@mail.ru, experimental data processing and analyzing of data;

Sabidolda Auganbek - Institute of Nuclear Physics, Almaty, Kazakhstan, engineer, asabidolda@mail.ru, obtaining experimental data.

Khojayev Romazan - Institute of Nuclear Physics, Almaty, Kazakhstan, engineer, ramazan_inp@mail.ru, obtaining experimental data.

REFERENCE

- [1] Rolfs C and Rodney WS (1988) *Cauldrons in the Cosmos*, The University of Chicago Press, Chicago and London. ISBN- 13: 978-0226724577
- [2] Bahcall JN and Pinsonneault MH (1992) Standard solar models, with and without helium diffusion, and the solar neutrino problem, *Rev Mod Phys*, 64: 885. DOI: 10.1103/RevModPhys.64.885 (in English).
- [3] Bahcall JN, Basu S and Pinsonneault MH (1998) How uncertain are solar neutrino predictions?, *Phys Lett B*, 433: 1-8. DOI: [10.1016/S0370-2693\(98\)00657-1](https://doi.org/10.1016/S0370-2693(98)00657-1) (in English).
- [4] Adelberg EG et al (1998) Solar fusion cross sections, *Rev Mod Phys*, 70:1265. DOI: 10.1103/RevModPhys.70.1265 (in English).
- [5] Kirsten TA (1999) Solar neutrino experiments: results and implications, *Rev Mod Phys* 71:1213. DOI: 10.1103/RevModPhys.71.1213 (in English).
- [6] Burtebaev N, Igamov SB, Peterson RJ, Yarmukhamedov R and Zazulin DM (2008) New measurements of the astrophysical S factor for $^{12}\text{C}(p,\gamma)^{13}\text{N}$ reaction at low energies and the asymptotic normalization coefficient (nuclear vertex constant) for the $p+^{12}\text{C}\rightarrow^{13}\text{N}$ reaction, *Phys Rev C*, 78: 035802. DOI: 10.1103/PhysRevC.78.035802 (in English).
- [7] Artemov SV, Bajajin AG, Igamov SB, Nie GK, Yarmukhamedov R (2008) Nuclear asymptotic normalization coefficients for $^{14}\text{N}\rightarrow^{13}\text{C}+p$ configurations and astrophysical S factor for radiative proton capture, *Physics of Atomic nuclei*, 71:998. DOI: 10.1134/S1063778808060045 (in English).
- [8] Artemov SV, Igamov SB, Tursunmakhatov QI, Yarmukhamedov R (2012) Extrapolation of astrophysical S factors for the reaction $^{14}\text{N}(p,\gamma)^{15}\text{O}$ to near-zero energies, *Phys Atom Nucl*, 75: 291. DOI: 10.1134/S1063778812020032 (in English).
- [9] Igamov SB, Artemov SV, Yarmukhamedov R, Burtebayev N, Sakuta SB (2016) Astrophysical S factors for the reaction $^6\text{Li}(p,\gamma)^7\text{Be}$ at ultralow energies, *Bulletin of National Nuclear Centre of the Republic of Kazakhstan [Vestnik Nazionalnogo Yadernogo Centra RK]* 56:82. (in English).
- [10] Arzumanov AA (1992) *Proceedings of the 13th particle accelerator conference, Dubna, Russia*. P. 118.
- [11] Butler JW (1959) *Table of (p, γ) Resonances*, NRL Report, 5282. (in English).
- [12] Lyons PB, Toevs JW and Sargood DG (1969) Total yield measurements in $^{27}\text{Al}(p,\gamma)^{28}\text{Si}$, *Nucl Phys A*, 130: 1. DOI: [10.1016/0375-9474\(69\)90954-3](https://doi.org/10.1016/0375-9474(69)90954-3) (in English).
- [13] Dubovichenko SB, Burtebaev N, Zazulin DM, Kerimkulov ZhK and Amar ASA (2011) Astrophysical S factor for the radiative-capture reaction $p\ ^6\text{Li}\rightarrow\ ^7\text{Be}$, *Phys Atom Nucl*, 74: 984. DOI: 10.1134/S1063778811050073 (in English).
- [14] Dubovichenko S, Burtebayev N, Dzhazairov-Kakhramanov A, Zazulin D, Kerimkulov Zh, Nassurlla M, Omarov C, Tkachenko A, Shmygaleva T, Stanislaw Kliczewski S, Sadykov T (2017) New measurements and phase shift analysis of $p^{16}\text{O}$ elastic scattering at astrophysical energies, *Chinese Physics C*, 41: 014001. DOI: [10.1088/1674-1137/41/1/014001](https://doi.org/10.1088/1674-1137/41/1/014001) (in English).
- [15] Barker PH and Connor RD (1967) ^{56}Co as a calibration source up to 3.5 MeV for gamma ray detectors, *Nucl Inst And Meth*, 57:147. DOI: [10.1016/0029-554X\(67\)90513-7](https://doi.org/10.1016/0029-554X(67)90513-7) (in English).
- [16] Maulen Nassurulla, Burtebayev N, Kerimkulov Zh K, Suzuki T, Sakuta SB, Marzhan Nassurulla, Khojayev R (2018) Investigation of deuteron scattering by ^7Li nuclei at energy of 14.5 MeV, *News of the National Academy of Sciences of the Republic of Kazakhstan, series physico-mathematical [Izvestiya Nazionalnoi Akademii Nauk RK serija fizicheskaja]* 6:15-22. <https://doi.org/10.32014/2018.2518-1726.12> (in English).
- [17] Rolfs C and Azuma RE (1974) Interference effects in $^{12}\text{C}(p,\gamma)^{13}\text{N}$ and direct capture to unbound states, *Nucl Phys A*, 227: 291. DOI: [10.1016/0375-9474\(74\)90798-2](https://doi.org/10.1016/0375-9474(74)90798-2) (in English).
- [18] <http://lise.nsl.msui.edu/lise.html>
- [19] Barker FC and Kajino T (1991) The $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ cross section at low energies, *Aust J Phys*, 44: 369. DOI: 10.1071/PH910369 (in English).
- [20] Holt RJ, Jackson HE, Laszewski RM, Monahan JE, Specht JR (1978) [Effects of channel and potential radiative transitions in the \$^{17}\text{O}\(\gamma,n\)^{16}\text{O}\$ reaction](https://doi.org/10.1103/PhysRevC.18.1962), *Phys Rev C*, 18: 1962. DOI: 10.1103/PhysRevC.18.1962 (in English).
- [21] Yarmukhamedov R (1997) Influence of the three-particle Coulomb effects on the spectroscopic information extracted from analysis of the $^{12,13}\text{C}(\gamma,\text{He},\text{d})^{13,14}\text{N}$ surface reactions, *Nuclear Physics [Yadernaya Fizika]* 60: 1017. (in Russian).
- [22] Artemov SV, Zaparov EA, Nei GK, Nadirbekov M, and Yarmukhamedov R (2002) Influence of the three-body coulomb effects on extracted value of nuclear vertex constants from transfer reactions, *Bulletin of the Russian Academy of Sciences [Izvestiya Akademii Nauk Rossii serija fizicheskaja]* 66:60. (in Russian).
- [23] Ajzenberg-Selove F (1986) Energy levels of light nuclei A = 13–15, *Nucl Phys A*, 449: 1-15. DOI: [10.1016/0375-9474\(86\)90119-3](https://doi.org/10.1016/0375-9474(86)90119-3) (in English).
- [24] Barker FC and Ferdous N (1980) The first excited state of ^9B , *Aust J Phys*, 33: 691. DOI: [10.1071/PH870307](https://doi.org/10.1071/PH870307) (in English).
- [25] Hebbard DF and Vogl JL (1960) Elastic scattering and radiative capture of protons by C^{13} , *Nucl Phys*, 21: 652. DOI: [10.1016/0029-5582\(60\)90084-5](https://doi.org/10.1016/0029-5582(60)90084-5) (in English).
- [26] Vogl JL (1963) Ph.D. thesis, California Institute of Technology. (in English).
- [27] Dufour M and Descouvemont P (1997) Multicenter study of the $^{12}\text{C}+n$ and $^{12}\text{C}+p$ systems, *Phys Rev C*, 56: 1831. DOI: 10.1103/PhysRevC.56.1831 (in English).
- [28] Caughlan GR and Fowler WA (1988) Thermonuclear reaction rates V, *At Data Nucl Data Tables*, 40: 283. DOI: [10.1016/0092-640X\(88\)90009-5](https://doi.org/10.1016/0092-640X(88)90009-5) (in English).
- [29] Roughton NA, Fritts MR, Peterson RJ, Zaidins CS, and Hansen CJ (1979) Thick-target measurements and astrophysical thermonuclear reaction rates: Proton-induced reactions, *At Data Nucl Data Tables*, 23: 177. DOI: [10.1016/0092-640X\(79\)90004-4](https://doi.org/10.1016/0092-640X(79)90004-4) (in English).
- [30] Angulo C, Arnould M et al (1999) A compilation of charged-particle induced thermonuclear reaction rates, *Nucl Phys A*, 656: 3. DOI: [10.1016/S0375-9474\(99\)00030-5](https://doi.org/10.1016/S0375-9474(99)00030-5) (in English).
- [31] King JD, Azuma RE, Vise JB et al (1994) Cross section and astrophysical S-factor for the $^{13}\text{C}(p,\gamma)^{14}\text{N}$ reaction, *Nucl Phys A*, 567: 354. DOI: [10.1016/0375-9474\(94\)90154-6](https://doi.org/10.1016/0375-9474(94)90154-6) (in English).