

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2020.2518-1726.21>

Volume 2, Number 330 (2020), 103 – 111

UDK 517.951

MRNTI 27.31.15

B.D. Koshanov¹, G.D. Koshanova², G.E. Azimkhan³, R.U. Segizbayeva⁴¹Institute of Mathematics and Mathematical modeling, Almaty, Kazakhstan;^{1,3}Abai Kazakh National Pedagogical University, Almaty, Kazakhstan;²Yasavi International Kazakh-Turkish University, Turkistan, Kazakhstan;⁴Civil Aviation Academy, Almaty, Kazakhstan.E-mail: koshanov@list.ru, gulnazira.koshanova@ayu.edu.kz,
g_azimhanova@mail.ru, segizbaeva-55@mail.ru

SOLVABILITY OF BOUNDARY VALUE PROBLEMS WITH NON-LOCAL CONDITIONS FOR MULTIDIMENSIONAL HYPERBOLIC EQUATIONS

Abstract. In this paper, we study the solvability of new nonlocal boundary value problems for hyperbolic equations in a multidimensional bounded domain. For the problem under study, the existence and uniqueness theorems of regular solutions are proved.

Keywords: hyperbolic equation, boundary value problems, regular solutions, existence, uniqueness.

1. Introduction and statement of the problem. It is well known that studies of the properties of correcting boundary value problems for differential equations are important both for applications and for mathematical and mechanical problems. The theory of boundary value problems for high-order elliptic equations attracts particular attention of mathematicians.

The aim of the work is to study the solvability in classes of regular solutions of a new boundary value problem for the equation

$$u_{tt} - \Delta u + c(x, t)u = f(x, t), (x, t) \in Q. \quad (1)$$

This equation is an ordinary second-order hyperbolic equation, and the solvability of natural boundary value and initial-boundary value problems for it is well studied [24-26].

The study of the solvability of boundary value problems for quasi-hyperbolic equations (by analogy with quasi-elliptic equations) apparently began with the works of V.N. Vragova [1,2]. A study of the solvability of problems for high orders of equations in Sobolev spaces was carried out in [1-3], a number of similar results were obtained in [4-11].

Investigations of nonlocal problems with integral conditions for linear parabolic equations, for differential equations of odd orders, and for some classes of non-stationary equations, have recently been actively studied in the works of A.I. Kozhanova [7,9,10]. In [1-6], the main attention was paid to situations related to degenerate equations of the form (1).

In [12], a criterion was obtained for the strong solvability of the mixed Cauchy problem for the Laplace equation.

It is known that the Dirichlet problem for polyharmonic equations is uniquely solvable for any right-hand side of the equation. In [13-16], a new representation of the Green function of the Dirichlet problem for the polyharmonic equation in a multidimensional ball is constructed explicitly. In [17,18], a representation of the Green function of the Neumann problem for the Poisson equation in a multidimensional unit ball was obtained.

The problems of finding solvability conditions for boundary value problems for a polyharmonic equation in a ball were investigated in [19]. In [20], for derivatives of the $2l$ -th order equation with constant (and only higher) real coefficients, normal derivatives were studied under boundary conditions. For these problems, sufficient conditions for the Fredholm solvability of the problem are obtained and formulas for the index of the problem are given.

In [21-23], the behavior of solutions of the Dirichlet problem for the Poisson equations and the biharmonic equation in an unbounded domain was studied.

In this paper, we obtain a theorem on the existence and uniqueness of a regular solution of a nonlocal in time boundary value problem for hyperbolic equations in a multidimensional bounded domain by the method of a priori estimation and passage to the limit.

Let Ω - be a bounded region of space R^n with a smooth boundary Γ , Q - a cylinder $\Omega \times (0, T)$, $0 < t < T < +\infty$, $S = \Gamma \times (0, T)$ - a side boundary Q , $f(x, t)$ - given functions defined for $x \in \overline{\Omega}$, $t \in [0, T]$

For equation (1) we study the time-nonlocal boundary value problem

$$u|_S = 0, \tag{2}$$

$$u(x, 0) = \alpha u(x, T), \tag{3}$$

$$u_t(x, 0) = \beta u_t(x, T), \alpha, \beta \in R. \tag{4}$$

In studies of this kind of nonlocal problems, the parameter continuation method, a priori estimation method, and passage to the limit are usually used. For a hyperbolic equation, the parameter continuation method is not applicable, since the smoothness of the right side of the equation will be lost.

We define a functional space in which the properties of uniqueness and existence of a solution to the boundary value problem (1) - (4) will be studied. Namely, we define space $W_2^{2,2}(Q)$ as the set of functions from space $L_2(Q)$, that have generalized derivatives of spatial variables up to the second order inclusive and with respect to variable t up to the second order inclusive, belonging to the same space. We define the norm in space $W_2^{2,2}(Q)$.

$$\|v\|_{W_2^{2,2}(Q)} = \left(\int_Q \left[v^2 + \sum_{i,j=1}^n v_{x_i x_j}^2 + (v_t)^2 \right] dx dt \right)^{1/2}$$

obviously, space $W_2^{2,2}(Q)$ with this norm will be a Banach space.

2. *The regularized nonlocal boundary value problem and the main result.* We consider the following regularized problem in order to apply the continuation method with respect to the parameter

$$u_{tt} - \Delta u + c(x, t)u - \varepsilon \Delta u_t = f(x, t) \in L_2(Q) \tag{5}$$

$$u|_S = 0 \tag{6}$$

$$u(x, 0) = \lambda \alpha u(x, T), \tag{7}$$

$$u_t(x, 0) = \lambda \beta u_t(x, T). \tag{8}$$

For the regularized problem (5) - (8), we obtain estimates in the corresponding spaces. To this end, we multiply equation (5) by u_t integrate over Q , then we obtain.

$$\int_Q u_{tt} u_t dx dt - \int_Q \Delta u u_t dx dt + \int_Q c u u_t dx dt - \varepsilon \int_Q \Delta u_t u_t dx dt = \int_Q f u_t dx dt;$$

$$I_1 = \frac{1}{2} \int_Q \frac{\partial}{\partial t} (u_t^2) dx dt = \frac{1}{2} \int_{\Omega} [u_t^2(0, T) - u_t^2(x, 0)] dx = \frac{(1 - \lambda^2 \beta^2)}{2} \int_{\Omega} u_t^2(x, T) dx;$$

Here we take into account the boundary condition (7) and $|\beta| < 1$, such that $\frac{1 - \lambda^2 \beta^2}{2} > 0$.

$$\begin{aligned}
 I_2 &= - \int_Q \Delta u u_t = \sum_{i=1}^n \int_Q u_{x_i} u_{x_i t} dx dt = \frac{1}{2} \sum_{i=1}^n \int_Q \frac{\partial}{\partial t} (u_{x_i}^2) dx dt = \\
 &= \frac{1}{2} \sum_{i=1}^n \int_{\Omega} [u_{x_i}^2(x, T) - u_{x_i}^2(x, 0)] dx = \frac{1 - \lambda^2 \alpha^2}{2} \sum_{i=1}^n \int_Q u_{x_i}^2(x, T) dx ; \\
 I_3 &= \int_Q c u u_t dx dt = \frac{1}{2} \int_Q \frac{\partial}{\partial t} (c u^2) dx dt - \frac{1}{2} \int_Q c_t u^2 dx dt = \\
 &= \frac{1}{2} \int_{\Omega} [c(x, T) u^2(x, T) - c(x, 0) u^2(x, 0)] dx - \frac{1}{2} \int_Q c_t u^2 dx dt = \\
 &= \frac{1}{2} \int_{\Omega} [c(x, T) - \lambda^2 \alpha^2 c(x, 0)] u^2(x, T) dx - \frac{1}{2} \int_Q c_t u^2 dx dt ; \\
 I_4 &= -\varepsilon \int_Q \Delta u_t u_t dx = -\varepsilon \sum_{i=1}^n \int_Q u_{x_i x_i t} u_t dx dt = \varepsilon \sum_{i=1}^n \int_Q u_{x_i t}^2 dx dt ; \\
 &\frac{1 - \lambda^2 \beta^2}{2} \int_{\Omega} u_t^2(x, T) dx + \frac{1 - \lambda^2 \alpha^2}{2} \sum_{i=1}^n \int_{\Omega} u_{x_i}^2(x, T) dx + \\
 &+ \frac{1}{2} \int_{\Omega} [c(x, T) - \lambda^2 \alpha^2 c(x, 0)] u^2(x, T) dx + \frac{1}{2} \int_Q \Delta u_t u_t dx dt + \varepsilon \sum_{i=1}^n \int_Q u_{x_i t}^2 dx dt = \int_Q f u_t dx dt ;
 \end{aligned}$$

We will require that the conditions are met

$$c(x, T) - \lambda^2 \alpha^2 c(x, 0) \geq 0 \quad , \quad c_t(x, T) \leq 0 .$$

Under the condition $\vartheta|_S = 0$ the embedding theorem holds, i.e. Friedrichs inequality [20]:

$$\int_{\Omega} \vartheta(x, T) dx \leq m_0 \sum_{i=1}^n \int_{\Omega} \vartheta_{x_i}^2(x, T) dx . \quad (F)$$

We apply it to the function $\vartheta = u_t$.

$$I_f = - \int_Q f u_t dx dt \leq \frac{\delta^2}{2} \int_Q u_t^2 dx dt + \frac{1}{2\delta^2} \int_Q f^2 dx dt \leq \frac{\delta^2 m_0}{2} \sum_{i=1}^n \int_Q u_{x_i t}^2 dx dt + \frac{1}{2\delta^2} \int_Q f^2 dx dt .$$

We choose $\frac{\delta^2 m_0}{2} = \frac{\varepsilon}{2}$.

Then we have

$$\frac{1 - \beta^2}{2} \int_{\Omega} u_t^2(x, T) dx + \frac{1 - \alpha^2}{2} \sum_{i=1}^n \int_{\Omega} u_{x_i}^2(x, T) dx + \varepsilon \sum_{i=1}^n \int_Q u_{x_i t}^2 dx dt \leq \frac{1}{2\varepsilon} \int_Q f^2 dx dt ; \quad (9)$$

At the beginning, we must obtain a uniform estimate for λ for a fixed ε . Equation (5) is multiplied by u_{tt} and integrated over Q , then we obtain

$$- \int_Q u_{tt} \Delta u_t dx dt + \int_Q \Delta u \Delta u_t dx dt - \int_Q c u \Delta u_t dx dt + \varepsilon \int_Q \Delta u_t^2 dx dt = - \int_Q f \Delta u_t dx dt .$$

Then we have

$$\frac{1 - \lambda^2 \beta^2}{2} \sum_{i=1}^n \int_{\Omega} u_{x_i t}^2(x, T) dx + \frac{1 - \lambda^2 \alpha^2}{2} \int_{\Omega} [\Delta u(x, T)]^2 dx + \varepsilon \int_Q (\Delta u_t)^2 dx dt =$$

$$= - \int_Q f \Delta u_t dxdt + \int_Q cu \Delta u_t dxdt ;$$

Estimating the right-hand side and taking into account $C_0 = \max_Q |c|$ we have

$$\begin{aligned} & \frac{1 - \lambda^2 \beta^2}{2} \sum_{i=1}^n \int_{\Omega} u_{x_i t}^2(x, T) dx + \frac{1 - \lambda^2 \alpha^2}{2} \int_{\Omega} [\Delta u(x, T)]^2 dx + \frac{\varepsilon}{2} \int_Q (\Delta u_t)^2 dxdt \leq \\ & \leq \frac{1}{\varepsilon} \int_Q f^2 + \frac{C_0^2}{\varepsilon} \int_Q u^2 dxdt = I_f. \end{aligned} \tag{10}$$

Applying the Newton-Leibniz formula:

$$u(x, t) = \int_0^t u_{\tau}(x, \tau) d\tau + u(x, 0) \tag{N-L}$$

using the boundary condition (7) we obtain

$$u(x, T) = \frac{1}{1 - \lambda \alpha} \int_0^T u_t(x, t) dt.$$

Using the inequality $|(a + b)^2 \leq 2(a^2 + b^2)|$ we obtain

$$\begin{aligned} u^2(x, t) & \leq 2 \left(\int_0^t u_t(x, t) dt \right)^2 + 2\alpha_1 \left(\int_0^T u_t(x, t) dt \right)^2 \leq \\ & \leq 2 \left(\int_0^t u_t^2 d\tau \right) \left(\int_0^t dt \right) + 2\alpha_1 \left(\int_0^T u_t^2 dt \right) \left(\int_0^T dt \right) \leq (2T + 2\alpha_1 T) \int_0^T u_t^2 dt; \end{aligned}$$

Integrate over Q :

$$\int_Q u^2(x, t) dxdt \leq 2(1 + \alpha_1) T^2 \int_Q u_t^2 dxdt; \tag{11}$$

Here $\alpha_1 = \left(\frac{\lambda \alpha}{1 - \lambda \alpha} \right)^2$.

This is the required inequality. Denote by $k_0 = 2(1 + \alpha_1) T^2$. Further, continuing inequality (11), we have:

$$I_f \leq \frac{1}{\varepsilon} \int_Q f^2 dxdt + \frac{C_0^2 k_0}{\varepsilon} \int_Q u_t^2 dxdt \leq \frac{1}{\varepsilon} \int_Q f^2 + \frac{C_0^2 k_0 m_0}{\varepsilon} \sum_{i=1}^n \int_Q u_{x_i t}^2 dxdt;$$

Using inequality (9), we have

$$I_f \leq \frac{1}{\varepsilon} \int_Q f^2 dxdt + \frac{C_0^2 k_0 m_0}{2\varepsilon} \frac{1}{\varepsilon^2} \int_Q f^2 dxdt = \frac{1}{\varepsilon} \left(1 + \frac{C_0^2 k_0 m_0}{2\varepsilon^2} \right) \int_Q f^2 dxdt;$$

The Newton-Leibniz representation (N-L) using (7) gives:

$$u(x, t) = \int_0^t u_{\tau}(x, \tau) d\tau + \frac{\lambda \alpha}{1 - \lambda \alpha} \int_0^T u_t(x, t) dt;$$

So we have

$$I_1 + I_2 + I_3 \leq \left(\frac{1}{\varepsilon} + \frac{C_0^2 k_0 m_0}{2\varepsilon^3} \right) \int_Q f^2 dxdt; \tag{12}$$

If in (11) we assume $u = \Delta u$, then we obtain:

$$\int_Q (\Delta u)^2 dxdt \leq k_0 \int_Q (\Delta u_t)^2 dxdt \leq \frac{2k_0}{\varepsilon} \left(\frac{1}{\varepsilon} + \frac{C_0^2 k_0 m_0}{2\varepsilon^3} \right) \int_Q f^2 dxdt; \tag{13}$$

It remains to obtain an estimate for the first term, i.e. for u_{tt} . To do this, we use the following well-known inequality:

$$\begin{aligned} |(a_1 + \dots + a_m)^2| &\leq m(a_1^2 + \dots + a_m^2) \\ \int_Q u_{tt}^2 dxdt &= \int_Q (\Delta u - cu + \varepsilon \Delta u_t + f)^2 dxdt \leq (4 \int_Q (\Delta u)^2 + 4C_0^2 \int_Q u^2 + 4\varepsilon \int_Q (\Delta u)^2 + 4 \int_Q f^2) dxdt. \end{aligned}$$

For all terms of the right-hand side of the last inequality, estimates are obtained for fixed ε , i.e.

$$\int_Q u_{tt}^2 dxdt \leq N(\varepsilon) \int_Q f^2 dxdt. \tag{14}$$

Next, let's bonus the continuation method with respect to the parameter for a fixed ε . Now erase λ , i.e. consider the boundary value problem for $\lambda=1$.

$$\begin{cases} u(x, 0) = \alpha u(x, T), \\ u_t(x, 0) = \beta u_t(x, T). \end{cases}$$

Obtaining the first a priori estimate uniform in ε :

$$\begin{aligned} &\frac{1-\beta^2}{2} \int_{\Omega} u_t^2(x, T) dx + \frac{1-\alpha^2}{2} \sum_{i=1}^n \int_{\Omega} u_{x_i}^2(x, T) dx + \\ &+ \frac{1}{2} \int_{\Omega} [c(x, T) - a^2 c(x, 0)] u^2(x, T) dx - \frac{1}{2} \int_{\Omega} c_t u^2 dxdt + \varepsilon \sum_{i=1}^n \int_Q u_{x_i t}^2 dxdt = \\ &= \int_Q f u_t dxdt \leq \frac{\delta^2}{2} \int_Q u_t^2 dxdt + \frac{1}{2\delta^2} \int_Q f dxdt. \end{aligned} \tag{15}$$

Hence, in particular, we have

$$\frac{1-\beta^2}{2} \int_{\Omega} u_t^2(x, T) dx \leq \frac{\delta^2}{2} \int_Q u_t^2 dxdt + \frac{1}{2\delta^2} \int_Q f^2 dxdt. \tag{16}$$

(5) $\times (A - t)u_t$, where A - is some positive number, $A > T$. For example, you can take $A = 2T$, then $A - t > A - T = T > 0$;

$$\begin{aligned} &\int_Q (A - t) u_{tt} u_t dxdt - \int_Q (A - t) \Delta u u_t dxdt + \\ &+ \int_Q (A - t) c u u_t dxdt - \varepsilon \int_Q (A - t) \Delta u_t u_t dxdt = \int_Q (A - t) f u_t dxdt; \end{aligned}$$

Suppose that condition $\frac{\partial}{\partial t}[(A-t)c(x,t)] \leq 0$, is satisfied, then we have

$$\begin{aligned} & \frac{A-T}{2} \int_{\Omega} u_t^2(x,T) dx + \frac{1}{2} \int_Q u_t^2 dxdt + \frac{A-T}{2} \sum_{i=1}^n \int_{\Omega} u_{x_i}^2(x,T) dx + \frac{1}{2} \sum_{i=1}^n \int_Q u_{x_i}^2 dxdt \\ & + \frac{A-T}{2} \int_{\Omega} c(x,T) u^2(x,T) dx - \frac{1}{2} \int_Q ((A-T)c)_t u^2 dxdt + \varepsilon \sum_{i=1}^n \int_Q (A-t) u_{x_i}^2 dxdt = \\ & = \int_Q f(A-t) u_t dxdt + \frac{A}{2} \int_{\Omega} u_t^2(x,0) dx + \frac{A}{2} \sum_{i=1}^n \int_{\Omega} u_{x_i}^2(x,0) dx + \frac{A}{2} \int_{\Omega} c(x,0) u^2(x,0) dx \quad (17) \end{aligned}$$

We use the boundary conditions (7), (8) and (16) we have

$$\int_{\Omega} u_t^2(x,T) dx \leq \frac{\delta^2}{1-\beta^2} \int_Q u_t^2 dxdt + \frac{1}{\delta^2(1-\beta^2)} \int_Q f^2 dxdt.$$

As a result, we get

$$\sum_i I_i \leq K_1 \delta^2 \int_Q u_t^2 dxdt + \frac{K_2}{\delta^2} \int_Q f^2 dxdt,$$

where K_1, K_2 - independent of ε .

We choose, in particular, $K_1 \delta^2 = \frac{1}{4}$, $\delta^2 = \frac{1}{4K_1}$. Then we get

$$\frac{1}{4} \int_Q u_t^2 dxdt + \frac{1}{2} \sum_{i=1}^n \int_Q u_{x_i}^2 dxdt + \varepsilon \sum_{i=1}^n \int_Q u_{x_i}^2 dxdt \leq 4K_1 K_2 \int_Q f^2 dxdt.$$

Hence, in particular, we have

$$\int_Q u_t^2 dxdt \leq 16K_1 K_2 \int_Q f^2 dxdt, \quad \sum_{i=1}^n \int_Q u_{x_i}^2 dxdt \leq 8K_1 K_2 \int_Q f^2 dxdt, \quad \varepsilon \sum_{i=1}^n \int_Q u_{x_i}^2 dxdt \leq 4K_1 K_2 \int_Q f^2 dxdt.$$

Adding all the terms, we get

$$\int_Q u_t^2 dxdt + \sum_{i=1}^n \int_Q u_{x_i}^2 dxdt + \varepsilon \sum_{i=1}^n \int_Q u_{x_i}^2 dxdt \leq K_0 \int_Q f^2 dxdt,$$

where K_0 - is independent of ε . As required.

Thus, the following theorem is proved.

Theorem. Let the conditions be satisfied: $c(x,T) - \lambda^2 \alpha^2 c(x,0) \geq 0$, $c_t(x,T) \leq 0$, $f, f_t \in L_2(Q)$. Then the boundary value problem (1) - (4) cannot have more than one solution in space $W_2^{2,2}(Q)$.

Funding. This work was supported by grant AP05135319 of the Ministry of Education and Science of the Republic of Kazakhstan.

Б.Д. Қошанов¹, Г.Д. Қошанова², Г.Е. Әзімхан³, Р.У. Сегізбаева⁴

¹Математика және математикалық моделдеу институты, Алматы, Қазақстан;

^{1,3}Абай атындағы Қазақ ұлттық педагогикалық университеті, Алматы, Қазақстан;

³Х.А. Яссауи атындағы Халықаралық Қазақ-Түрік университеті, Түркістан, Қазақстан;

⁴Азаматтық авиация академиясы, Алматы, Қазақстан

КӨПӨЛШЕМДІ ШЕКТЕЛГЕН ОБЛЫСТА ГИПЕРБОЛАЛЫҚ ТЕНДЕУЛЕР ҮШІН ЛОКАЛЬДЫ ЕМЕС ШЕТТІК ЕСЕПТЕРДІҢ ШЕШІМДІЛІГІ

Аннотация Бұл жұмыста шектелген көпөлшемді цилиндрлі облыста $Q = \Omega \times (0, T)$, $\Omega \subset R^n$, $S = \Gamma \times (0, T)$, $\Gamma = \partial\Omega$ $0 < t < T < +\infty$ келесі гиперболалық теңдеу үшін

$$u_{tt} - \Delta u + c(x, t)u = f(x, t), \quad (x, t) \in Q \quad (1)$$

келесі түрдегі локалды емес шеттік есеп

$$u(x, t)|_S = 0, \quad (2)$$

$$u(x, 0) = \alpha u(x, T), \quad (3)$$

$$u_t(x, 0) = \beta u_t(x, T) \quad (4)$$

карастырылады, мұндағы $f(x, t) - x \in \bar{\Omega}$, $t \in [0, T]$ берілген функциялар $\alpha, \beta \in R$ - сандар.

Азғындалған квазигиперболалық теңдеулер үшін шеттік есептердің шешімділігін зерттеу В.Н. Врэгтың жұмыстарынан бастау алады.

Сызықты параболалық теңдеулер үшін интегралдық шарты бар, тақ ретті дифференциалдық теңдеулер үшін, стационарлы емес кейбір теңдеулердің класында локалды емес шеттік есептердің шешімділігінің мәселесі, соңғы жылдары, А.И. Кожановтың жұмыстарында зерттелген.

$$u_{tt} - \Delta u + c(x, t)u - \varepsilon \Delta u_t = f(x, t) \in L_2(Q) \quad (5)$$

$$u(x, t)|_S = 0, \quad (6)$$

$$u(x, 0) = \lambda \alpha u(x, T), \quad (7)$$

$$u_t(x, 0) = \lambda \beta u_t(x, T) \quad (8)$$

(1)-(4) есебінің регулярлы шешімін зерттеу үшін қосалқы (5)-(8) регулярланған есебін қарастырамыз. Осы алынған қосалқы (5)-(8) есеп үшін: λ параметрі бойынша жалғастыру әдісін, сәйкес функционалды кеңістіктерде априорлы бағалаулар әдістерін қолданамыз. ε -нан бірқалыпты тәуелді бағалауларда $\varepsilon \rightarrow 0$ ұмтылдырыу арқылы, зерттеліп отырған есептің регулярлы шешімінің бар болуы және жалғыздығы туралы теорема дәлелденеді.

Түйін сөздер: гиперболалық теңдеу, шеттік есептер, регулярлы шешімдер, бар болуы, жалғыздығы.

Б.Д. Кошанов¹, Г.Д. Кошанова², Г.Е. Азимхан³, Р.У. Сегізбаева⁴

¹Институт математики и математического моделирования, Алматы, Казахстан;

^{1,3}Казахский национальный педагогический университет им. Абая, Алматы, Казахстан;

²Международный Казахско-турецкий университет имени Х.А. Ясави, Туркестан, Казахстан;

⁴Академия гражданской авиации, Алматы, Казахстан

РАЗРЕШИМОСТЬ КРАЕВЫХ ЗАДАЧ С НЕЛОКАЛЬНЫМИ УСЛОВИЯМИ ДЛЯ МНОГОМЕРНЫХ ГИПЕРБОЛИЧЕСКИХ УРАВНЕНИЙ

Аннотация. В данной работе исследуется разрешимость новых нелокальных краевых задач для гиперболических уравнений в многомерной ограниченной области.

В ограниченной цилиндрической области $Q = \Omega \times (0, T)$, $\Omega \subset R^n$, $S = \Gamma \times (0, T)$, $\Gamma = \partial\Omega$ $0 < t < T < +\infty$ рассмотрим гиперболическое уравнение

$$u_{tt} - \Delta u + c(x,t)u = f(x,t), \quad (x,t) \in Q \quad (1)$$

со следующими нелокальными по времени краевыми условиями

$$u(x,t)|_S = 0, \quad (2)$$

$$u(x,0) = \alpha u(x,T), \quad (3)$$

$$u_t(x,0) = \beta u_t(x,T) \quad (4)$$

где $c(x,t), f(x,t)$ – заданные функции, определенные при $x \in \bar{\Omega}, t \in [0, T], \alpha, \beta \in R$ – некоторые числа.

Исследование разрешимости краевых задач для вырожденных квазигиперболических уравнений (по аналогии с квазиэллиптическими уравнениями) началось с работ В.Н. Врагова.

Исследования нелокальных задач с интегральными условиями для линейных параболических уравнений, для дифференциальных уравнений нечетных порядков и для некоторых классов нестационарных уравнений в последнее время активно исследуются в работах А.И. Кожанова.

Для исследования нелокальной краевой задачи (1)-(4) рассмотрим следующую регуляризованную задачу

$$u_{tt} - \Delta u + c(x,t)u - \varepsilon \Delta u_t = f(x,t) \in L_2(Q) \quad (5)$$

$$u|_S = 0 \quad (6)$$

$$u(x,0) = \lambda \alpha u(x,T), \quad (7)$$

$$u_t(x,0) = \lambda \beta u_t(x,T). \quad (8)$$

Для этой регуляризованной задачи (5)-(8) мы применим метод продолжения по параметру. Для задачи (5)-(8) получим априорные оценки в соответствующих функциональных пространствах. И в конце переходим к пределу $\varepsilon \rightarrow 0$.

Таким образом, для изучаемой задачи доказывается теорема существования и единственности регулярного решения.

Ключевые слова: гиперболическое уравнение, краевые задачи, регулярные решения, существование, единственность.

Information about authors:

Koshanov Bakhytbek Danebekovich - Institute of Mathematics and Mathematical Modeling, Chief scientific researcher, Abai Kazakh National Pedagogical University, Doctor of Physical and Mathematical Sciences, professor, koshanov@list.ru <https://orcid.org/0000-0002-0784-5183>;

Koshanova Gulnazira Danebekovna - Senior lecturer, Yasavi International Kazakh-Turkish University, Candidate of Pedagogical Sciences, gulnazira.koshanova@ayu.edu.kz, <https://orcid.org/0000-0002-5990-1490>;

Azimkhan Gulsara Yerbolatkyzy - Abai Kazakh National Pedagogical University, magistr, g_azimhanova@mail.ru, <https://orcid.org/0000-0001-7130-4413>;

Segizbayeva Rabigash Umirzakovna - Civil Aviation Academy, Candidate of Physical and Mathematical Sciences, Associate professor, segizbaeva-55@mail.ru

REFERENCES

[1] Enemies V.N. On the theory of boundary value problems for equations of mixed type // Differ. equations. 13: 6 (1977), 1098-1105.

[2] Enemies V.N. On the formulation and solvability of boundary value problems for equations of mixed - composite type, Mathematical analysis and related problems of mathematics, Nauka, Novosibirsk, 1978, 5–13.

[3] Egorov I.E., Fedorov V.E. Nonclassical equations of higher order mathematical physics, Computing Center SB RAS, Novosibirsk, 1995.

[4] Terekhov A.N. A boundary value problem for an equation of mixed type. Application of functional analysis methods to solving problems of mathematical physics and computational mathematics, Sat. scientific tr., Institute of mathematics. SB USSR Academy of Sciences, Novosibirsk, 1979, 128-137.

[5] Egorov I.E., Zakharova T.I. On the Fredholm property of a boundary value problem for an equation of mixed type // Math. Notes of YSU, 20:1 (2013), 20 - 26.

[6] Egorov I.E. On a boundary value problem for an equation of mixed type with a spectral parameter // Matem. notes NEFU, 21: 1 (2014), 11-17.

- [7] Kozhanov A.I., Sharin E.F. The conjugation problem for some nonclassical differential equations of high order // *Ukrain. mate. visnik*, **11:2** (2014), 181-202.
- [8] Pinigina N.R. On the question of the correctness of boundary value problems for non-classical differential equations of high order // *Asian - European Journal of Mathematics*, **10:3** (2017), 7-17.
- [9] Kozhanov A.I., Pinigina N.R. Boundary value problems for nonclassical differential equations of higher order // *Math. deputy.*, 2017.
- [10] Kozhanov A.I., Koshanov B.D., Sultangazieva Zh.B. New boundary value problems for a fourth-order quasi-hyperbolic type // *Siberian Electronic Math. News*, **16**, (2019), 1410-1436.
- [11] Koshanov B.D., Sultangazieva Zh.B., Emir Kady oglu A.N., Smatova G.D. The spectral problem for nonclassical differential equations of the sixth order // *Bulletin of the Karaganda University, series Mathematics*, **97:1** (2020), 17-25.
- [12] Kalmenov T.Sh., Iskakova U.A. A criterion for the strong solvability of the mixed Cauchy problem for the Laplace equation // *Doklady RAS*, **414: 2**, (2007), 455-458.
- [13] Kal'menov T.Sh., Koshanov B.D. Representation for the Green's function of the Dirichlet problem for polyharmonic equations in a ball. *Siberian Mathematical Journal*. **49:3** (2008), 423-428.
- [14] Kal'menov T.Sh., Koshanov B.D., Nemchenko M.Yu. Green function representation in the Dirichlet problem for polyharmonic equations in a ball // *Doklady Mathematics*, **78:1** (2008), 528-530.
- [15] Kal'menov T.Sh., Koshanov B.D., Nemchenko M.Yu. Green function representation for the Dirichlet problem of the polyharmonic equation in a sphere // *Complex Variables and Elliptic Equations*, **53:2** (2008), 177-183.
- [16] Koshanov B.D., Edil K. The Green function of the Dirichlet problem for a biharmonic equation and a polynomial solution for the Poisson equation in a circle // *Bulletin of the NAS of the Republic of Kazakhstan, Series of Physics and Mathematics*, **3:307**, (2016), 62-72.
- [17] Sadybekov M.A., Torebek B.T., Turmetov B.Kh. On an explicit form of the Green function of the third boundary value problem for the Poisson equation in a circle, *AIP Conference Proceedings*, **1611**, (2014), 255-260.
- [18] Sadybekov M.A., Torebek B.T., Turmetov B.Kh. Representation of Green's function of the Neumann problem for a multi-dimensional ball, *Complex Variables and Elliptic Equations*, **61:1** (2016), 104-123.
- [19] Kanguzhin B.E., Koshanov B.D. Necessary and sufficient conditions for the solvability of boundary value problems for polyharmonic equation // *Ufa Mathematical Journal*, **52:2** (2010), 41-52.
- [20] Koshanov B.D., Soldatov A.P. The boundary value problem with normal derivatives for an elliptic equation in the plane // *Differ. Equations*, **52:12** (2016), 1666-1681. DOI: 10.1134/S03740641160
- [21] Koshanov B.D., Adilbekov E.N., Duysen E. Dimension of the space of solutions of the Dirichlet problem for the Poisson equations and the biharmonic equation in an unbounded domain - I // *Bulletin of the NAS of the Republic of Kazakhstan, Series of Physics and Mathematics*, **2:312** (2017), 116-125.
- [22] Koshanov B.D., Adilbekov E.N., Duysen E. Dimension of the space of solutions of the Dirichlet problem for the Poisson equations and the biharmonic equation in an unbounded domain - II // *Bulletin of the NAS of the Republic of Kazakhstan, Series of Physics and Mathematics*, **3:313** (2017), 126-131.
- [23] Koshanov B.D., Nurikenova J. On solvability of the generalized Dirichlet-Neiman problem for a high order elliptic equation // *Bulletin of the NAS of the Republic of Kazakhstan, Series of Physics and Mathematics*, **2:312** (2017), 95-103.
- [24] V.S. Vladimirov, *Equations of Mathematical Physics*, Nauka, M., 1988.
- [25] V.A. Trenogin, *Functional Analysis*, Science, M., 1980.
- [26] O.A. Ladyzhenskaya, N.N. Uraltseva, *Linear and quasilinear equations of elliptic type*, Nauka, Moscow, 1973.