

ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

2020 • 6

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944



ALMATY, NAS RK

Б а с р е д а к т о р ы
х.ғ.д., проф., ҚР ҰҒА академигі
М.Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

Адекенов С.М. проф., академик (Қазақстан) (бас ред. орынбасары)
Бенберин В.В., проф., академик (Қазақстан)
Березин В.Э., проф., корр.-мүшесі (Қазақстан)
Величкин В.И. проф., корр.-мүшесі (Ресей)
Вольдемар Вуйцик проф. (Польша)
Елешев Р.Е., проф., академик (Қазақстан)
Жамбакин Қ.Ж., проф., академик (Қазақстан)
Иванов Н.П., проф., академик (Қазақстан)
Илолов М.И. проф., академик (Тәжікстан)
Кригер Виктор проф. (Германия)
Кененбаев С.Б., проф., академик (Қазақстан)
Леска Богуслава проф. (Польша)
Локшин В.Н. проф., академик (Қазақстан)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Нургожин Т.С., проф., корр.-мүшесі (Қазақстан)
Перни Стефано проф. (Ұлыбритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Ұлыбритания)
Рамазанов Т.С. проф., академик (Қазақстан)
Раманкулов Е.М., проф., корр.-мүшесі (Қазақстан)
Садыкулов Т., проф., академик (Қазақстан)
Семенов В.Г., проф., академик (Россия)
Сикорски Марек проф., (Польша)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Уразалиев Р.А., проф., академик (Қазақстан)
Харин С.Н. проф., академик (Қазақстан)
Харун Парлар проф. (Германия)
Чечин Л.М. проф., корр.-мүшесі (Қазақстан)
Энджун Гао проф. (Қытай)

«Қазақстан Республикасы Ұлттық ғылым академиясының баяндамалары»

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.).

Қазақстан Республикасының Ақпарат және қоғамдық даму министрлігінің Ақпарат комитетінде 29.07.2020 ж. берілген № KZ93VPY00025418 мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *наноматериалдар алу, биотехнология және экология саласындағы бірегей зерттеу нәтижелерін жариялау.*

Мерзімділігі: жылына 6 рет.

Тиражы: 500 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19, 272-13-18,
<http://reports-science.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор
д.х.н., проф., академик НАН РК
М. Ж. Журинов

Редакционная коллегия:

Адекенов С.М. проф., академик (Казахстан) (зам. гл. ред.)
Бенберин В.В., проф., академик (Казахстан)
Березин В.Э., проф., чл.-корр. (Казахстан)
Величкин В.И. проф., чл.-корр. (Россия)
Вольдемар Вуйцик проф. (Польша)
Елешев Р.Е., проф., академик (Казахстан)
Жамбакин К.Ж., проф., академик (Казахстан)
Иванов Н.П., проф., академик (Казахстан)
Илолов М.И. проф., академик (Таджикистан)
Кригер Виктор проф. (Германия)
Кененбаев С.Б., проф., академик (Казахстан)
Леска Богуслава проф. (Польша)
Локшин В.Н. проф., академик (Казахстан)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Нургожин Т.С., проф., чл.-корр. (Казахстан)
Перни Стефано проф. (Великобритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Великобритания)
Рамазанов Т.С. проф., академик (Казахстан)
Раманкулов Е.М., проф., чл.-корр. (Казахстан)
Садыкулов Т., проф., академик (Казахстан)
Семенов В.Г., проф., академик (Россия)
Сикорски Марек проф., (Польша)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Уразалиев Р.А., проф., академик (Казахстан)
Харин С.Н. проф., академик (Казахстан)
Харун Парлар проф. (Германия)
Чечин Л.М. проф., чл.-корр. (Казахстан)
Энджун Гао проф. (Китай)

Доклады Национальной академии наук Республики Казахстан»

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и общественного развития Республики Казахстан № **KZ93VPY00025418**, выданное 29.07.2020 г.

Тематическая направленность: *публикация оригинальных результатов исследований в области получения наноматериалов, биотехнологии и экологии.*

Периодичность: 6 раз в год.

Тираж: 500 экземпляров

Адрес редакции: 050010, г.Алматы, ул.Шевченко, 28; ком. 219, 220; тел. 272-13-19, 272-13-18,

<http://reports-science.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2020 г.

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

M.Zh. Zhurinov

E d i t o r i a l b o a r d :

Adekenov S.M. prof., academician (Kazakhstan) (deputy editor in chief)
Benberin V.V., prof., academician (Kazakhstan)
Berezin V.Ye., prof., corr. member. (Kazakhstan)
Velichkin V.I. prof., corr. member (Russia)
Voitsik Valdemar prof. (Poland)
Eleshev R.E., prof., academician (Kazakhstan)
Zhambakin K.Zh., prof., academician (Kazakhstan)
Ivanov N.P., prof., academician (Kazakhstan)
Iolov M.I. prof., academician (Tadjikistan)
Krieger Viktor prof. (Germany)
Kenenbayev S.B., prof., academician (Kazakhstan)
Leska Boguslava prof. (Poland)
Lokshin V.N. prof., academician (Kazakhstan)
Nekludov I.M. prof., academician (Ukraine)
Nur Izura Udzir prof. (Malaysia)
Nurgozhin T.S., prof., corr. member. (Kazakhstan)
Perni Stephano prof. (Great Britain)
Potapov V.A. prof. (Ukraine)
Prokopovich Polina prof. (Great Britain)
Ramankulov E.M., prof., corr. member. (Kazakhstan)
Sadykulov T., prof., academician (Kazakhstan)
Semenov V.G., prof., academician (Russia)
Sikorski Marek prof., (Poland)
Ramazanov T.S. prof., academician (Kazakhstan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief
Urazaliev R.A., prof., academician (Kazakhstan)
Kharin S.N. prof., academician (Kazakhstan)
Kharun Parlar prof. (Germany)
Chechin L.M. prof., corr. member (Kazakhstan)
Endzhun Gao prof. (China)

Reports of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2224-5227

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Social Development of the Republic of Kazakhstan No. **KZ93VPY00025418**, issued 29.07.2020.Thematic scope: *publication of original research results in the field of obtaining nanomaterials, biotechnology and ecology.*

Periodicity: 6 times a year.

Circulation: 500 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://reports-science.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

REPORTS OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ISSN 2224-5227

<https://doi.org/10.32014/2020.2518-1483.132>

Volume 6, Number 334 (2020), 29 – 34

UDC 637.334; 579.61

MRSTI 68.05.31

A. U. Issayeva^{1*}, B. Leska², A. A. Uspabaeva³, A. Ye. Tleukeeva³, A. Abubakirova³

¹ Shymkent University, Shymkent, Kazakhstan;

² A. Mickiewicz Poznań State University, Poznań, Poland;

³ M. Auezov South Kazakhstan University, Shymkent, Kazakhstan.

E-mail: akmaral.issayeva@bk.ru, bogunial@amu.edu.pl, aseltleukeeva@mail.ru,
uspabaeva73@mail.ru, azhar.baikal79@mail.ru

BIODIVERSITY OF THE MICROFLORA OF THE SALT LAKES DZHAKSY-KLYCH AND BUGA-DZHAILY

Abstract. The article illustrates the results of microbiological and chemical studies of samples of salt lakes in the south of Kazakhstan, such as Dzhaksy-Klych and Buga-Dzhaily. It was found that the samples from the bottom of the dried up Small Aral Sea are dominated by ions of magnesium, potassium, calcium, iron and copper. Moreover, the presence of pesticides was revealed: acetamiprid, fenhexamide, folpet-like compounds, tetrahydroptolamide, which are used in the agro-industry. Samples of Lake Dzhaksy-Klych were isolated from different layers of halite, magnesium, and sulfate salts. Salt-containing raw material of Lake Buga-Dzhaily contains 96.0 ± 8.9 wt.% NaCl, the content of magnesium ions ranges from 0.17 ± 0.01 - 4.01 ± 0.0 wt.%, The presence of sulfur is within $0,26 \pm 0.02\%$. 1.71 ± 0.12 wt%. The amount of heavy metals is below the permissible concentration. It was found that the increased content of magnesium in the samples proves the presence of mixed sodium-magnesium layers. The number of heterotrophic microorganisms in the samples of salt lakes is approximately 103-104 CFU/g, enterobacteria -103 CFU / g, micromycetes in the amount of $(6.0 \pm 0.5) \times 10^3$ CFU/g were found only in the sample of sulfate salts and silt sampled at a distance of 1.0 m from the coastline. The distribution and density of microflora on the coastline indicate an increased anthropogenic factor due to salt extraction.

Key words: salt lakes, salt-containing raw materials, microflora, heterotrophic microorganisms, micromycetes, pesticides.

Introduction. Continental salt water bodies differ markedly in chemical and microbiological composition, while the seas are similar in this composition. The composition of these waters directly depends on the geological, climatic and biogeochemical conditions and can be neutral, alkaline and acidic. Salt water bodies are poor in the presence of microorganisms. Moreover, the variety of these organisms is limited. Continental salt water bodies are often confined to areas with an arid climate, for example, in the south of Kazakhstan.

Numerous studies show a low level of biodiversity of microorganisms, which are widespread in the water itself and the coastal area. Most organisms, adapting to extreme conditions, begin to adapt at the cellular and functional levels [1,2]. Such changes have been investigated for further industrial application [3].

According to the chemical composition of some sodium salts, salinization of water can be sulfate, chloride, soda or mixed, with the most harmful effects of Na⁺ and Cl⁻ ions. The salinity of water is the main factor in the adaptation of aquatic organisms to environmental conditions. Adaptive mechanisms are based on the regulation of the concentration of ions in the intracellular fluid and its osmotic pressure [4,5]. The concentration of inorganic ions in the aquatic environment can vary over a very wide range, which leads to an increase in the ionic gradient between the body and the environment.

The maintenance of osmotic pressure inside the cell under conditions of different salinity of the aquatic environment is carried out by two mechanisms: a change in the concentration of organic osmotically active substances in the cytoplasm and a change in the content of inorganic ions in it [6,7]. The molecular aspects of adaptation of prokaryotes to changes in the salinity of the aquatic environment have been established [8]. According to [9, 10], NaCl in concentrations higher or lower than optimal, but still within acceptable limits, can play the role of a stress factor that causes morphophysiological and behavioral changes in organisms. The influence of salinization of water bodies on the stability and change in the structure of the population of organisms of aquatic organisms has been established [11,12,13].

On the other hand, in studies by Loukas et al. shows that the salinity level of the reservoir is a less significant limiting factor than expected. The formation of biocenoses of salt water bodies is influenced by a combination of factors such as pH of the environment, oxygen content and ionic composition of water, anthropogenic impact, hydrological and geographical characteristics of the reservoir, and paleoclimatic parameters [14,15,16, 17]. The influence of climatic conditions on the landscape is shown in the work on the basis of a model of biological and ecological productivity of landscape systems of the South Kazakhstan region, including Almaty, Zhambyl, Turkestan and Kyzylorda regions using long-term data from 32 meteorological stations located in the region [18].

In this regard, the purpose of the study was to study the effect of saline raw materials on the biodiversity of microflora in the Dzhaksy-Klych and Buga-Dzhaily lakes located in the south of Kazakhstan.

Objects and methods of research. As a research objects were used salt-containing raw from lakes Dzhaksy-Klych and Buga-Dzhaily: halite, halite-sulfate, sulphate, sulphate-magnesium salt, brine, surface brine (halite layer), brine (sulfate formation) etc.

Lake Dzhaksy-Klych –the biggest salt lake of the Caspian lowlands located in the Aral sea region. The lake is of marine origin and consists of two parts with an area of 18 and 58 km. The thickness of the salt deposit is about 2 m.

Lake Buga-Dzhaily is one of the largest lakes in the system of salt lakes located in the Suzak district of Turkestan region. Every year, up to 150 thousand tons of table salt is extracted by open method.

Sampling was carried out in accordance with GOST 33770-2016. Samples were taken from the Dzhaksy-Klych Deposit by “ONYX-R” LLP and the authors of the article; from the Buga-Dzhaily Deposit- by “As-Dinar” LLP.

Microscopic studies were performed using "Tauda" and "Mikmed-5" microscopes, light microscopy methods.

Microorganisms were isolated and cultured on the appropriate nutrient media: heterotrophs - on MPA, micromycetes - on Chapek medium, enterobacteria-on Endo-Ploskirev medium. Haloresistance of isolated cultures of microorganisms was determined by their ability to grow at various concentrations of NaCl from 0 to 20%.

In the study of sulfur-containing raw materials, a mass spectrometer with inductively coupled plasma was used, the elements were determined in accordance with ST RK ISO 17294-2-2006. Chemical analysis was carried out according to GOST 13685-84. Research conditions: temperature-250C; humidity -83.0%; pressure-714 mm Hg.

Statistical processing of the results was performed by calculating the arithmetic mean and the standard deviation. All determinations were carried out in 3-and 5-fold repetitions. The data was processed using an IBM Pentium personal computer based on Excel application software packages.

Research results and discussion. To use salt-containing raw materials from two salt lakes in the South of Kazakhstan as bases for cosmetology products, it was necessary to make sure that they were harmless to human health. In this regard, physical, chemical and microbiological studies were carried out. In previous studies (Aladin et al., 2008), it was found that the content of sulfates in the Aral sea salts exceeds 31% (of the total amount of salts), and sodium chloride is only 54%. The ionic composition of the salts is as follows: sodium - from 2.83 to 13.73%; sulfate-ion-7.5-30.14; calcium - up to 1.08; magnesium-3.03; potassium-0.93; carbonate-ion-0.18; chlorine-2.09; water - up to 55.23%. The analysis of the elemental composition of salt samples taken during expedition trips in 2019 from the dried bottom of the Small Aral sea revealed the predominance of magnesium, potassium, calcium, iron and copper ions in them (table 1). The chemical composition of the Aral salts belongs to the chloride group, although in some cases there are also deposits of sulfide compounds, while the proportion of sodium chloride in samples from 15 sampling points is 89.98%, and sulfates-no more than 2.0%.

Table 1 – Chemical composition of salts of the bottom of the Small Aral sea (main components)

Elements	Mas. %	mkg/l
B	0,001039747	593,7200
Mg	0,046351029	63073,6600
Al	0,003912761	78,0350
P	0,005430454	203,2210
K	0,018169232	11514,1000
Ca	0,236304506	67786,4576
Ti	0,000126723	5,2341
Fe	0,025324563	1263,2342
Cu	0,002123332	2132,5411
Zn	0,002343253	3,1234
Ag	0,000045364	2,2534
Co	0,000012432	0,3243
Ba	0,0006243001	7,2364
Cr	0,0034524321	32,4323
Ni	0,000254364	4.6300
Mo	0.0000354231	15,6534
Na ₂ SO ₄	1,89	1,98 g/l
NaCl	89,98	1188,98 g/l

Due to the fact that the Aral sea region is characterized as an agro-industrial region where a variety of pesticides are widely used for agricultural production, analyses were conducted for the content of pesticides in samples of salt-containing samples. In the analyzed samples was not detected biphenyl, diphenylamine, but is revealed acetamiprid, phenhexamid, pholpet-like compositions, tetrahydrofolate. In total, about 400 active substances were searched on LC-MS/MS and GC-MS/MS. Almost all compounds were found on GC-MS/MS, acetamiprid on LC-MS/MS. Comparing the detector responses (peak size), by far the largest residue was found for phengexamide (a fungicide) used to protect against the gray mold *Botrytis cinerea* in crops including strawberries, raspberries, or blueberries. Other substances are found only in trace amounts.

In the average samples of lake Buga-Dzhaily salt, the pH of the salt solution did not exceed 6.34 ± 0.5 . No lead, arsenic, zinc, or iron were found in the samples. Chemical analysis of averaged samples of salt-containing raw materials showed that the NaCl content in the feed salt ranges from 96.68-98.06 wt.%; in the food salt-92.88 – 93.94 wt.%; in the brine -26.13 ± 0.01 wt.%; in the sedimentary silt -11.02 wt.%. Elemental analysis showed the presence of magnesium ions, the content of which in the feed salt is in the limit, wt.% -0.17 ± 0.01 ; in the food salt -1.87 ± 0.11 ; brine -4.01 ± 0.0 ; sludge -0.79 ± 0.0 . In addition, all analyzed samples revealed the presence of sulfur, the amount of which in the samples of feed salt was in the range, wt.% -0.26 ± 0.02 ; in brine -1.71 ± 0.12 ; in silt sediment -0.49 ± 0.0 ; in food salt, an increase in the sulfur content to 0.98 ± 0.05 was noted. The content of other elements, including heavy metal ions, is below the safety limit. Analysis of the elemental composition of sample salt samples showed that in addition to sodium chloride and sulfate (0.71-1.05% wt.) and insoluble residue (0.07-0.28% wt.) in decimal and hundredths of % wt. boron, magnesium, potassium, and calcium are noted. The prevalence of magnesium in these samples shows the probability of mixed sodium-magnesium layers in the Buga-Dzhaily deposit.

Microscopy of samples of surface brine and brine taken from the sulfate layer of lake Dzhaksy-Klych showed the absence of living objects. Crystals of various sizes and cuboidal shapes were found. In the brine taken from the halite formation from a depth of 0.3 m, in addition to cuboidal forms of salt crystals, amorphous formations and a few amoeboid structures were found. The microflora is represented along with small coccoid and rod-shaped motile bacteria and clusters of small convoluted motile microorganisms noted earlier in the studies of E. G. Dobrynin(1984) of the Eupatoria solprom garden brine.

As a result of microbiological studies, it was found that the number of heterotrophic microorganisms in the samples taken from lake Dzhaksy-Klych ranges from 10^4 CFU/g. However, the number of heterotrophic microorganisms increases to 10^5 CFU/g in samples of halite salt samples taken at a distance of 3-5 m from the coastline. The smallest number of microorganisms was found in samples of sulfate salts,

where the number of heterotrophic microorganisms was within the range of $(4.2 \pm 0.3) \times 10^3$ CFU/g. The number of enterobacteria in all samples ranges from 10^3 CFU/g. Micromycetes in the amount of $(6.0 \pm 0.5) \times 10^3$ CFU / g were found only in a sample of sulfate salts and silt taken at a distance of 1.0 m from the coastline.

In samples of salt-containing raw materials of lake Buga - Dzhaily, the number of heterotrophic microorganisms was in the range of 10^4 CFU/g, enterobacteria- 10^3 CFU/g. In almost all samples, micromycetes of 10^3 CFU/g were detected, with the exception of samples of deep silt and halite salt taken from a depth of 1.0 m from the center of the lake. From the studied samples, taking into account the morphological and cultural properties of microorganisms, 10 cultures of microorganisms were isolated, the dominant part of which consisted of representatives of the genera-*Micrococcus*, *Bacillus*, *Pseudomonas*. The genus *Micrococcus* is represented by three dominant species-*M.luteus*, *M. roseus*, *Micrococcus sp.* Enterobacteria are represented by the species – *Enterobacter sp.* Micromycetes are represented by the genera *Aspergillus*, *Mucor*, *Penicillium*, *Fusarium*. The presence of these taxonomic groups of microorganisms and the nature of the distribution of microflora indicates the contamination of the surface of salt-containing raw materials as a result of human activity.

Haloresistance of isolated cultures of microorganisms was determined by their ability to grow at various concentrations of NaCl from 0 to 20%. As follows from the presented data, it was found that *M. roseus* grows intensively on a medium with 3-10% NaCl, good growth is observed on a medium with 13% NaCl, and there is no growth on a medium with 17-25% NaCl. *Micrococcus sp.* it grows intensively on medium with 3-5% NaCl, on medium with 10-17% NaCl it grows well, and on medium with 20-25% NaCl there is no growth. *M. luteus* and *Bacillus sp.* they grow intensively on a medium with 0-17% NaCl, on a medium with 20% NaCl, good growth is observed, with an increase in the percentage of NaCl to 25%, there is no growth. Cultures of *Pseudomonas sp.* and *Enterobacter sp.* grow intensively on medium with 3-10% NaCl, with an increase in the concentration of NaCl to 13-17% culture *Pseudomonas sp.* it grows well, with a further increase in the salt concentration to 20% or higher, the growth of this culture stops, and the growth of *Enterobacter sp.* it is suppressed already on the medium with 17-25% NaCl. *Aspergillus sp.* it grows intensively on a medium with 3-13% NaCl, on a medium with 17-20% NaCl there is good growth, and on a medium with 25% NaCl there is no growth. *Mucor sp.* on the medium with 3-5% NaCl, good growth is observed, in other variants, the growth of this culture is absent. On a medium with 3-10% NaCl *Penicillium sp.* it grows intensively, and good growth is observed in the environment with 13-17% NaCl. *Fusarium sp.* it grows intensively on medium with 3-5% NaCl, with an increase in the percentage of NaCl by 10-13%, there is a good growth, and with an increase from 17% to 25%, the growth is completely absent.

Conclusion. Thus, the results of physico-chemical and microbiological studies showed that the salt-containing raw materials of the Dzhaksy-Klych and Buga-Dzhaily deposits contain 96.0 ± 8.9 wt.% NaCl, the content of magnesium ions ranges from 0.17 ± 0.01 - 4.01 ± 0.0 wt.%, the presence of sulfur is noted in the range of 0.26 ± 0.02 - 1.71 ± 0.12 wt.%. the Content of other elements, including heavy metal ions, is below the safety level, in decimal and hundredths of % wt. boron, potassium and calcium are noted.

It was found that the number of heterotrophic microorganisms ranges from 10^3 - 10^4 CFU/g, enterobacteria - 10^3 CFU/g, and micromycetes were found only in samples of salt-containing raw materials from the Buga-Dzhaily deposit. The nature of the distribution and abundance of microflora indicates the anthropogenic presence in the area of salt production. It was found that the cultures of *M. luteus*, *Bacillus sp.*, *Aspergillus sp.*, *Penicillium sp.* and *Fusarium sp.* isolated from salt-containing substrates are resistant to 17-20% NaCl.

Acknowledgement.

The work was carried out under the grant of the Ministry of education and science of the Republic of Kazakhstan AR05131728: "Development of production technology and obtaining prototypes of new cosmetic products based on pharmacological studies of domestic salt-containing and vegetable raw materials" (2018-2020).

А.Ө. Исаева^{1*}, Б. Леска², А.А. Успабаева³, А.Е. Тлеукеева³, А. Абубакирова³

¹ Шымкент университеті, Шымкент, Қазақстан;

² А.Мицкевич атындағы Познань мемлекеттік университеті, Познань, Польша;

³ М.Әуезов атындағы Оңтүстік Қазақстан университеті, Шымкент, Қазақстан

ЖАҚСЫ ҚЫЛЫШ ЖӘНЕ БҰҒА-ЖАЙЛЫ ТҰЗДЫ КӨЛ МИКРОФЛОРАСЫНЫҢ БИОЛОГИЯЛЫҚ ӘРТҮРЛІЛІГІ

Аннотация. Мақалада Қазақстанның оңтүстігіндегі Жақсы қылыш және Бұға-жайлы сынды тұзды көл үлгілерін микробиологиялық және химиялық зерттеу нәтижелері көрсетілген.

Континентальды тұзды су айдындары химиялық және микробиологиялық құрамы бойынша айтарлықтай ерекшеленеді, ал теңіздер де құрамы бойынша ұқсас келеді. Судың құрамы тікелей геологиялық, климаттық және биогеохимиялық жағдайға байланысты бейтарап, сілтілі және қышқыл болуы мүмкін. Тұзды су объектілерінде микроорганизмдер популяциясы өте төмен және таралуы тұздың концентрациясына тікелей қатысты.

Адаптивті механизмдер жасушаішілік сұйықтықтағы ион концентрациясын және оның осмотық қысымын реттеуге негізделген. Су ортасындағы бейорганикалық ион концентрациясы кең ауқымда өзгеруі мүмкін, бұл организм мен қоршаған орта арасындағы ионды градиентті жоғарылатады. Қазақстанның оңтүстігіндегі екі тұзды көлден алынған тұзды шикізатты косметикаға негіз ретінде пайдалану үшін оның адам денсаулығына зиянсыз екендігіне көз жеткізу мәселесін алға қойдық. Осыған байланысты физикалық, химиялық және микробиологиялық зерттеулер жүргізілді.

Құрғаған Арал теңізінің түбінен алынған сынамаларда магний, калий, кальций, темір және мыс иондары басым екендігі анықталды. Сонымен қатар, пестицидтердің: агроөнеркәсіпте қолданылатын ацетамиприд, фенгексамид, фолпет тәрізді қосылыстар, тетрагидрофтоламид бар екендігі айқындалды. Жақсы қылыш көлінің сынамалары галит, магний және сульфат тұзының түрлі қабаттарынан оқшауланған. Бұға-жайлы көліндегі тұзды шикізат құрамында $96,0 \pm 8,9$ масс% NaCl, магний иондарының мөлшері $0,17 \pm 0,01$ – $4,01 \pm 0,0\%$ құрайды, күкірттің мөлшері $0,26 \pm 0,02\%$ аралығында кездеседі. $1,71 \pm 0,12$ мас.%. Ауыр металл мөлшері қажетті концентрациядан төмен. Сынамалардағы магний құрамының жоғарылауы натрий-магний аралас қабаттарының бар екендігін дәлелдейтіні анықталды. Бұл тұзды көлдегі гетеротрофты микроорганизмдер саны шамамен 103-104 КҚБ/г құрайды, энтеробактериялар 103 КФБ/г, $(6,0 \pm 0,5) \times 10^3$ CFU/гт мөлшеріндегі микромицеттер тек сульфат үлгісінде анықталған тұз бен лай жағалау сызығынан 1,0 м қашықтықта сыналды. Жағалау сызығындағы микрофлораның таралуы мен тығыздығы тұз шығару эсерінен антропогендік фактордың жоғарылағанын көрсетеді.

Түйін сөздер: тұзды көл, құрамында тұзы бар шикізат, микрофлора, гетеротрофты микроорганизмдер, микромицеттер, пестицидтер.

А.У. Исаева^{1*}, Б. Леска², А.А. Успабаева³, А.Е. Тлеукеева³, А. Абубакирова³

¹ Шымкентский университет, Шымкент, Казахстан;

² Познаньский государственный университет им. А. Мицкевича, Познань, Польша;

³ Южно-Казахстанский государственный университет им. М. Ауэзова, Шымкент, Казахстан

БИОРАЗНООБРАЗИЕ МИКРОФЛОРЫ СОЛЕННЫХ ОЗЕР ДЖАКСЫ-КЛЫЧ И БУГАЖАЙЛЫ

Аннотация. В статье представлены результаты микробиологических и химических исследований образцов соленых озер юга Казахстана, таких как Джаксы-Клыч и Буга-Джайлы.

Континентальные соленые водоемы существенно различаются по химическому и микробиологическому составу, а моря похожи по составу. Состав этих вод напрямую зависит от геологических, климатических и биогеохимических условий и может быть нейтральным, щелочным и кислым. Популяция микроорганизмов в соленых водоемах очень мала, и их распространение зависит от концентрации соли.

Адаптивные механизмы основаны на регуляции концентрации ионов во внутриклеточной жидкости и ее осмотического давления. Концентрация неорганических ионов в водной среде может широко варьироваться, что приводит к увеличению ионного градиента между организмом и окружающей средой. Для применения солесодержащего сырья из двух соленых озер на юге Казахстана в качестве основы для косметических средств, необходимо было убедиться, что оно безвредно для здоровья человека. В связи с этим были проведены физико-химические и микробиологические исследования.

Установлено, что в пробах со дна высохшего Малого Аральского моря преобладают ионы магния, калия, кальция, железа и меди. Кроме того, выявлено наличие пестицидов: ацетамиприда, фенгексамида, фолпетоподобных соединений, тетрагидрофтоламида, которые используются в агропромышленном комплексе. Пробы озера Джаксы-Клыч были выделены из разных слоев галита, магния и сульфатных солей. Солесодержащее сырье озера Буга-Джайлы содержит $96,0 \pm 8,9$ мас.% NaCl, содержание ионов магния колеблется от $0,17 \pm 0,01$ до $4,01 \pm 0,0$ мас.%,

Содержание серы находится в пределах $0,26 \pm 0,02\%$. $1,71 \pm 0,12$ мас.%. Количество тяжелых металлов ниже допустимой концентрации. Установлено, что повышенное содержание магния в образцах свидетельствует о наличии смешанных натрий-магниевых слоев. Количество гетеротрофных микроорганизмов в образцах соленых озер составляет примерно 103-104 КОЕ/г, энтеробактерий -103 КОЕ/г, микромицеты в количестве $(6,0 \pm 0,5) \times 10^3$ КОЕ/г обнаружены только в образце сульфатных солей и илов, отобранных на расстоянии 1,0 м от береговой линии. Распределение и плотность микрофлоры на береговой линии указывают на повышенный антропогенный фактор из-за добычи соли.

Ключевые слова: соленые озера, солесодержащее сырье, микрофлора, гетеротрофные микроорганизмы, микромицеты, пестициды.

Information about authors:

Issayeva Akmaral Umurbekovna, ScD, Professor, Director of Ecology and Biology research Institute, Shymkent University, Shymkent;

Leska Boguslawa, Dr hab, Professor, Department of Chemistry, A. Mickiewicz Poznań State University, Poznań, Poland;

Uspabaeva Aigul Amankulovna, PhD, head of the biotechnology laboratory, M. Auezov South Kazakhstan University, Shymkent;

Tleukeeva Assel Yerzhanovna, doctoral student, M. Auezov South Kazakhstan University, Shymkent;

Abubakirova Azhar, doctoral student, M. Auezov South Kazakhstan University, Shymkent

REFERENCES

- [1] Aladin N., Micklin P. & Plotnikov I. (2008) Biodiversity of the Aral Sea and its importance to the possible ways of rehabilitating and conserving its remnant water bodies. In: NATO Science for Peace and Security Series – C: Environmental Security. Environmental Problems of Central Asia and their Economic, Social and Security Impacts. Edited by Jianguo Qi, Kyle T. Evered. Springer:73–98.
- [2] Berger E., Frör O. & Schäfer R.B. (2019) Salinity impacts on river ecosystem processes: a critical mini-review. *Phil. Trans. R. Soc. B* 374, 2018. 0010. (doi:10.1098/rstb.2018.0010)
- [3] Biswas J& Paul AK.(2017) Diversity and production of extracellular polysaccharide by halophilic microorganisms. *Biodiversity Int J*. 1(2):32–39. DOI: 10.15406/bij.2017.01.00006
- [4] DasSarma S& DasSarma P. (2015) Halophiles and their enzymes: negativity put to good use. *Curr Opin Microbiol*. 25:120-6. doi: 10.1016/j.mib.2015.05.009. Epub 2015 Jun 9.
- [5] Derevenskaya O. & N Urazaeva. (2018) Evaluation of the lake Lybiazhie (Kazan, Russia) state by indicators of communities of hydrobionts. *IOP Conf. Ser.: Earth Environ. Sci.* 107 012129
- [6] Dobrynin E.G. (1984) Microbiological processes of organic matter circulation in hyperhaline reservoirs [*Microbiologicheskie process krugovorota organicheskogo veshchestva v gypergalinnih vodoemah*], PhD thesis-Borok, 1984. 237 p. (in Russ.)
- [7] Ejsmont-Karabin J & Karabin A (2013) The suitability of zooplankton as lake ecosystem indicators: crustacean trophic state index *Polish Journal of Ecology* 61(3) pp 561-573
- [8] Horrigan N. (2005) Response of stream macroinvertebrates to changes in salinity and the development of a salinity index. / N. Horrigan [et al.] // *Mar. Freshwat. Res.* 56. : 825–833.
- [9] Haberman J& Haldna M (2014) Indices of zooplankton community as valuable tools in assessing the trophic state and water quality of eutrophic lakes: long term study of Lake Vörtsjärv *J.Limnol.* 73 (2) pp 263-273, DOI: 10.4081/jlimnol.2014.828J.
- [10] GOST 33770-2016. Food salt. Sampling and sample preparation. Determination of organoleptic parameters.
- [11] Loukas A., Kappas I. & Abatzopoulos T.J. (2018) HaloDom: a new database of halophiles across all life domains. *J Biol Res (Thessalon)*. 15;25:2. doi: 10.1186/s40709-017-0072-0. eCollection 2018 Dec.
- [12] Paul S., Bag S.K., Das S., Harvill ET&Dutta C. (2008) Molecular signature of hypersaline adaptation: insights from genome and proteome composition of halophilic prokaryotes. *Genome Biol.* 9;9(4):R70. doi: 10.1186/gb-2008-9-4-r70.
- [13] Rivera-Ingraham G.A. & Lignot J-H. (2017). Osmoregulation, bioenergetics and oxidative stress in coastal marine invertebrates: raising the questions for future research. *J. Exp. Biol.* 220, 1749-1760. (doi:10.1242/jeb.135624)
- [14] Velasco J., Gutiérrez-Cánovas C., Botella-Cruz M., Sánchez-Fernández D., Arribas P., Carbonell J. A., Millán A. & Pallarés S. (2019). Effects of salinity changes on aquatic organisms in a multiple stressor context. *Phil. Trans. R. Soc. B*. 374, 2018.0011, <http://doi.org/10.1098/rstb.2018.0011>
- [15] Venâncio C., Castro B.B., Ribeiro R., Antunes S.C., Abrantes N., Soares AMVM & Lopes I. (2019). Sensitivity of freshwater species under single and multigenerational exposure to seawater intrusion. *Phil. Trans. R. Soc. B* 374, 20180252. (doi:10.1098/rstb.2018.0252)
- [16] Wang Y., Mopper S & Hasenstein K.H. (2001) Effects of salinity on endogenous levels of ABA, IAA, JA, and SA in *Iris hexagona*. *Journal of Chemical Ecology*. 27: 327 – 342.
- [17] Adilbektegi G.A., Mustafayev J.S., Uvatayeva T.K., Dulatbekova Z.N., Mosiej Jozef. (2019) Quantitative and qualitative assessment of biological and ecological potential of the landscapes of Southern Kazakhstan, News of the academy of sciences of the Republic of Kazakhstan, <https://doi.org/10.32014/2019.2518-170X.160>

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

<http://reports-science.kz/index.php/en/archive>

Редакторы: *М. С. Ахметова, Д. С. Аленов, А. Ахметова*

Верстка на компьютере *А. М. Кульгинбаевой*

Подписано в печать 04.12.2020.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
8,25 п.л. Тираж 500. Заказ 6.