

ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

2020 • 3

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫң

БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944



ALMATY, NAS RK

Бас редакторы
х.ғ.д., проф., ҚР ҰҒА академигі
М.Ж. Жұрынов

Редакция алқасы:

Адекенов С.М. проф., академик (Қазақстан) (бас ред. орынбасары)
Величкин В.И. проф., корр.-мүшесі (Ресей)
Вольдемар Вуйчик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Белорус)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Тәжікстан)
Кригер Виктор проф. (Германия)
Леска Богуслава проф. (Польша)
Локшин В.Н. проф., чл.-корр. (Қазақстан)
Нараев В.Н. проф. (Ресей)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Уздир проф. (Малайзия)
Перни Стефано проф. (Ұлыбритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Ұлыбритания)
Омбаев А.М. проф., корр.-мүшесі (Қазақстан)
Өтелбаев М.О. проф., академик (Қазақстан)
Садыбеков М.А. проф., корр.-мүшесі (Қазақстан)
Сатаев М.И. проф., корр.-мүшесі (Қазақстан)
Северский И.В. проф., академик (Қазақстан)
Сикорски Марек проф., (Польша)
Рамазанов Т.С. проф., академик (Қазақстан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Харин С.Н. проф., академик (Қазақстан)
Чечин Л.М. проф., корр.-мүшесі (Қазақстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Қытай)
Эркебаев А.Ә. проф., академик (Қыргыстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының баяндамалары»
ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.).
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 01.06.2006 ж.
берілген №5540-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылдан 6 рет.

Тиражы: 500 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19, 272-13-18,
<http://reports-science.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2020

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Г л а в н ы й р е д а к т о р
д.х.н., проф., академик НАН РК
М. Ж. Журинов

Р е д а к ц и о н на я кол л е г и я:

Адекенов С.М. проф., академик (Казахстан) (зам. гл. ред.)
Величкин В.И. проф., чл.-корр. (Россия)
Вольдемар Вуйчик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Беларусь)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Таджикистан)
Кригер Виктор проф. (Германия)
Леска Богуслава проф. (Польша)
Локшин В.Н. проф., чл.-корр. (Казахстан)
Нараев В.Н. проф. (Россия)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Перни Стефано проф. (Великобритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Великобритания)
Омбаев А.М. проф., чл.-корр. (Казахстан)
Отелбаев М.О. проф., академик (Казахстан)
Садыбеков М.А. проф., чл.-корр. (Казахстан)
Сатаев М.И. проф., чл.-корр. (Казахстан)
Северский И.В. проф., академик (Казахстан)
Сикорски Марек проф., (Польша)
Рамазанов Т.С. проф., академик (Казахстан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Харин С.Н. проф., академик (Казахстан)
Чечин Л.М. проф., чл.-корр. (Казахстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Китай)
Эркебаев А.Э. проф., академик (Кыргызстан)

Доклады Национальной академии наук Республики Казахстан»

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5540-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 500 экземпляров

Адрес редакции: 050010, г.Алматы, ул.Шевченко, 28; ком. 219, 220; тел. 272-13-19, 272-13-18,
<http://reports-science.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2020 г.

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

M.Zh. Zhurinov**E d i t o r i a l b o a r d:****Adekenov S.M.** prof., academician (Kazakhstan) (deputy editor in chief)**Velichkin V.I.** prof., corr. member (Russia)**Voitsik Valdemar** prof. (Poland)**Goncharuk V.V.** prof., academician (Ukraine)**Gordiyenko A.I.** prof., academician (Belarus)**Duka G.** prof., academician (Moldova)**Ilolov M.I.** prof., academician (Tadzhikistan)**Krieger Viktor** prof. (Germany)**Leska Boguslava** prof. (Poland)**Lokshin V.N.** prof., corr. member (Kazakhstan)**Narayev V.N.** prof. (Russia)**Nekludov I.M.** prof., academician (Ukraine)**Nur Izura Udzir** prof. (Malaysia)**Perni Stephano** prof. (Great Britain)**Potapov V.A.** prof. (Ukraine)**Prokopovich Polina** prof. (Great Britain)**Ombayev A.M.** prof., corr. member (Kazakhstan)**Otelbayev M.O.** prof., academician (Kazakhstan)**Sadybekov M.A.** prof., corr. member (Kazakhstan)**Satayev M.I.** prof., corr. member (Kazakhstan)**Severskyi I.V.** prof., academician (Kazakhstan)**Sikorski Marek** prof., (Poland)**Ramazanov T.S.** prof., academician (Kazakhstan)**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief**Kharin S.N.** prof., academician (Kazakhstan)**Chechin L.M.** prof., corr. member (Kazakhstan)**Kharun Parlar** prof. (Germany)**Endzhun Gao** prof. (China)**Erkebayev A.Ye.** prof., academician (Kyrgyzstan)**Reports of the National Academy of Sciences of the Republic of Kazakhstan.**

ISSN 2224-5227

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5540-ЖК, issued 01.06.2006.

Periodicity: 6 times a year.

Circulation: 500 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://reports-science.kz/index.php/en/archive>

Agriculture

REPORTS OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

ISSN 2224-5227

Volume 3, Number 331 (2020), 20 – 28

<https://doi.org/10.32014/2020.2518-1483.51>

UDC 636.295/296

D. A. Baimukanov

Educational Scientific and Production Center Bayserke-Agro LLP,
Talgar district, Almaty region, Kazakhstan.
E-mail: dbaimukanov@mail.ru

REGULARITIES OF DEVELOPMENT OF COLTS OF THE KAZAKH BACTRIAN BREED

Abstract. Based on the studies, it was found that for parental couples with a coefficient of wool clip up to 0.8 at the age of one year, the amount of wool clip was 2.4 kg, which is significantly lower by 12.5% compared to herdmates from parents with a coefficient of wool clip of 0.9-1.4, and 29.2% lower in comparison with young one-year-olds obtained from parental couples with the coefficient of wool clip of 1.5 and above.

It was established that the real wool cut was lower in two-year-old females obtained from parents with a coefficient of wool clip of up to 0.8 and 1.5 and higher. In two-year-old females obtained from parental couples with the coefficient of the wool clip of 0.9-1.4, the real wool cut was 9.1% higher than predicted. In the future, during the selection, it is recommended to give preference to individuals with a coefficient of shear of wool 0.9-1.4.

In colts of the Kazakh Bactrian of the South Kazakhstan type obtained from female camels corresponding to the technological parameters of dairy productivity, exceeded their mates of the wool-meat direction of productivity in live weight by 17.8% ($P < 0.01$), of the milk and meat direction - in ($P < 0.1$)

After weaning, the intensity of growth and development is significantly reduced, which affected the increase in absolute live weight from 6 months to 9 months of age. After weaning, the colts from female camels of dairy productivity, the live weight increased by 11.9%, of the wool and meat productivity - by 34.0%, of the milk and meat direction of productivity - by 23.3%. That is, early weaning is not beneficial for colts from camels of the dairy direction of productivity. From one year to 15 months of age, young livestock from dairy camels have increased live weight by 17.1%, from wool and meat camels - by 19.1%, from milk and meat camels - by 16.6%.

From the age of 9 months to 15 months, the colts increased their live weight obtained from mothers with the coefficient of wool clip up to 0.8 by 58.7%, with the coefficient of 0.9-1.4 - by 55.3%, with the coefficient of 1.5 and above - by 91.1 %. The height between the humps increases by 15.7% -11.4% -11.7% respectively. That is, all female camels are of wool-meat productivity directions. The effectiveness of conducting assessment and selection by the coefficient of wool clip has been proved in the conditions of Bayserke-Agro LLP.

Key words: Kazakh Bactrian, assessment, selection, technological parameters, colts, early ripeness.

The relevance of the topic. Camel breeding in the Republic of Kazakhstan is developing dynamically. There is an interest in agricultural enterprises of various forms of property to develop a productive direction of camel breeding.

A deterrent to the development of camel breeding in the Almaty region is the small number of females and producers. The rotation system of sampler-producers is violated, as a result of this the frequency of using moderate and even related mating of camels of reproductive age increases.

Many elaborations of local scholars regarding the use of approved methods for selection and breeding using commercial livestock are not effective.

One of the urgent problems in camel breeding is the long fructification period of the Kazakh Bactrian camels and low fertility up to 42%. Despite the available breeding achievements in Kazakh Bactrians, the effective methods for assessing dairy, meat and wool productivity have not yet been developed [1].

Reasonable methods of reproduction of the breeding stock of camels have not been worked out, taking into account the technological parameters of productivity.

In connection with the above, the choice of the research direction to improve breeding efficiencies and increase productivity through targeted selection and breeding of purebred camels of the Kazakh Bactrian in the conditions of the Bayserke-Agro Educational Scientific and Production Center is correct.

Therefore, the development of effective criteria for evaluating camels by productivity based on a study of the regularities of growth and development of camels is relevant.

The aim of the research. To develop processing technologies for increasing the productive capacity of camels and improving the breeding efficiencies of the purebred animals.

Scientific novelty. For the first time, the influence of technological parameters of productivity of the Kazakh Bactrian camels on their productivity and early ripeness was established in the post-dairy period. For the first time, the regularities of postembryonic growth and development of young camels in the Almaty region were found.

The practical value of the research. Criteria for reliable assessment and selection of camels of the Kazakh Bactrian breed by the productivity direction have been developed. Using various options for the selection of parental pairs made it possible to obtain offspring with the given productive and technological parameters. Scientific provisions, conclusions, and recommendations have been widely used in the selection of camels of the Kazakh Bactrian breed of the South Kazakhstan type under the conditions of Bayserke-Agro LLP.

Research methods. The object of the study was purebred Kazakh Bactrians of the South-Kazakhstan type.

Body measurements were studied according to the Instructions on the bonitation of camels 2014 [2]. The height between the humps was measured with an accuracy of 1.0 cm. The oblique body length and chest girth were measured with an accuracy of 1.0 cm, and the metacarpus girth was measured with up to 0.5 cm.

The live weight of camels was determined by individual weighing of animals on a stationary scale, as well as by calculation method at the request of Patent No. 15886 [3].

Morphofunctional features of the udder of camels were studied by the method of A. Baimukanov of 1972 [4]. The fat content in milk was determined using Milkotester device (2017). Total protein in milk was defined using AM-2 milk analyzer (2017).

The regularities of growth and development of the Kazakh Bactrian of the South Kazakhstan type born in 2019 was studied according to the common method [5]. We studied live weight and height at the withers in the postembryonic period.

The technological parameters for the selection of camels by dairy productivity were determined by the degree of full value of lactation and its impact on milk yield and fat content. Gradation of camels by the degree of full value of lactation was carried out in three ranks: up to 65 - 74; 75-84; 85 and more. The degree of full value of lactation in camels was determined according to the recommendation of professor A. Baimukanov by the formula [6].

$$СПЛ = УФ \times 100 / УСхн,$$

where СПЛ – degree of full value of lactation; УФ – real milk yield for the entire lactation period; УС – average daily milk yield in the third month of lactation; n – the number of days of lactation.

The milking capacity coefficient was determined by the requirement of the Patent of the Republic of Kazakhstan No. 22213 through the ratio of the real milk yield for the active period of lactation to live weight [7].

$$KM = YM / \bar{ЖM}$$

where KM – the milking capacity coefficient; YM – the milk yield per lactation; ЖM – live weight.

Gradation by milking capacity coefficient was carried out in three ranks: up to 1.4; 1.5-1.9; 2.0 and more.

The influence of the fertility index on the real milk yield in experimental camels of the Kazakh Bactrian of the South Kazakhstan type was carried out according to the common method. The fertility index was determined by the formula proposed by professor A. Baimukanov [8].

$$\Pi = 365 (n-1) \times 100/N,$$

where Π – the fertility index; n – number of coltings; N – the number of days between the first and last coltings.

The gradation by the fertility index was conducted in three ranks: up to 42; 42 - 47; 47 and more.

The influence of the gradation of the coefficient of wool clip (CWC) on the intensity of achieving the best finish and live weight was carried out in young males born in 2017 while the coefficient of wool clip is calculated by the formula [9].

$$KHIII = \frac{HIII}{\mathbb{X}M} \times 100$$

where $KHIII$ – coefficient of wool clip; $HIII$ – the amount of wool clip; $\mathbb{X}M$ – live weight.

The gradation by the coefficient of wool clip was carried out in three ranks: up to 0.8; 0.9-1.4; 1.5 and above.

Biometric processing of digital materials was carried out according to the common method [10].

Research results. Based on the studies, it was found that for parental couples with a coefficient of wool clip up to 0.8 at the age of one year, the amount of wool clip was 2.4 kg, which is significantly lower by 12.5% compared to herdmates from parents with a coefficient of wool clip of 0.9-1.4, and 29.2% lower in comparison with young one-year-olds obtained from parental couples with a coefficient of wool clip of 1.5 and above (table 1).

Table 1 – Influence of the coefficient of wool clip on the indices of practical shearing of wool in young camels of the Kazakh Bactrian

Traits	Parental coefficient of wool clip		
	up to 0.8	0.9 -1.4	1.5 and more
The number of estimated daughters, animals	5	5	6
Live weight at birth, kg	29.3±1.7	37.7±1.1	35.5±1.6
Live weight at weaning, kg	245.8±9.8	269.2±14.5	239.1±19.1
Wool clip at one-year-old age, kg	2.4±0.3	2.7±0.3	3.1±0.2
Predicted wool clip at the age of two years, kg	2.9	3.3	4.2
Real wool clip at the age of two years, kg	2.8±0.2	3.6±0.4	3.7±0.4
The yield of pure fiber at the age of one year, %	95.2	95.5	95.5
The yield of pure fiber at the age of two years, %	96.5	96.6	96.5

It was established that the real wool clip was lower in two-year-old females obtained from parents with the coefficient of wool clip of up to 0.8 and 1.5 and more. In two-year-old females, the real wool clip obtained from parental couples with the coefficient of wool clip of 0.9-1.4 was 9.1% higher than the predicted.

With that in mind, we believe that the coefficient of wool clip is a reliable criterion for assessing camels of the Kazakh Bactrian breed in terms of shearing. In the future, during the selection, it is advisable to give preference to individuals with the coefficient of wool clip of 0.9-1.4.

Concerning the study of the yield of pure fiber, it was established that the produced wool is characterized by high technological qualities due to targeted selection with camels of the Kazakh Bactrian breed. At the same time, the pure fiber yield of 95.2-96.6% indicates the presence of inbreeding in the selection and breeding of parental couples. Therefore, to avoid mortality from the offspring in subsequent years, it is necessary to purchase sampler-producers from other regions of Kazakhstan.

In colts, as in calves, the early ripeness depends on the dairy productivity of their mothers. The regularities of milk formation as well as the formation of dairy productivity in cattle [11] are to some extent identical to camels.

In Kazakh Bactrian colts of the South Kazakhstan type obtained from camels of the dairy productivity exceeded their mates of the wool-meat direction of productivity in live weight by 17.8% ($P<0.01$), milk and meat in ($P<0.1$) (table 2).

Table 2 – Growth and development of colts in the post-dairy period, depending on the technological parameters of the breeding traits of their mothers (n=5)

Technological parameters of productivity	Age	Live weight, kg	Height between humps, cm
Dairy	At birth	40.3±1.9	109.7±2.1
	3 months	114.0±4.4	130.3±3.8
	6 months	169.9±6.3	145.1±3.3
	9 months	190.2±4.2	148.9±1.5
	12 months	232.7±5.7	155.4±1.1
	15 months	272.5±6.4	159.2±1.3
Wool - meat	At birth	34.2±1.5	104.0±2.4
	3 months	94.6±3.7	123.4±3.1
	6 months	144.5±5.1	137.2±3.5
	9 months	193.6±3.5	145.6±2.2
	12 months	240.1±4.7	151.1±2.7
	15 months	289.6±5.3	156.3±1.9
Milk-meat	At birth	36.8±1.9	107.1±1.8
	3 months	108.9±2.8	128.6±2.1
	6 months	156.2±6.1	141.9±2.5
	9 months	192.6±4.9	146.3±2.9
	12 months	242.4±5.2	152.8±2.7
	15 months	282.7±3.6	158.2±2.3

After weaning, the intensity of growth and development was significantly reduced, which affected the increase in absolute live weight from 6 months to 9 months of age. After weaning, the colts from female camels of dairy productivity, the live weight increased by 11.9%, of the wool and meat productivity - by 34.0%, of the milk and meat direction of productivity - by 23.3%. That is, early weaning is not beneficial for colts from camels of the dairy direction of productivity.

In young Kazakh Bactrians obtained from females of dairy productivity, the increase in live weight for the next 90 days (from 9 months to 12 months) was 42.5 kg, from the group of camels of wool-meat productivity direction - 46.5 kg and obtained from milk-meat female camels - 49.8 kg.

From one year to 15 months of age, young livestock from dairy camels have increased live weight by 17.1%, from wool and meat camels - by 19.1%, from milk and meat camels - by 16.6%.

From 2018 to 2020, the influence of the technological parameters of female camels on the growth and development of colts after weaning was studied. Weaning was performed at the age of 9 months (table 3).

When studying female camels of the main herd with technological parameters according to the degree of full value of lactation (DFL), it was found that with a degree of full value lactation of 75-84%, the height at the withers (8.8 cm) is intensively increased in colts from 9 months of age to one year old, in comparison with mates received from females with DFL of 65-74% (5.7 cm) and 85% and above (8.5 cm). Likewise, the live weight of colts from females is increasing intensively: with DFL of 75-84% (37.6 kg), with DFL of 65-74% (39.4 kg), and with DFL of 85% and above (41.1 kg) (table 3).

In the next three months, the colts have an increase in live weight: in those from female camels with DFL of 65-74% - 37.4 kg or 16.2%, with DFL of 75-84% - 16.2 kg or 6.8%, with DFL of 85% and above - 24.2 kg or 10.3%.

It has been found that in female camels with the milking capacity (MC) of up to 1.4; 5-1.9 and 2.0 and higher in the post-embryonic period, the live weight increases in camels from 9 months of age to one-year-old by 32.9% -19.9% -21.8%. From 12 months to 15 months of age by 14.1% -15.5% -12.45%. That is, in the first three months, the colts from female camels with MC of up to 1.4 grow intensively, in the next three months - the colts from camels with MC of 1.5-1.9.

Table 3 – The influence of the technological parameters of mothers on the growth and development of young stock in the post-dairy period

Traits	Age	Technological parameters		
		The degree of full value of lactation, %		
		65-74	75-84	85 and higher
Live weight, kg	9 months	192.1±3.7	202.3±7.4	193.6±5.8
	12 months	231.5±4.2	239.9±5.7	234.9±8.1
	15 months	268.9±9.4	256.1±6.1	259.1±7.5
Height between humps, cm	9 months	142.4±1.1	137.5±0.9	139.3±1.2
	12 months	148.1±0.9	146.3±1.2	147.8±1.4
	15 months	155.8±1.5	154.6±1.1	155.2±1.3
Milking capacity				
		up to 1.4	1.5-1.9	2.0 and higher
Live weight, kg	9 months	177.9±8.2	196.9±4.2	190.3±6.4
	12 months	236.4±5.7	236.2±6.4	231.9±8.2
	15 months	269.9±8.2	272.9±4.2	260.8±6.4
Height between humps, cm	9 months	139.2±1.3	145.1±1.1	141.4±1.1
	12 months	145.7±1.1	151.5±1.4	150.3±1.2
	15 months	152.3±1.2	158.2±1.3	154.5±1.6
Fertility index				
		up to 42	42-47	47 and higher
Live weight, kg	9 months	181.6±6.2	191.5±3.7	172.1±8.5
	12 months	237.1±8.7	246.5±5.4	235.6±6.8
	15 months	271.6±5.9	281.3±4.3	275.4±9.2
Height between humps, cm	9 months	140.9±1.7	141.4±1.5	141.1±1.6
	12 months	143.4±1.5	148.1±1.8	146.8±1.4
	15 months	164.3±1.9	156.4±1.6	151.7±1.3
Coefficient of wool clip				
		up to 0.8	0.9 -1.4	1.5 and higher
Live weight, kg	9 months	168.9±7.1	187.6±4.9	190.1±8.3
	12 months	221.3±5.8	245.1±6.2	233.7±6.8
	15 months	268.1±6.3	291.5±5.8	281.2±7.1
Height between humps, cm	9 months	131.3±1.7	139.5±1.3	147.1±1.5
	12 months	141.1±1.4	146.1±1.8	152.8±1.1
	15 months	151.9±1.3	155.4±1.6	164.3±1.3

The colts in the post-dairy period obtained from females with the fertility index of up to 42% develop more intensively, compared with mates received from camels with the fertility index of 42-47%, 47 and higher. From the age of 9 months to 15 months, the height at the withers increases by 16.6% -10.6% - 7.5%. In terms of live weight, young camels from mothers with the fertility index of 47% and above 60.0% intensively grow, compared with the mates of 49.6% and 46.9%.

From the age of 9 months to 15 months, the colts increased their live weight obtained from mothers with the coefficient of wool clip up to 0.8 by 58.7%, with the coefficient of 0.9-1.4 - by 55.3%, with the coefficient of 1.5 and above - by 91.1 %. The height between the humps increases by 15.7% -11.4% - 11.7% respectively. That is, all female camels are of wool-meat productivity directions. The effectiveness of conducting assessment and selection by the coefficient of wool clip has been proved in the conditions of Bayserke-Agro LLP.

Conclusions. To increase the early ripeness of the Kazakh Bactrian colts, it is needed to increase the selection of the breeding stock with the degree of full value of lactation of 75-84%, with the dairy productivity rank of 1.5-1.9, with the fertility index of 42-47% and the coefficient of wool clip of 0.9-1.4.

The studies have been carried out under the program of the Ministry of Agriculture of the Republic of Kazakhstan for 2018 - 2020. IRN: BR06249249 Development of the integrated system for increasing productivity and improving breeding qualities of farm animals by the example of Bayserke-Agro LLP.

Д. А. Баймұқанов

«Байсерке-Агро оку-ғылыми өндірістік орталығы» ЖШС, Алматы обласы, Қазақстан

ҚАЗАҚ ҚОС ӨРКЕШТІ ТҮЙЕЛЕРИНІҢ БОТАЛАРЫНЫң ДАМУ ЗАҢДЫЛЫҚТАРЫ

Аннотация. Қазақ бактериан түйесінің қырқылған шудасын бағалау нәтижелері бойынша сұрыптау критерилерін жасау, зерттеудің өзекті бағыты болып табылады, себебі келер ұрпақтарды алынатын шуда шығымын болжауға мүмкіндік береді. Жүргізілген зерттеулер негізінде анықталғандай, шуда шығымының коэффициенті 0,8 болған ата – аналық жұптардардың ұрпақтарында 1 жасындағы боталарында шуда шығымы 2,4 кг, жұн шығымы 0,9 – 1,4 коэффициенте ата – аналық жұптарда 29,2% төмен және жұн шығымы 1,5 және одан жоғары коэффициентте 1 жас боталардан төмен.

Шуда шығымы 0,8 және 1,5 және жоғары коэффициенті ата – аналық жұптардан алынған 2 жастағы інгендерде жұн шығымы қүтілгеннен төмен болды. Жұн шығымы 0,9 – 1,4 коэффициентке ие ата – аналық жұптардан алынған інгендерден шуда шығымы болжалғаннан 9,1 % жоғары болды. Осыған сүйене отырып, қазақ бактериан түйелерінің шудаларының шығымы анықтаушы критерий болып табылады. Ары қарай, сұрыптауда жұн шығымы 0,9 – 1,4 коэффициентті түрлер алынды.

Қазақ бактериан түйелерінің бағытталған селекциясы барысында алынған шудалар жоғары технологиялық сапаларға ие. Алынған таза талшықтың шығым көрсеткіші 95,2-96,6%, бұл ата – аналық жұптарда тандауда және сұрыптауда инбридингтің бар екендігін дәлелдейді. Сондыктan ұрпақтың төмедеуі болмауы үшін Қазақстанның басқа аймақтарынан буралар сатып алу керек.

Сәйкес технологиялық параметрлермен алынған қазақ бактериан боталарының онтүстік қазақстан типінде сүттілігі жүнді – етті бағыттағы өнімділігі бойынша тірі салмағы күрдастарынан басым 17,8% ($P<0,01$), сүтті - етті ($P<0,1$).

Қарқынды өсу мен даму тоқтағаннан кейін 6 айлық 9 айлық аралықта тірі салмақта абсолютті ұлғаюы айтартылтай төмендеді. Сүтті бағыттағы боталарды інгендерден бөлгеннен кейін тірі салмақ 11,9%, жүнді – етті 34,0%, сүтті - етті - 23,3%. Боталарды інгендерден ерте бөлу тиімсіз.

Сүтті бағыттағы қазақ қос өркешті інгендерінен алынған боталар кейінгі 90 күнде (9 айдан 12 айға дейін) тірі салмақтың өсімі 42,5 кг құрады, жүнді – етті бағыттағылардан – 46,5 кг және сүтті – етті бағыттағы інгендерден – 49,8 кг.

Сүтті бағыттағы інгендерден алынған төлдердің 1 жастаң және 15 айлық жаста тірі салмағы 17,1%, жүнді – еттіде - 19,1%, сүтті – еттіде – 16,6% ұлғайды.

2018 ж. және 2020 жыл аралығында төлдерді інгендерден бөлгеннен кейін боталардың өсуі мен дамуына технологиялық параметрлердің әсері зерттелді. Анасынан бөлу 9 айда іске асырылды.

Негізгі үйірдегі інгендердің технологиялық параметрлерінің толық лактациялық дәрежесі бойынша зерттеу 9 айлық төдерді толық лактация дұрыс болғанда 9 айдан 1 жасқа дейінгі аралықта шоқтық биіктігі өсу қарқындылығы (8,8 см), інгендердің лактация дәрежесі 65-74% бойынша аналогтарымен салыстырғанда (5,7 см) және 85% және (8,5 см) жоғары. ЛДТ інгендердің боталарында тірі салмақтың және өсу қарқындылығы 75-84% (37,6 кг), 65-74% (39,4 кг), 85% және жоғары (41,1 кг).

ЛДТ соңғы үш айда боталарда 65-74% - 37,4 кг немесе 16,2%, 75-84% - 16,2 кг немесе 6,8%, 85% және жоғары – 24,2 кг немесе 10,3%. 15 айлық жасқа жеткенде барлық боталарда шоқтық биіктігі 154,6-155,8 см бірдей болды. Алынған нәтижелер боталардың емізуден ажырағаннан кейін өсу мен даму қарқындылығына ЛТД ұтымдылығының әсері анықталмады.

Сүттілік коэффициенті 1,4 дейінгі, 9 – 1,9 және 2,0 және жоғары інгендерде туылғаннан кейінгі кезеңде боталардың тірі салмағы 9 айдан 12 ай аралығында 32,9%-19,9%-21,8%, 12 – 15 ай аралығында 14,1%-15,5%-12,45%. Алғашқы үш айда СК 1,4 інгендердің боталары қарқынды өседі, кейінгі үш айда СК 1,5-1,9 інгендерден алынған боталар қарқынды өсті.

Шоқтық биіктігінің ұлғауы бойынша бақылаудағы боталарда 9 айдан бір жасқа дейін 6,5 см-6,4 см-7,9 см, 12 айдан 15 айға дейін 6,6 см-6,7 см-4,2 см. 9 айдан 15 айға дейінгі аралықта шоқтық биіктігінің ұлғауы 13,3 см-13,1 см-13,1 см. Алынған нәтижелер қазак қос өркешті түйелерінде инбридинг қауіптілігін растайды.

Өнімділік индексі 42 % дейінгі інгендерден алынған боталар емізу мезгілінен кейін 42 – 47 % өнімділік индексі құрдастарымен салыстырғанда қарқынды дамиды. 9 айдан 15 ай аралығында шоқтық биіктігі 16,6%-10,6%-7,5% жоғарлайды. Тірі салмағы бойынша өнімділік индексі 47% және 60,0 % жоғары аналардан туылған боталар ие. Аналогтарымен салыстырғанда 49,6% және 46,9%.

Негізгі үйірде өнімділік индексі бойынша інгендерді сұрыптаудың тиімділігі анықталмады. Өндіруші буралар сатып алу керек.

9 айдан 15 ай аралығында шуда шығымы 0,8 коэффицентіне ие інгендерден алынған боталардың тірі салмағы 58,7%, 0,9-1,4 - 55,3%, 1,5 және жоғары 91,1%. Өркештер арасындағы биіктік 15,7%-11,4%-11,7% ұлғаяды. Корыта келгенде, барлық інгендер жүнді – етті бағытта. «Байсерке-Агро» ЖШС жағдайында шуданың қырқым коэффициенті бойынша бағалау мен сұрыптау жасау тиімді.

Түйін сөздер: қазак бактерианы, бағалау, сұрыптау, технологиялық параметрлер, боталар, жетілу.

Д. А. Баймukanов

ТОО “Учебный научно-производственный центр Байсерке-Агро”, Алматинская область, Казахстан

ЗАКОНОМЕРНОСТИ РАЗВИТИЯ ВЕРБЛЮЖАТ ПОРОДЫ КАЗАХСКИЙ БАКТРИАН

Аннотация. Разработка критерииев отбора по результатам оценки верблюдов породы казахский бактриан по коэффициенту настрига шерсти является актуальным направлением исследований, так как позволяет достоверно прогнозировать настриг шерсти у полученного потомства. На основании проведенных исследований установлено, что у родительских пар с коэффициентом настрига шерсти до 0,8 в верблюжата в годовалом возрасте имели настриг шерсти 2,4 кг, что достоверно ниже на 12,5% в сравнении со сверстницами, полученных от родителей с коэффициентом настрига шерсти 0,9-1,4, и на 29,2% ниже в сравнении с молодняком годовалого возраста, полученных от родительских пар с коэффициентом настрига шерсти 1,5 и выше.

Установлено, что фактический настриг шерсти оказался ниже у двухлетних самок, полученные от родителей с коэффициентом настрига шерсти до 0,8 и 1,5 и выше. У двухлетних самок, полученных от родительских пар с коэффициентом настрига шерсти 0,9-1,4, фактический настриг шерсти оказался на 9,1% выше прогнозируемого. Исходя из этого, считаем, что коэффициент настрига шерсти является достоверным критерием оценки верблюдов породы казахский бактриан по показателям настрига шерсти. В дальнейшем при отборе отдавать предпочтение особям с коэффициентом настрига шерсти 0,9-1,4.

Касательно изучения выхода чистого волокна установлено, что продуцируемая шерсть характеризуется высокими технологическими качествами, обусловленными целенаправленной селекцией с верблюдами породы казахский бактриан. В то же время, показатель выходы чистого волокна 95,2-96,6% свидетельствует о наличии инбридинга при отборе и подборе родительских пар. Поэтому во избежание падежа у приплода в последующие годы необходимо провести закуп бура-производителей из других регионов Казахстана.

У верблюжат казахского бактриана южно-казахстанского типа, полученных от верблюдоматок, соответствующие технологическим параметрам продуктивности – молочное превосходили сверстниц шерстно-мясного направления продуктивности по живой массе на 17,8% ($P<0,01$), молочно-мясного на ($P<0,1$).

После отъема интенсивность роста и развития значительно снижаются, что повлияло на увеличение абсолютной живой массы с 6 месячного возраста до 9 месячного. После отъема у верблюжат от верблюдоматок молочной продуктивности живая масса увеличилась на 11,9%, шерстно-мясной – 34,0%, молочно-мясного – 23,3%. То есть, ранний отъем не выгоден для верблюжат от верблюдоматок молочного направления продуктивности.

У молодняка казахского бактриана, полученных от верблюдоматок молочной продуктивности, за последние 90 дней (с 9 мес. до 12 мес.) увеличение живой массы составило 42,5 кг, от группы верблюдоматок шерстно-мясного направления продуктивности – 46,5 кг и верблюдоматок молочно-мясного направления – 49,8 кг.

От годовалого возраста до 15 месячного возраста у молодняка от верблюдоматок молочной продуктивности живая масса увеличилась на 17,1%, шерстно-мясного – на 19,1%, молочно-мясного – 16,6%.

С 2018 г по 2020 г. изучали влияние технологических параметров верблюдоваток на интенсивность роста и развития верблюжат после отъема. Отъем производили в 9 мес. возрасте.

При изучении верблюдоваток основного стада с технологическими параметрами по степени полноценности лактации (СПЛ) установлено, что при степени полноценности лактации 75-84% у верблюжат с 9 месячного возраста до годовалого возраста интенсивно растет высота в холке (8,8 см), в сравнении со сверстницами, полученными от верблюдоваток со СПЛ 65-74% (5,7 см) и 85% и выше (8,5 см). Аналогично интенсивно увеличивается и живая масса у верблюжат от верблюдоваток со СПЛ 75-84% (37,6 кг), 65-74% (39,4 кг), 85% и выше (41,1 кг).

В последующие три месяца у верблюжат увеличение живой массы составило от верблюдоваток со СПЛ 65-74% - 37,4 кг или 16,2%, 75-84% - 16,2 кг или 6,8%, 85% и выше – 24,2 кг или 10,3%. При достижении 15 месячного возраста высота в холке у всех верблюдоваток оказалась одинаковой 154,6-155,8 см. Полученные данные не позволили установить эффективность влияния СПЛ на интенсивность роста и развития молодняка в постмолочный период.

Установлено, что у верблюдоваток с коэффициентом молочности (КМ) до 1,4; 5-1,9 и 2,0 и выше в постэмбриональный период живая масса увеличивается у верблюжат с 9 месячного возраста до годовалого на 32,9%-19,9%-21,8%, с 12 месячного до 15 месячного возраста на 14,1%-15,5%-12,45%. То есть, в первые три месяца интенсивно растут верблюжата от верблюдоваток с КМ до 1,4, в последующие три месяца – верблюжата от верблюдоваток с КМ 1,5-1,9.

По высоте в холке увеличение составило у подопытных верблюжат от 9 месячного возраста до годовалого возраста 6,5 см-6,4 см-7,9 см, от 12 месячного возраста – до 15 месячного возраста 6,6 см-6,7 см-4,2 см. С 9 месячного возраста до 15 месячного возраста увеличение высоты в холке составило 13,3 см-13,1 см-13,1 см. Полученные данные подтверждают наше опасение о инбридинге при разведении верблюдов казахского бактриана.

Верблюжата в постмолочный период, полученные от верблюдоваток, с индексом плодовитости до 42% развиваются более интенсивно, в сравнении со сверстницами, полученными от верблюдоваток с индексом плодовитости 42-47%, 47 и выше. С 9 месячного возраста до 15 месячного возраста увеличивается высота в холке на 16,6%-10,6%-7,5%. По живой массе интенсивно растут верблюжата от матерей с индексом плодовитости 47% и выше 60,0%, в сравнении со сверстницами 49,6% и 46,9%.

То есть, не выявлена эффективность отбора по индексу плодовитости верблюдоваток основного стада. Подтверждается наше положение о необходимости закупа производителей.

Увеличили живую массу верблюжата с 9 месячного возраста до 15 месячного возраста, полученные от матерей с коэффициентом настрига шерсти до 0,8 на 58,7%, 0,9-1,4 на 55,3%, 1,5 и выше 91,1%. Высота между горбами увеличивается на 15,7%-11,4%-11,7%. То есть, все верблюдоватки являются шерстномясного направления продуктивности. Доказана эффективность ведения оценки и отбора по коэффициенту настрига шерсти в условиях ТОО «Байсерке-Агро».

Ключевые слова: казахский бактриан, оценка, отбор, технологические параметры, верблюжата, скороспелость.

Information about the author:

Baimukanov Dastanbek Asylbekovich, Corresponding Member of the National Academy of Sciences of the Republic of Kazakhstan, Doctor of Agricultural Sciences, Chief Researcher of the Baiserke-Agro ESPC, Talgar district, Almaty region, Kazakhstan; dbaimukanov@mail.ru; <https://orcid.org/0000-0002-4684-7114>

REFERENCES

- [1] Baimukanov D.A., Yuldashbayev Yu.A., Doshanov D.A. (2016) Camel breeding (Bachelor degree) [Verblyudovodstvo (Bakalavriat)]. Textbook (ISBN 978-5-906818-14-0). Moscow. KURS Publishing House, SIC INFRA 184 p. (in Russ.).
- [2] Instructions for bonitation of camels of Bactrian and Dromedary breeds with the basics of breeding [Instruktsiya po bonitirovke verblyudov porod baktrianov i dromedarov s osnovami plemennoy raboty] (2014). Astana 25 p. (in Russ.).
- [3] The method of professor Baimukanov A. and Baimukanov D.A. Determination of live weight in camels (2008). Patent of the Republic of Kazakhstan for invention No. 15886. [Sposob professora Baymukanova A. i Baymukanov D.A. opredeleniya zhivoy massy u verblyudov (2008). Patent RK na izobreteniye №15886.] Publ. 15.08.2008, bull. No. 8 (Baimukanov A., Baimukanov D.A.) (in Russ.).

- [4] Baimukanov A. (1972). Morphofunctional features of the udder of camels (03.00.13 –Physiology of humans and animals). Abstract diss. Biol.Science: [Morfofunktsional'nyye osobennosti vymeni u verblyudits (03.00.13 –Fiziologiya cheloveka i zhivotnykh). Avtoref.diss.biol.nauk] 30.05.1972. Alma-Ata: AZVI. 18 p. (in Russ.).
- [5] Baimukanov D.A. (2009). Selection of camels of the Kazakh Bactrian breed and methods for their improvement. [Selektsiya verblyudov porody kazakhskiy baktrian i metody ikh sovershenstvovaniya] Monograph (ISBN9965-413-90-8). Almaty, Bastau. 280 p. (in Russ.).
- [6] The method of selection of camels of the Kazakh Bactrian of dairy direction (2010) [Sposob selektsii verblyudov kazakhskogo baktriana molochnogo napravleniya]. Patent of the Republic of Kazakhstan No. 16226. Publ. 15.01.2010, Bulletin ... No. 1. (Baimukanov D.A., Baimukanov A., Imangaziev Z., Koshshan B.L., Zholdybayev T.) (in Russ.).
- [7] The method of selection of camels of the Kazakh Bactrian by milking capacity [Sposob otbora verblyudov kazakhskogo baktriana po molochnosti] (2010). Patent of the Republic of Kazakhstan No. 22213. Publ. 15.01.2010, bull. No. 1. (Baimukanov A., Turumbetov B.S., Baimukanov D.A., Alikhanov O.) (in Russ.).
- [8] The method of selection of camels of the Kazakh Bactrian [Sposob selektsii verblyudov kazakhskogo baktriana] (2010). Patent of the Republic of Kazakhstan No. 16747. Published 08.16.2010, bull. No. 8. (Baimukanov D.A., Baimukanov A., Imangaziev Z., Koshshan B.L., Zholdybayev T.) (in Russ.).
- [9] The method of selection of the Kazakh Bactrian by wool productivity [Sposob otbora kazakhskogo baktriana po sherstnoy produktivnosti] (2010) Patent of the Republic of Kazakhstan No. 22214. Publ. 15.01.2010, bull. No. 1. (Baimukanov A., Turumbetov B.S., Baimukanov D.A., Tastanov A.) (in Russ.).
- [10] Baimukanov D.A., Tarchokov T.T., Alentayev A.S., Yuldashbayev Yu.A., Doshanov D.A. (2016) Fundamentals of genetics and biometrics [Osnovy genetiki i biometrii]. Textbook (ISBN 978-601-310-078-4). Almaty Evero. 128 p.
- [11] Bekenov D.M., Spanov A.A., Chindaliyev A.E., Baimukanov A.D., Sultanbai D.T., Zhaksylykova G.K., Kalimoldinova A.S. (2019) Comparative study of fruitfulness of cow insemination of a milking herd at various levels of productivity in the conditions of Bayserke-Agro LLP // Bulletin of the National Academy of Sciences of the Republic of Kazakhstan. Vol. 6, N 382 (2019). P. 335–338. ISSN 2518-1467 (Online), ISSN 1991-3494 (Print). <https://doi.org/10.32014/2019.2518-1467.178>

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www:nauka-nanrk.kz

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

<http://reports-science.kz/index.php/en/archive>

Редакторы: М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов

Верстка на компьютере А.Д. Абдрахимовой

Подписано в печать 10.06.2020.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.
16,5 п.л. Тираж 500. Заказ 3.