

ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

**ACADEMIC JOURNAL
OF PHYSICAL AND CHEMICAL SCIENCES**

**№1
2026**

ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

2026 • 1



**ACADEMIC JOURNAL
OF PHYSICAL AND
CHEMICAL SCIENCES**

PUBLISHED SINCE JANUARY 1944

ALMATY, NAS RK

EDITOR-IN-CHIEF

ZHURINOV Murat Zhurinovich, Doctor of Chemical Sciences, Professor, Academician of IAAS and NAS RK, General Director Oil refining and Petrochemistry Research Institute (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

DEPUTY EDITOR-IN-CHIEF:

KALIMOLDAYEV Maksat Nuradilovich, Doctor of Physical and Mathematical Sciences, Professor, Academician of NAS RK (Almaty, Kazakhstan), <https://www.scopus.com/authid/detailuri?authorId=56153126500>, <https://www.webofscience.com/wos/author/record/2428551>

ABILMAGZHANOV Arlan Zainutallaevich, PhD in Chemistry, General Director of JSC "Institute of Fuel, Catalysis and Electrochemistry named after D.V. Sokolsky", (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57197468109>, <https://www.webofscience.com/wos/author/record/2024265>

EDITORIAL BOARD:

ADEKENOV Sergazy Mynzhasarovich, Doctor of Chemical Sciences, Professor, Academician of NAS RK, Director of the JSC "Phytochemistry Research and Production Center", (Karaganda, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=7006153118>, <https://www.webofscience.com/wos/author/record/48648658>

RAMAZANOV Tlekkabul Sabitovich, Doctor of Physical and Mathematical Sciences, Professor, Academician of NAS RK (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6701328029>, <https://www.webofscience.com/wos/author/record/13503476>

ABIEV Rufat, Doctor of Technical Sciences (Biochemistry), Professor, Head of the Department of Optimization of Chemical and Biotechnological Equipment, St. Petersburg State Technological Institute (St. Petersburg, Russia), <https://www.scopus.com/authid/detail.uri?authorId=6602431781>, <https://www.webofscience.com/wos/author/record/1405661>

OLIVIERO Rossi Cesare, PhD (Chemistry), Professor at the University of Calabria (Calabria, Italy), <https://www.scopus.com/authid/detail.uri?authorId=57221375979>, <https://www.webofscience.com/wos/author/record/399768>

TIGINYANU Ion Mihailovich, Doctor of Physical and Mathematical Sciences, Academician, President of the Academy of Sciences of Moldova, Technical University of Moldova (Chisinau, Moldova), <https://www.scopus.com/authid/detail.uri?authorId=7006315935>, <https://www.webofscience.com/wos/author/record/524462>

SANG SU Kwak, PhD (Biochemistry, Agricultural Chemistry), Professor, Chief Scientist, Research Center for Plant Systems Engineering, Korea Research Institute of Bioscience and Biotechnology (KRIBB), (Daecheon, Korea), <https://www.scopus.com/authid/detail.uri?authorId=59286321700>, <https://www.webofscience.com/wos/author/record/30028581>

BERSIMBAYEV Rakhmetkazhi Iskenderovich, Doctor of Biological Sciences, Professor, Academician of NAS RK, L.N. Gumilyov Eurasian National University (Astana, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=7004012398>, <https://www.webofscience.com/wos/author/record/19854255>

CALANDRA Pietro, PhD (Physics), Professor, Institute for the Study of Nanostructured Materials (Rome, Italy), <https://www.scopus.com/authid/detail.uri?authorId=7004303066>, <https://www.webofscience.com/wos/author/record/616360>

BOSHKAEV Kuantai Avgazyevich, PhD, Associate Professor, Department of Theoretical and Nuclear Physics, Al-Farabi Kazakh National University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=54883880400>, <https://www.webofscience.com/wos/author/record/2080231>

BURKITBAEV Mukhambetkali, Doctor of Chemical Sciences, Professor, Academician of NAS RK, (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=8513885600>, <https://www.webofscience.com/wos/author/record/29017135>

QUEVEDO Hernando, Professor, National Autonomous University of Mexico (UNAM), Institute of Nuclear Sciences (Mexico City, Mexico), <https://www.scopus.com/authid/detail.uri?authorId=55989741100>, <https://www.webofscience.com/wos/author/record/30353742>

ZHUSUPOV Marat Abzhanovich, Doctor of Physical and Mathematical Sciences, Professor of the Department of Theoretical and Nuclear Physics, Al-Farabi Kazakh National University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6602166928>, <https://www.webofscience.com/wos/author/record/566>

KOVALEV Alexander Mikhailovich, Doctor of Physical and Mathematical Sciences, Academician of NAS of Ukraine, Institute of Applied Mathematics and Mechanics (Donetsk, Ukraine), <https://www.scopus.com/authid/detail.uri?authorId=7202799321>, <https://www.webofscience.com/wos/author/record/65533963>

TAKIBAEV Nurgali Zhabagaevich, Doctor of Physical and Mathematical Sciences, Professor, Academician of NAS RK, Al-Farabi Kazakh National University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=24077239000>, <https://www.webofscience.com/wos/author/record/1671760>

KHARIN Stanislav Nikolaevich, Doctor of Physical and Mathematical Sciences, Professor, Academician of NAS RK, Kazakh-British Technical University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6701353063>, <https://www.webofscience.com/wos/author/record/2023295>

ABISHEV Medeu Erzhanovich, Doctor of Physical and Mathematical Sciences, Professor, Corresponding Member of NAS RK, (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=26530759900>, <https://www.webofscience.com/wos/author/record/1556025>

ACADEMIC JOURNAL OF PHYSICAL AND CHEMICAL SCIENCES.

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

Owner: «Central Asian Academic Research Center» LLP (Almaty).

The certificate of registration of a periodical printed publication in the Committee of Information of the Ministry of Information and Social Development of the Republic of Kazakhstan № KZ93VPY00121157 issued 05.06.2025

Thematic scope: *physics and chemistry*.

Periodicity: 4 times a year.

<http://reports-science.kz/index.php/en/archive>

БАС РЕДАКТОР

ЖҰРЫНОВ Мұрат Жұрынулы, химия ғылымдарының докторы, профессор, ХҒАҚ және ҚР ҰҒА академигі, Мұнай өңдеу және мұнай-химиясы ғылыми-зерттеу институтының бас директоры (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

БАС РЕДАКТОРДЫҢ ОРЫНБАСАРЛАРЫ:

КАЛИМОЛДАЕВ Мақсат Нұрәділұлы, физика-математика ғылымдарының докторы, профессор, ҚР ҰҒА академигі (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=56153126500>, <https://www.webofscience.com/wos/author/rec-ord/2428551>

ӘБИЛМАҒЖАНОВ Арлан Зайнуталлайұлы, химия ғылымдарының кандидаты, Д.В. Сокольский атындағы «Жанармай, катализ және электрохимия институты» АҚ Бас директоры (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=57197468109>, <https://www.webofscience.com/wos/author/record/2024265>

РЕДАКЦИЯ АЛҚАСЫ:

ӘДЕКЕНОВ Серғазы Мынжасарұлы, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, «Фитохимия» ғылыми-өндірістік орталығы» АҚ директоры (Қарағанды, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=7006153118>, <https://www.webofscience.com/wos/author/record/48648658>

РАМАЗАНОВ Тілеккабыл Сәбитұлы, физика-математика ғылымдарының докторы, профессор, ҚР ҰҒА академигі (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=6701328029>, <https://www.webofscience.com/wos/author/record/13503476>

ӘБИЕВ Руфат, техника ғылымдарының докторы (биохимия), профессор, Санкт-Петербург мемлекеттік технологиялық институты «Химиялық және биотехнологиялық аппаратураны оңтайландыру» кафедрасының меңгерушісі (Санкт-Петербург, Ресей), <https://www.scopus.com/author/detail.uri?authorId=6602431781>, <https://www.webofscience.com/wos/author/record/1405661>

ОЛИВЬЕРО Росси Сесаре, PhD (химия), Калабрия университетінің профессоры (Калабрия, Италия), <https://www.scopus.com/author/detail.uri?authorId=57221375979>, <https://www.webofscience.com/wos/author/record/399768>

ТИГИНЯНУ Ион Михайлович, физика-математика ғылымдарының докторы, академик, Молдова Ғылым академиясының президенті, Молдова техникалық университеті (Кишинев, Молдова), <https://www.scopus.com/author/detail.uri?authorId=7006315935>, <https://www.webofscience.com/wos/author/record/524462>

САНГ-СУ Квак, PhD (биохимия, агрохимия), профессор, Корей Биоғылым және биотехнология ғылыми-зерттеу институты (KRIBB), өсімдіктердің инженерлік жүйелері ғылыми-зерттеу орталығының бас ғылыми қызметкері (Дэчон, Корея), <https://www.scopus.com/author/detail.uri?authorId=59286321700>, <https://www.webofscience.com/wos/author/record/30028581>

БЕРСІМБАЕВ Рахметқажы Есендірұлы, биология ғылымдарының докторы, профессор, ҚР ҰҒА академигі, Л.Н. Гумилев атындағы Еуразия ұлттық университеті (Астана, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=7004012398>, <https://www.webofscience.com/wos/author/record/19854255>

КАЛАНДРА Пьетро, PhD (физика), нанокүрылымды материалдарды зерттеу институтының профессоры (Рим, Италия), <https://www.scopus.com/author/detail.uri?authorId=7004303066>, <https://www.webofscience.com/wos/author/record/616360>

БОШКАЕВ Қуантай Авғазыұлы, PhD теориялық және ядролық физика кафедрасының доценті, әл-Фараби атындағы Қазақ ұлттық университеті (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=54883880400>, <https://www.webofscience.com/wos/author/record/2080231>

БҮРКІТБАЕВ Мұхамбетқали, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=8513885600>, <https://www.webofscience.com/wos/author/record/29017135>

QUEVEDO Hernando, профессор, Мексика ұлттық автономиялық университеті (UNAM), Ядролық ғылымдар институты (Мехико, Мексика), <https://www.scopus.com/author/detail.uri?authorId=55989741100>, <https://www.webofscience.com/wos/author/record/30353742>

ЖҮСПІНОВ Марат Абжанұлы, физика-математика ғылымдарының докторы, теориялық және ядролық физика кафедрасының профессоры, әл-Фараби атындағы Қазақ ұлттық университеті (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=6602166928>, <https://www.webofscience.com/wos/author/record/566>

КОВАЛЕВ Александр Михайлович, физика-математика ғылымдарының докторы, Украина ҰҒА академигі, Қолданбалы математика және механика институты (Донецк, Украина), <https://www.scopus.com/author/detail.uri?authorId=7202799321>, <https://www.webofscience.com/wos/author/record/65533963>

ТАКИБАЕВ Нұрғали Жабағаұлы, физика-математика ғылымдарының докторы, профессор, ҚР ҰҒА академигі, әл-Фараби атындағы Қазақ ұлттық университеті (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=24077239000>, <https://www.webofscience.com/wos/author/record/1671760>

ХАРИН Станислав Николаевич, физика-математика ғылымдарының докторы, профессор, ҚР ҰҒА академигі (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=6701353063>, <https://www.webofscience.com/wos/author/record/2023295>

ӘБИШЕВ Медеу Ержанұлы, физика-математика ғылымдарының докторы, профессор, ҚР ҰҒА академигі (Алматы, Қазақстан), <https://www.scopus.com/author/detail.uri?authorId=26530759900>, <https://www.webofscience.com/wos/author/record/1556025>

ACADEMIC JOURNAL OF PHYSICAL AND CHEMICAL SCIENCES

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

Меншіктеуші: «Орталық Азия академиялық ғылыми орталығы» ЖШС (Алматы қ.).

Ақпарат агенттігінің мерзімді баспасөз басылымын, ақпарат агенттігін және желілік басылымды қайта есепке қою туралы ҚР Мәдениет және Ақпарат министрлігі «Ақпарат комитеті» Республикалық мемлекеттік мекемесі **05.06.2025 ж.** берген № **KZ93VPY00121157** Куәлік.

Тақырыптық бағыты: *физика, химия.*

Мерзімділігі: жылына 4 рет.

<http://reports-science.kz/index.php/en/archive>

ГЛАВНЫЙ РЕДАКТОР

ЖУРИНОВ Мурат Журинович, доктор химических наук, профессор, академик МАН и НАН РК, Генеральный директор Научно-исследовательского института нефтепереработки и нефтехимии (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

ЗАМЕСТИТЕЛИ ГЛАВНОГО РЕДАКТОРА:

КАЛИМОЛДАЕВ Максат Нурадилович, доктор физико-математических наук, профессор, академик НАН РК (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=56153126500>, <https://www.webofscience.com/wos/author/record/2428551>

АБИЛЬМАГЖАНОВ Арлан Зайнуталлаевич, кандидат химических наук, Генеральный директор АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского», (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57197468109>, <https://www.webofscience.com/wos/author/record/2024265>

РЕДАКЦИОННАЯ КОЛЛЕГИЯ:

АДЕКЕНОВ Сергазы Мынжасарович, доктор химических наук, профессор, академик НАН РК, директор АО «Научно-производственного центра «Фитохимия» (Караганда, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=7006153118>, <https://www.webofscience.com/wos/author/record/48648658>

РАМАЗАНОВ Тлеккабул Сабитович, (заместитель главного редактора), доктор физико-математических наук, профессор, академик НАН РК (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6701328029>, <https://www.webofscience.com/wos/author/record/13503476>

АБИЕВ Руфат, доктор технических наук (биохимия), профессор, заведующий кафедрой «Оптимизация химической и биотехнологической аппаратуры», Санкт-Петербургский государственный технологический институт (Санкт-Петербург, Россия), <https://www.scopus.com/authid/detail.uri?authorId=6602431781>, <https://www.webofscience.com/wos/author/record/1405661>

ОЛИВЬЕРО Россини Чезаре, доктор философии (PhD, химия), профессор Университета Калабрии (Калабрия, Италия), <https://www.scopus.com/authid/detail.uri?authorId=57221375979>, <https://www.webofscience.com/wos/author/record/399768>

ТИГИНЯНУ Ион Михайлович, доктор физико-математических наук, академик, президент Академии наук Молдовы, Технический университет Молдовы (Кишинев, Молдова), <https://www.scopus.com/authid/detail.uri?authorId=7006315935>, <https://www.webofscience.com/wos/author/record/524462>

САНГ-СУ Квак, доктор философии (PhD, биохимия, агрохимия), профессор, главный научный сотрудник, Научно-исследовательский центр инженерных систем растений, Корейский научно-исследовательский институт бионауки и биотехнологии (KRIBB) (Дэчон, Корея), <https://www.scopus.com/authid/detail.uri?authorId=59286321700>, <https://www.webofscience.com/wos/author/record/30028581>

БЕРСИМБАЕВ Рахметкажи Искендерович, доктор биологических наук, профессор, академик НАН РК, Евразийский национальный университет им. Л.Н. Гумилева (Астана, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=7004012398>, <https://www.webofscience.com/wos/author/record/19854255>

КАЛАНДРА Пьетро, доктор философии (PhD, физика), профессор Института по изучению наноструктурированных материалов (Рим, Италия), <https://www.scopus.com/authid/detail.uri?authorId=7004303066>, <https://www.webofscience.com/wos/author/record/616360>

БОШКАЕВ Куантай Авгазиевич, PhD, преподаватель, доцент кафедры теоретической и ядерной физики, Казахский национальный университет им. Аль-Фараби (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=54883880400>, <https://www.webofscience.com/wos/author/record/2080231>

БУРКИТБАЕВ Мухамбетали, доктор химических наук, профессор, академик НАН РК, (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=8513885600>, <https://www.webofscience.com/wos/author/record/29017135>

QUEVEDO Hernando, профессор, Национальный автономный университет Мексики (UNAM), Институт ядерных наук (Мехико, Мексика), <https://www.scopus.com/authid/detail.uri?authorId=55989741100>, <https://www.webofscience.com/wos/author/record/30353742>

ЖУСУПОВ Марат Абжанович, доктор физико-математических наук, профессор кафедры теоретической и ядерной физики, Казахский национальный университет имени аль-Фараби (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6602166928>, <https://www.webofscience.com/wos/author/record/566>

КОВАЛЕВ Александр Михайлович, доктор физико-математических наук, академик НАН Украины, Институт прикладной математики и механики (Донецк, Украина), <https://www.scopus.com/authid/detail.uri?authorId=7202799321>, <https://www.webofscience.com/wos/author/record/65533963>

ТАКИББАЕВ Нурғали Жабағевич, доктор физико-математических наук, профессор, академик НАН РК, Казахский национальный университет им. Аль-Фараби (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=24077239000>, <https://www.webofscience.com/wos/author/record/1671760>

ХАРИН Станислав Николаевич, доктор физико-математических наук, профессор, академик НАН РК (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6701353063>, <https://www.webofscience.com/wos/author/record/2023295>

АБИШЕВ Медеу Ержанович, доктор физико-математических наук, профессор, академик НАН РК, (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=26530759900>, <https://www.webofscience.com/wos/author/record/1556025>

ACADEMIC JOURNAL OF PHYSICAL AND CHEMICAL SCIENCES

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

Собственник: ТОО «Центрально-азиатский академический научный центр» (г. Алматы).

Свидетельство № KZ93VPY00121157 о повторной регистрации периодического печатного издания информационного агентства, информационного агентства и сетевого издания, выданное Республиканским государственным учреждением «Комитет информации» Министерства культуры и информации Республики Казахстан **05.06.2025**

Тематическая направленность: *физика, химия*.

Периодичность: 4 раза в год.

<http://reports-science.kz/index.php/en/archive>

CONTENTS

PHYSICS

Aimaganbetova Z.K., Kulshymbayev Y.A., Zhanturina N.N., Beketova G.K.
 First-principles calculation of the electronic properties of the Double Halide
 Perovskite $\text{Cs}_2\text{Ag}_{0.2}\text{Na}_{0.4}\text{In}_{0.6}\text{Ti}_{0.4}\text{Cl}_6$ based on the quantum ESPRESSO software.....14

Amangeldinova S., Zhuniskhan S., Kalzhigitov N., Kurmangaliyeva V.
 Study of the cluster structure of ^5He and ^5Li mirror nuclei in two-cluster
 approximation.....35

Chokin K., Otunchi Ye., Kozhahmetova A., Kasenova A., Shongalova A.
 Development and testing of a laboratory pyrometallurgical installation for recycling
 lithium-ion batteries.....46

Issayeva A., Beisebayeva A., Madybekova G., Shynazbekova Sh., Issa A.
 Comparative analysis of physico-chemical characteristics of drinking, spring
 and natural water in the South Kazakhstan.....65

Kim V.Yu., Aimuratov Y.K.
 Search for transient cosmic events by scanning the sky with wide-field telescopes.....78

Koshtybayev T.B., Tatenov A.M., Aliyeva M.E., Tugelbaeva G.T., Zhanaliyeva G.Zh.
 Study of the electromagnetic field based on thermodynamics principles.....89

Mukamedenkyzy V., Akberdiyev B.
 Numerical investigation of the effect of inclination angle on the stability
 of mechanical equilibrium in Ar–N₂ binary gas mixtures.....105

Myasnikova L.N., Uzakbaeva S.S., Shanina Z.K., Bekeshev A.Z.
 Kinetic properties of high-density polyethylene filled with chromium spinel powder.....119

Nurbayev B.M., Dmitriyeva E.A., Kemelbekova A.E.
 The role of low-dimensional layered structures in enhancing the stability of tin-based
 perovskite materials.....136

Sattinova Z., Ermakhanova F., Assilbekov B., Taimuratova L.
 Influence of various cooling conditions and heat transfer coefficients on solidification
 during the formation of beryllium ceramic products.....149

Shestakova L.I., Serebryanskiy A.V., Spassyyuk R.R., Omarov Ch.T.
 Search for gas of comet-meteor origin in the inner Solar System: caii ion emission.....165

Ualikhanova U., Tursynkazy F., Syzdykova A.M., Altayeva G.S., Altaibayeva A.B.
 Studying the amplitude of $f(T)$ gravitational waves using Bessel functions.....179

Zhexenbayeva G.A., Nasirova D.M., Aimanova G.K., Shomshekova S.A. Photometric study of the symbiotic object V725 Tau.....	194
Zhusupova N.K., Zhadyranova A.A. Bounce cosmology in $f(T, \mathcal{T})$ gravity based on energy condition analysis.....	205
Ziyatbekova G., Abdimanapova P., Sagyntay O., Nurym A., Ilinov R. Using artificial intelligence to predict diseases based on medical data.....	225

CHEMISTRY

Almassov N.Zh., Zhumagaliyeva A.N., Duisenbekov S.E., Zhakiyev N.K. Design and optimization of hybrid renewable energy systems for hydrogen production in Kazakhstan.....	236
Amangeldi B., Zhanikulov N., Taimasov B., Aitureev M.M., Dauletiyarov M. Calculation of the Raw material composition for obtaining white Portland cement clinker.....	251
Baeshov A., Tashenov E.A., Atykhanova S.B., Koshkarbayeva Sh.T. Preparation of cadmium sulfide by electrochemical method using a composite sulfur-graphite electrode.....	267
Baisalova G.Zh., Azhikhanova Zh., Taltenov A.A., Kuzhatova P. Determination of the total phenolic content in perennial herbaceous plants of the flora of Kazakhstan.....	277
Darmenbayeva A.S., Rajasekharan R., Zhussipnazarova G.M., Mukazhanova Zh.B., Begenova B.E. Composites based on chitosan and cellulose: synthesis, properties, and application prospects.....	287
Erkasov R.Sh., Zhamkenova A.S., Sergazina S.M., Nurmukhanbetova N.N., Kassenova N.B. Halide-dependent modulation of hydrogen bonding in Mn(II) complexes with protonated acetamide: a QAIM, NCI, and energy decomposition study.....	304
Kalimoldina L.M., Shaikhova Zh.E., Kaliyeva B.K., Bubish Sh., Askarova Sh.K. The effect of silver nanoparticles on the germination of bean, lemon, tangerine and avocado seeds.....	320
Kurtebayeva A.A., Alvarez-Torrellas S., Gomes H.T., Orynbayev S.A., Kalmakhanova M.S. Activated-carbon-enhanced polymeric membranes for efficient elimination of emerging contaminants.....	334

Massenova A.T., Zhumakanova A.S., Torlopov I.I., Rakhmetova K.S., Abilmagzhanov A.Z. Optimization of the hierarchical zeolite ZSM-5 synthesis process by steam-assisted alkaline modification.....	350
Mutushev A.Zh., Nuraly A.M., Sanat A.S., Shaukharova M.A., Yessimsiitova Z.B. The effect of light-converting films on the accumulation of bioactive compounds and the quality of fruits.....	366
Nefedov A.N., Taikenova A.T. Current state of organic corrosion inhibitor application in oil refining.....	379
Omarov B.T., Altybayev Zh.M., Serikbayeva B.S. Production of biohumus by vermicomposting of organic wastes and study of its agroecological effectiveness.....	399
Rakhman D.M., Kappasuly A., Makhayeva D.N., Kazybayeva D.S., Irmukhametova G.S. Development and investigation of mucoadhesive hydrogels based on gellan–cysteine complexes.....	414
Sabyrzhanova A.E., Bolatkyzy N., Berganaeva G.E., Dyusebaeva M.A. Study of amino acids and fatty acids in the aerial part of <i>Plantago major</i>	428
Satayeva S., Akhmetova F., Urazova A., Aituganova S., Yerniyazova K. The influence of PEPA concentration on the physical, mechanical, and operational properties of ED-20 epoxy adhesives.....	439
Zamanbek A.Zh., Koshkarbayeva Sh.T., Satayev M.S. Methods of Obtaining Silver Nanoparticles and Antibacterial Properties.....	450
Zhortarova A.A., Salkeyeva L.K., Minayeva Ye.V., Ibrayev M.K., Fazylov S.D. New possibilities for the synthesis and phosphorylation of phosphonoacetic acid ester.....	462

МАЗМҰНЫ

ФИЗИКА

Аймағанбетова З.К., Құлшымбаев Е.А., Жантурина Н.Н., Бекетова Г.К. Quantum Espresso бағдарламасы негізінде Cs ₂ Ag _{0.2} Na _{0.4} In _{0.6} Ti _{0.4} Cl ₆ кос галогенді перовскиттің электрондық қасиеттерін бірінші принциптік есептеу.....	14
Амангелдинова С., Жүнісхан С., Калжигитов Н., Курманғалиева В. Екі кластерлік жуықтауда 5He және 5Li айналық ядроларының кластерлік құрылымын зерттеу.....	35
Чокин К., Отунчи Е., Кожаметова А., Касенова А., Шонғалова А. Литий-ионды аккумуляторларды қайта өңдеуге арналған зертханалық пирометаллургиялық қондырғыны әзірлеу және сынау.....	46
Исаева А.Б., Бейсебаева А.С., Мадыбекова Г.М., Шиназбекова Ш.С., Иса А.Б. Сравнительный анализ физико-химических характеристик питьевой, родниковой и природной воды юга Казахстана.....	65
Ким В.Ю., Аймуратов Е.К. Кең бұрышты телескоптармен аспанды сканерлеу арқылы өтпелі ғарыштық оқиғаларды іздеу.....	78
Коштыбаев Т.Б., Татенов А.М., Алиева М.Е., Тугелбаева Г.Т., Жаналиева Г.Ж. Электромагниттік өрісті термодинамикалық бастамалар тұрғысында зерттеу.....	89
Мукамеденқызы В., Ақбердиев Б. Ar–N ₂ бинарлы газ қоспаларындағы механикалық тепе-теңдік тұрақтылығына қиғаш бұрыштың әсерін сандық зерттеу.....	105
Мясникова Л.Н., Узакбаева С.С., Шанина З.К., Бекешев А.З. Хром-шпинельді ұнтақ қосылған жоғары тығыздықты полиэтиленнің кинетикалық қасиеттері.....	119
Нұрбаев Б.М., Дмитриева Е.А., Кемелбекова А.Е. Қалайы негізіндегі перовскитті материалдардың тұрақтылығын арттырудағы төменөлшемді қабатты құрылымдардың рөлі.....	136
Саттинова З., Ермаханова Ф., Асылбеков Б., Таймуратова Л. Бериллий керамикалық бұйымдарын қалыптастыру кезінде әр түрлі салқындату жағдайлары мен жылу беру коэффициенттерінің қатаюға әсері.....	149
Шестакова Л.И., Серебрянский А.В., Спасюк Р.Р., Омаров Ш.Т. Күн жүйесінің ішкі аймағындағы комета-метеорлық тектегі газды іздеу: CaII иондарының жарқырауы.....	165

Уалиханова У.А., Тұрсынқазы Ф., Сыздықова А.М., Алтаева Г.С., Алтайбаева А.Б.
Бессель функцияларын пайдаланып $f(T)$ гравитациялық толқындардың
амплитудасын зерттеу.....179

Жексенбаева Г.А., Насирова Д.М., Айманова Г.К., Шомшекова С.А
V725 Тау симбиотикалық объектiсiн фотометрлiк зерттеу.....194

Жусупова Н.К., Жадыранова А.А.
Энергия шарттарын талдауға негiзделген $f(T, T)$ серпiлiс космологиясы.....205

Зиятбекова Г.З., Абдиманапова П.Б., Сағынтай О.А., Нұрым А.А., Ильинов Р.А.
Жасанды интеллект көмегiмен медициналық деректер бойынша
ауруларды болжау.....225

ХИМИЯ

Алмасов Н.Ж., Жумагалиева А.Н., Дүйсенбеков С.Е., Жакиев Н.К.
Қазақстанда сутегi өндiруге арналған гибрирдiк жаңартылатын энергия жүйелерiн
жобалау және оңтайландыру.....236

Амангелдi Б., Жаникулов Н., Таймасов Б., Айтуреев М., Даулетияров М.
Ақ портландцемент клинкерiн алу үшiн шикiзат шихта құрамын есептеу.....251

Баешов А., Ташенов Е.А., Атыханова С.Б., Кошкарбаева Ш.Т.
Композициялы күкiрт-графит электродын қолдану арқылы кадмий
сульфидiн электрохимиялық әдiспен алу.....267

Байсалова Г.Ж., Ажиханова Ж., Талтенов А.А., Құжатова П.
Қазақстан флорасындағы көпжылдық шөптесiн өсiмдiктердiң фенолдық
қосылыстарының жиынтық мөлшерiн анықтау.....277

**Дарменбаева А.С., Rajasekharan R., Жусипназарова Г.М., Мукажанова Ж.Б.,
Бегенова Б.Е.**
Хитозан және целлюлоза негiзiндегi композиттер: синтез, қасиеттерi және қолдану
перспективалары.....287

**Еркасов Р.Ш., Жамкенова А.С., Сергазина С.М., Нурмуханбетова Н.Н.,
Касенова Н.Б.**
Mn (II) кешендерiндегi сутектiк байланыстардың энергиясы мен табиғатына
галогеннiң әсерi: QTAIM, NCI және энергия декомпозициясы.....304

Калимолдина Л.М., Шаихова Ж.Е., Калиева Б.К., Бубиш Ш., Аскарова Ш.К.
Күмiс нанобөлшектерiнiң бұршақ, лимон, мандарин, авокадо тұқымдарының
өнуiне әсерi.....320

Қуртебаева А.А., Álvarez-Torrellas S., Gomes Н.Т., Орынбаев С.Ә., Калмаханова М.С. Алаңдаушылық тудыратын ластаушы заттарды тиімді жою үшін белсендірілген көмір полимерлі мембраналар.....	334
Масенова А.Т., Жұмақанова А.С., Торлопов И.И., Рахметова К.С., Абильмагжанов А.З. ZSM-5 иерархиялық цеолитін бумен сілтілі модификациялау арқылы алу процесін онтайландыру.....	350
Мутушев А.Ж., Нұралы Ә.М., Санат А.С., Шаукарова М.А., Есимситова З.Б. Жарық түрлендіретін пленкалардың биоактивті қосылыстардың жинақталуына және жеміс сапасына әсері.....	366
Нефедов А.Н., Тайекенова А.Т. Мұнай өңдеу өнеркәсібінде органикалық коррозия ингибиторларын қолданудың қазіргі жағдайы.....	379
Омаров Б.Т., Алтыбаев Ж.М., Серикбаева Б.С. Органикалық қалдықтарды вермикомпостинг арқылы биогумус өндіру және оның агроэкологиялық тиімділігін зерттеу.....	399
Рахман Д.М., Қаппасұлы Ә., Махаева Д.Н., Қазыбаева Д.С., Ирмухаметова Ғ.С. Геллан–цистеин кешендері негізінде мукоадгезиялық гидрогельдерді әзірлеу және зерттеу.....	414
Сабыржанова А.Е., Болатқызы Н., Берганаева Г.Е., Дюсебаева М.А. Plantago Major жер үсті бөлігінің құрамындағы амин қышқылдары мен май қышқылдарын зерттеу.....	428
Сатаева С., Ахметова Ф., Уразова А., Айтуганова С., Ерниязова К. ЭД-20 эпоксидті желімдерінің физика-механикалық және эксплуатациялық қасиеттеріне ПЭПА концентрациясының ықпалы.....	439
Заманбек А.Ж., Кошкарбаева Ш.Т., Сатаев М.С. Күміс нанобөлшектерінің алыну әдістері мен антибактериалдық қасиеттері.....	450
Жоргарова А.А., Салькева Л.К., Минаева Е.В., Ибраев М.К., Фазылов С.Д. Фосфоносірке қышқылының эфирін синтездеу мен фосфорландырудың жаңа мүмкіндіктері.....	462

СОДЕРЖАНИЕ

ФИЗИКА

Аймаганбетова З.К., Кулшымбаев Е.А., Жантурина Н.Н., Бекетова Г.К. Расчет по первому принципу электронных свойств двойного галогенидного перовскита Cs ₂ Ag _{0.2} Na _{0.4} In _{0.6} Ti _{0.4} Cl ₆ на основе программы Quantum Espresso.....	14
Амангелдинова С., Жүнісхан С., Калжигитов Н., Курмангалиева В. Исследование кластерной структуры зеркальных ядер ⁵ He и ⁵ Li в двухкластерном приближении.....	35
Чокин К., Отунчи Е., Кожаметова А., Касенова А., Шонгалова А. Разработка и испытания лабораторной пирометаллургической установки для переработки литий-ионных аккумуляторов.....	46
Исаева А.Б., Бейсебаева А.С., Мадыбекова Г.М., Шиназбекова Ш.С., Иса А.Б. Сравнительный анализ физико-химических характеристик питьевой, родниковой и природной воды юга Казахстана.....	65
Ким В.Ю., Аймуратов Е.К. Поиск транзиентных космических событий методом сканирования неба широкоугольными телескопами.....	78
Коштыбаев Т.Б., Татенов А.М., Алиева М.Е., Тугелбаева Г.Т., Жаналиева Г.Ж. Исследование электромагнитного поля на основе термодинамических принципов.....	89
Мукамеденкызы В., Акбердиев Б. Численное исследование влияния угла наклона на устойчивость механического равновесия в бинарной газовой смеси Ar–N ₂	105
Мясникова Л.Н., Узакбаева С.С., Шанина З.К., Бекешев А.З. Кинетические свойства высокоплотного полиэтилена с добавлением хром-шпинельного порошка.....	119
Нурбаев Б.М., Дмитриева Е.А., Кемелбекова А.Е. Роль низкоразмерных слоистых структур в повышении стабильности перовскитных материалов на основе олова.....	136
Саттинова З., Ермаханова Ф., Асылбеков Б., Таймуратова Л. Влияние различных условий охлаждения и коэффициентов теплопередачи на затвердевание при формировании бериллиевых керамических изделий.....	149

Шестакова Л.И., Серебрянский А.В., Спасюк Р.Р., Омаров Ш.Т. Поиск газа кометно-метеорного происхождения во внутренней области Солнечной Системы: Свечение ионов СаII.....	165
Уалиханова У.А., Турсынказы Ф., Сыздыкова А.М., Алтаева Г.С., Алтайбаева А.Б. Изучение амплитуды $f(T)$ гравитационных волн с использованием функций Бесселя.....	179
Жексенбаева Г.А., Насирова Д.М., Айманова Г.К., Шомшекеева С.А. Фотометрическое исследование симбиотического объекта V725 Tau.....	194
Жусупова Н.К., Жадыранова А.А. Космология отскока в $f(T, \mathcal{J})$ гравитации на основе анализа энергетических условий.....	205
Зиятбекова Г.З., Абдимананова П.Б., Сағынтай О.А., Нұрым А.А., Ильинов Р.А. Использование искусственного интеллекта для прогнозирования заболеваний на основе медицинских данных.....	225

ХИМИЯ

Алмассов Н.Ж., Жумагалиева А.Н., Дуйсенбеков С.Е., Жакиев Н.К. Проектирование и оптимизация гибридных возобновляемых источников энергии для производства водорода в Казахстане.....	236
Амангелді Б., Жаникулов Н., Таймасов Б., Айтуреев М., Даулетияров М. Расчёт состава сырьевой шихты для получения белого порландцементного клинкера.....	251
Башов А., Ташенов Е.А., Атыханова С.Б., Кошкарбаева Ш.Т. Получение сульфида кадмия электрохимическим методом с использованием композитного сера-графитового электрода.....	267
Байсалова Г.Ж., Ажиханова Ж., Талтенов А.А., Кужатова П. Определение суммы фенольных соединений в многолетних травянистых растениях флоры Казахстана.....	277
Дарменбаева А.С., Rajasekharan R., Жусиппазарова Г.М., Мукажанова Ж.Б., Бегенова Б.Е. Композиты на основе хитозана и целлюлозы: синтез, свойства и перспективы применения.....	287
Еркасов Р.Ш., Жамкенова А.С., Сергазина С.М., Нурмуханбетова Н.Н., Касенова Н.Б. Влияние галогена на энергетику и природу водородных связей в Mn(II): QTAIM, NCI и энергодекомпозиция.....	304

Калимолдина Л.М., Шаихова Ж.Е., Калиева Б.К., Бубиш Ш., Аскарова Ш.К. Влияние наночастиц серебра на прорастание семян фасоли, лимона, мандарина, авокадо.....	320
Куртебаева А.А., Álvarez-Torrellas S., Gomes Н.Т., Орынбаев С.А., Калмаханова М.С. Полимерные мембраны с активированным углем для эффективного удаления загрязняющих веществ вызывающих обеспокоенность.....	334
Масенова А.Т., Жұмақанова А.С., Торлопов И.И., Рахметова К.С., Абильмагжанов А.З. Оптимизация процесса получения иерархического цеолита ZSM-5 паровой щелочной модификацией.....	350
Мутушев А.Ж., Нуралы А.М., Санат А.С., Шаукарова М.А., Есимсиитова З.Б. Влияние светопреобразующих плёнок на накопление биоактивных соединений и качество плодов.....	366
Нефедов А.Н., Тайекенова А.Т. Современное состояние применения органических ингибиторов коррозии в нефтепереработке.....	379
Омаров Б.Т., Алтыбаев Ж.М., Серикбаева Б.С. Получение биогумуса путем вермикомпостирования органических отходов и исследование его агроэкологической эффективности.....	399
Рахман Д.М., Қаппасұлы Ә., Махаева Д.Н., Казыбаева Д.С., Ирмухаметова Г.С. Разработка и исследование мукоадгезивных гидрогелей на основе комплексов геллан–цистеин.....	414
Сабыржанова А.Е., Болаткызы Н., Берганаева Г.Е., Дюсебаева М.А. Исследование аминокислот и жирных кислот в составе надземной части <i>Plantago Major</i>	428
Сатаева С., Ахметова Ф., Уразова А., Айтуганова С., Ерниязова К. Влияние концентрации ПЭПА на физические, механические и эксплуатационные свойства эпоксидных клеев ЭД-20.....	439
Заманбек А.Ж., Кошкарбаева Ш.Т., Сатаев М.С. Методы получения наночастиц серебра и антибактериальные свойства.....	450
Жоргарова А.А., Салькева Л.К., Минаева Е.В., Ибраев М.К., Фазылов С.Д. Новые возможности синтеза и фосфорилирования фосфонуксусного эфира.....	462

ACADEMIC JOURNAL
OF PHYSICAL AND CHEMICAL SCIENCES
ISSN 2224-5227
Volume 1.
Number 357 (2026), 165–178

<https://doi.org/10.32014/2026.2518-1483.412>

UDC: 523.62-726; 523.62-65; 523.62
IRSTI: 41.21.25; 89.15.35; 89.51.33

©**Shestakova L.I., Serebryanskiy A.V., Spassyyuk R.R.*, Omarov Ch.T., 2026.**

Fesenkov Astrophysical Institute LLP, Almaty, Kazakhstan.

*E-mail: spassyyuk@fai.kz

SEARCH FOR GAS OF COMET-METEOR ORIGIN IN THE INNER SOLAR SYSTEM: CaII ION EMISSION

Shestakova Lyubov — PhD, Fesenkov Astrophysical Institute, Almaty, Kazakhstan,

E-mail: shest1952@mail.ru, <https://orcid.org/0000-0002-2223-5332>;

Serebryanskiy Alexander — PhD, Fesenkov Astrophysical Institute, Almaty, Kazakhstan,

E-mail: aserebryanskiy@gmail.com, <https://orcid.org/0000-0002-4313-7416>;

Spassyyuk Ruslan — Junior researcher, Fesenkov Astrophysical Institute, Almaty, Kazakhstan,

E-mail: spassyyuk@fai.kz, <https://orcid.org/0000-0002-7780-2533>;

Omarov Chingis — PhD, Professor, Fesenkov Astrophysical Institute, Almaty, Kazakhstan,

E-mail: chingis.omarov@fai.kz, <https://orcid.org/0000-0002-1672-894X>.

Abstract. Interplanetary dust of asteroidal and cometary origin gradually drifts toward the Sun under the drag produced by solar radiation and the solar wind. Upon reaching the sublimation zone, the dust evaporates and breaks down into atoms, which are then ionized and may subsequently be carried away by the solar wind. Some atoms and singly charged ions experience resonant-line radiation pressure that significantly exceeds the gravitational pull of the Sun. Elements of this kind can be accelerated by radiation pressure to high velocities, provided that their lifetime before the next ionization event is sufficiently long. Gas clouds formed by the disintegration of compact objects such as comets or meteoroid streams in the vicinity of the Sun may be separated by radiation pressure into several clouds with homogeneous chemical composition and travel outward as far as Earth's orbit. Previous calculations have shown that a number of atoms and singly charged ions can attain high velocities within their lifetime before further ionization. Solar radiation acts most efficiently on the singly ionized species of magnesium and calcium. Earlier studies demonstrated that emission in the H and K lines of CaII can be detected during solar eclipses using highly sensitive, high spectral resolution instruments. Such unique observations have been accomplished only twice, in the works of Gulyaev and Shcheglov during the solar eclipses of 26 February 1998 and 11 August 1999. The difficulty of these observations is compounded by the short duration of eclipses and the presence of atmospheric disturbances. Observations made outside Earth's atmosphere using orbital instruments are free from such limitations.

Given the presence of inhomogeneities and density enhancements in the form of meteoroid streams within the inner Solar System, the likelihood of detecting clouds containing distinct groups of atoms or ions increases.

Keywords: dust, gas, meteoroid streams, resonance emission, calcium atoms and ions

Funding. *The work is carried out within the framework of the Project No. BR24992759 Development of the concept for the first Kazakhstani orbital cislunar telescope - Phase I”, financed by the Ministry of Science and Higher Education of the Republic of Kazakhstan.*

For citations: *Shestakova L.I., Serebryanskiy A.V., Spassyyuk R.R., Omarov Ch.T. Search for Gas of Comet-Meteor Origin in the Inner Solar System: CaII Ion Emission. Academic Journal of Physical and Chemical Sciences. 2026. No.1. Pp. 165–178. DOI: <https://doi.org/10.32014/2026.2518-1483.412>*

©Шестакова Л.И., Серебрянский А.В., Спасюк Р.Р.*, Омаров Ш.Т., 2026.

«В.Г. Фесенков атындағы Астрофизикалық институты» ЖШС,

Алматы, Қазақстан.

*E-mail: spassyyuk@fai.kz

КҮН ЖҮЙЕСІНІҢ ІШКІ АЙМАҒЫНДАҒЫ КОМЕТА-МЕТЕОРЛЫҚ ТЕКТЕГІ ГАЗДЫ ІЗДЕУ: САП ИОНДАРЫНЫҢ ЖАРҚЫРАУЫ

Шестакова Любовь — PhD, В.Г. Фесенков атындағы Астрофизикалық институты, Алматы, Қазақстан,

E-mail: shest1952@mail.ru, <https://orcid.org/0000-0002-2223-5332>;

Серебрянский Александр — PhD, В.Г. Фесенков атындағы Астрофизикалық институты, Алматы, Қазақстан,

E-mail: aserebryanskiy@gmail.com, <https://orcid.org/0000-0002-4313-7416>;

Спасюк Руслан — Кіші ғылыми қызметкер, В.Г. Фесенков атындағы Астрофизикалық институты, Алматы, Қазақстан,

E-mail: spassyyuk@fai.kz, <https://orcid.org/0000-0002-7780-2533>;

Омаров Шыңғыс — PhD, профессор, В.Г. Фесенков атындағы Астрофизикалық институты, Алматы, Қазақстан,

E-mail: chingis.omarov@fai.kz, <https://orcid.org/0000-0002-1672-894X>.

Аннотация. Астероидтық және кометалық шығу текті планетааралық шаң Күн сәулеленуі мен күн желі әсерінен болатын тежелулердің ықпалымен біртіндеп Күнге жақындайды. Сублимация аймағына жеткен кезде шаң буланып, атомдарға ыдырайды, олар кейін иондалып, әрі қарай күн желімен ілесіп әкетілуі мүмкін. Кейбір атомдар мен алғашқы иондар резонанстық сызықтардағы сәулелік қысымды сезінеді, ол Күннің тартылыс күшінен едәуір асып түседі. Мұндай элементтер, егер келесі иондалуға дейінгі өмір сүру уақыты жеткілікті ұзақ болса, сәулелік қысым әсерінен жоғары жылдамдықтарға дейін үдетілуі мүмкін. Күнге жақын маңда ықшам нысандардың кометалардың немесе метеор ағындарының



ыдырауы нәтижесінде түзілген газ бұлттары сәулелік қысым әсерінен химиялық құрамы біртекті бірнеше бұлтқа бөлініп, Жер орбитасына дейін жете алады. Бұрын жүргізілген есептеулер келесі иондалуға дейінгі өмір сүру уақыты ішінде жоғары жылдамдықтарға жете алатын бірқатар атомдар мен алғашқы иондардың бар екенін көрсетті. Күн радиациясының әсері Mg және Ca элементтерінің бірінші иондарына ең тиімді түрде әсер етеді. Бұрын жасалған зерттеулер Ca II-дің H және K сызықтарының сәуле шығаруын жоғары сезімталдыққа және жоғары спектралдық айырымдылыққа ие аспаптарды пайдалана отырып, күн тұтылулары кезінде тіркеуге болатынын көрсетті. Мұндай бірегей зерттеулер бар болғаны екі рет Гуляев пен Щегловтың еңбектерінде 1998 жылғы 26 ақпандағы және 1999 жылғы 11 тамыздағы күн тұтылулары кезінде жүзеге асырылды. Мұндай зерттеулердің күрделілігі тұтылулардың қысқа мерзімділігімен және атмосфералық кедергілердің болуымен күшейе түседі. Жер атмосферасынан тыс, орбиталық аппараттардың көмегімен жүргізілетін бақылауларда мұндай шектеулер жоқ. Ішкі Күн жүйесінде метеор ағындары түріндегі біртектес еместіктер мен тығыздалулардың болуын ескерсек, жекелеген атомдар немесе иондар топтарын қамтитын бұлттарды анықтау ықтималдығы арта түседі.

Түйін сөздер: шаң, газ, метеорлық ағындар, резонанстық жарқырау, Ca атомдары мен иондары

©Шестакова Л.И., Серебрянский А.В., Спасюк Р.Р.*, Омаров Ш.Т., 2026.

ТОО «Астрофизический институт имени В.Г. Фесенкова»,

Алматы, Казахстан.

*E-mail: spassyuk@fai.kz

ПОИСК ГАЗА КОМЕТНО-МЕТЕОРНОГО ПРОИСХОЖДЕНИЯ ВО ВНУТРЕННЕЙ ОБЛАСТИ СОЛНЕЧНОЙ СИСТЕМЫ: СВЕЧЕНИЕ ИОНОВ Ca II

Шестакова Любовь — PhD, Астрофизический институт имени В.Г. Фесенкова, Алматы, Казахстан,
E-mail: shest1952@mail.ru, <https://orcid.org/0000-0002-2223-5332>;

Серебрянский Александр — PhD, Астрофизический институт имени В.Г. Фесенкова, Алматы, Казахстан,

E-mail: aserebryanskiy@gmail.com, <https://orcid.org/0000-0002-4313-7416>;

Спасюк Руслан — Младший научный сотрудник, Астрофизический институт имени В.Г. Фесенкова, Алматы, Казахстан,

E-mail: spassyuk@fai.kz, <https://orcid.org/0000-0002-7780-2533>;

Омаров Чингис — PhD, Астрофизический институт имени В.Г. Фесенкова, Алматы, Казахстан,
E-mail: chingis.omarov@fai.kz, <https://orcid.org/0000-0002-1672-894X>.

Аннотация. Межпланетная пыль астероидного и кометного происхождения постепенно приближается к Солнцу под действием торможения солнечным излучением и солнечным ветром. При достижении области сублимации пылевые частицы испаряются и распадаются на атомы, которые затем ионизируются и далее могут увлекаться солнечным ветром. Некоторые атомы и однократно ионизованные ионы испытывают действие светового давления в резонансных

линиях, значительно превышающего силу солнечного тяготения. Такие элементы могут разгоняться световым давлением до больших скоростей, если время их жизни до следующей ионизации достаточно велико. Облака газа, образовавшиеся в результате распада вблизи Солнца компактных объектов - комет или метеорных потоков, — могут под действием светового давления разделяться на несколько облаков с однородным химическим составом и достигать орбиты Земли. Выполненные ранее расчеты показали, что существует ряд атомов и однократно ионизованных ионов, способных достичь высоких скоростей за время своего существования до следующей ионизации. Наиболее эффективно солнечная радиация воздействует на однократно ионизованные ионы Mg и Ca. Ранее проведенные исследования показали, что регистрация свечения линий H и K Ca II возможна во время солнечных затмений с использованием высокочувствительных приборов, обладающих высоким спектральным разрешением. Такие уникальные исследования были реализованы лишь дважды - в работах Гуляева и Щеглова во время солнечных затмений 26 февраля 1998 года и 11 августа 1999 года. Трудность подобных исследований усугубляется кратковременностью затмений и наличием атмосферных помех. Наблюдения за пределами земной атмосферы с помощью орбитальных аппаратов таких ограничений не имеют. С учетом наличия во внутренней Солнечной системе неоднородностей и уплотнений в виде метеорных потоков возрастает вероятность обнаружения облаков, содержащих отдельные группы атомов или ионов.

Ключевые слова: пыль, газ, метеорные потоки, резонансное свечение, атомы и ионы Ca

Introduction. The behavior of interplanetary dust is governed by the Poynting-Robertson drag effect, whose action leads to the fact that dust particles initially moving on circular orbits begin to approach the Sun along a spiral. It is obvious that the dust reaches the sublimation region, where it is heated to high temperatures and evaporates. For a long time, experimental investigation of this region was not carried out, since the near-solar region is a rather difficult object for observations. Attention was first drawn to the dust sublimation region near the Sun after the reports by (Peterson et al., 1967; MacQueen et al., 1968) on the detection of local maxima of dust thermal emission at a wavelength of 2.2 μm at heliocentric distances of 3.5, 4.0, 8.7, and 9.2 R_{\odot} near the ecliptic plane, symmetrically with respect to the Sun.

In the review by (Mann et al., 2004), an extensive list of infrared (IR) observations of the F-corona is presented, including both indications of the presence of thermal emission and its absence. IR observations revealed no signs of thermal radiation at distances up to 15 R_{\odot} during the solar eclipses of 1980 and 1991. At the same time, during the eclipse of 11 June 1983 an intense maximum of IR emission was detected to the west of the Sun, with a complete absence of thermal emission on the eastern side. Apparently, we are dealing with non-stationary dust formations clouds in the near-solar region.

It is obvious that dust in the sublimation region evaporates and is converted into a gas whose temperature is several orders of magnitude lower than the temperature of the

surrounding gas of the solar corona. The atoms of the resulting gas will actively interact with solar radiation and coronal particles. The relaxation time of atoms and primary ions with the coronal gas due to collisions is sufficiently large at distances from $4R_{\odot}$ to $10R_{\odot}$ compared with their lifetime, which is determined by ionization under the action of the Sun's ultraviolet (UV) radiation.

As for secondary ions, their lifetime until the next ionization may exceed the relaxation time with the coronal gas, and if these ions have not left the near-solar region, then they mix with the corona. Further ionization then proceeds mainly due to collisions with coronal particles, since the potential of each subsequent ionization increases to such an extent that the probability of ionization by radiation becomes much smaller because of the lack of high-energy quanta in the solar emission spectrum.

The process of transformation of cold gas into typically coronal gas can be considered complete after the onset of ionization equilibrium, which, under solar coronal conditions, occurs at high stages of ionization. Evidence for the presence in the near-solar region of gas with low degrees of ionization, not typical of the solar corona, can be regarded as consequences of the injection into the near-solar region of interplanetary material in the form of a flux of meteoroids, comets, or zodiacal dust.

What are the specific characteristics of the interaction between neutral atoms (and primary ions) and solar radiation prior to further ionization? Firstly, many neutral atoms possess resonance lines corresponding to transitions from the ground state in spectral regions near the peak of solar radiation. Certain atomic transitions may have significant oscillator strengths, experiencing radiation pressure in resonance lines that exceeds the Sun's gravitational pull. This allows specific atoms to exit the circumsolar region and accelerate to high velocities, potentially reaching solar escape velocity. Since the lifetime of primary ions significantly exceeds that of neutral atoms, the highest velocities are attained by elements that exhibit high radiation pressure in both their neutral and singly ionized states.

The first estimates of the intensity of resonance emission of some atoms and primary ions near the dust sublimation region were made in (Shestakova et al., 1990). That work concluded that Mg and Ca ions can reach the highest velocities, whose lifetimes allow them to accelerate to high speeds. It is precisely these elements that, after the first ionization, can exhibit the maximum intensity of resonance emission, which can be detected in observations during total solar eclipses or by means of space experiments.

The best candidates for such experiments are the resonance emissions of MgII. The cosmic abundance of Mg exceeds that of Ca by approximately an order of magnitude, and the estimated emission intensity also exceeds the intensity of the CaII K line by roughly an order of magnitude. Unfortunately, the resonance lines of MgI and MgII lie in the near-ultraviolet region, with vacuum wavelengths of about 2853\AA and 2796\AA , respectively. Such observations require more complex instrumentation than observations of lines in the optical range.

Of all atoms and ions having spectral emission lines in the optical range, the most effective may be the detection of the emission of calcium ions. The lifetime of these ions until the next ionization exceeds the lifetime of neutral calcium atoms by three orders

of magnitude and is almost four orders of magnitude greater than the lifetime of sodium atoms.

(Gulyaev and Shcheglov, 1999) were the first to confidently register the resonance emission of CaII during the eclipse of 26 February 1998 in Guadeloupe using a Fabry-Perot etalon (FPE) at several sections of the interferogram to the west of the Sun in the heliocentric distance interval from $5R_{\odot}$ to $20R_{\odot}$. The emission intensity in the CaII K line observed at the center of the terrestrial-sky absorption line during the eclipse turned out to be approximately equal to the intensity of the continuous spectrum in the interval between the lines. In total, nine emission features were recorded in the spectrum. One emission feature proved to be unshifted; two others showed negative velocities of -160 and -168 km/sec. For the remaining six emission features, the shifts of the K line were in the range $2.2\text{-}3.7\text{\AA}$, which corresponds to a radial velocity from 170 to 280 km/sec with a probable error of ± 20 km/sec. The derived velocities for these emission features turned out to be higher than the Keplerian velocities of circular orbital motion for the distances of the region considered, despite the fact that radial velocities were observed, i.e., projections of the spatial velocities onto the line of sight. Naturally, the true spatial velocities may be much larger.

In (Shestakova et al., 2004), an attempt was made to model the orbit of the parent body using emission details (features) of Ca-ion emission obtained from the observations of (Gulyaev and Shcheglov et al., 2001) during the eclipse of 26 February 1998. The simultaneously observed extended emission region cannot be interpreted as the passage of a single parent body. The most probable source of the appearance of cold gas in the outer solar corona may be explained by a scenario in which streams of meteoroids or remnants of comets, moving on elongated orbits, penetrate into the near-solar region. Ca ion particles leaving the parent orbit at each moment of time form an isochrone. All the observed emission is contained within an isochrone of $25 \cdot 10^3$ seconds. During this time, near perihelion the meteor stream (parent bodies) manages to traverse more than 180° in the plane of the sky. Applying the model to the observed results, taking into account the emission configuration and the observed radial velocities, made it possible to estimate the parameters of a parabolic orbit of the parent bodies with a perihelion distance of about $2R_{\oplus}$. The stream of bodies crossed the ecliptic almost at a right angle from south to north, extending along the orbit near perihelion by almost 180° . The total mass of the dispersed material, assuming a chemical composition of CI-type chondrites, is $M_{\text{CI}} = 1 \times 10^{11}$ g. For a porous comet-type body with an average abundance for the Solar System, the mass estimate is $M_{\text{com}} = 2.6 \times 10^{13}$ g, which is equivalent to a body of radius 0.22 km at a density of 0.6 g/cm^3 .

In (Shestakova et al., 2004; 2013), calculations were performed of the acceleration of calcium ions by radiation pressure upon detachment from different points along the orbit from a parent body moving on a parabolic orbit with a perihelion distance $q = 2R_{\oplus}$. It was shown that atoms attain the highest velocities when they detach from the parent body at perihelion. The farther from the Sun the detachment of an atom occurs, the lower its terminal velocity.

Detailed calculations of the acceleration of CaII ions were carried out in (Shestakova

et al., 2013), where an attempt was made to construct a numerical model of the motion of calcium ions after detachment from parent bodies comets, dust grains, or meteoroids following their approach to the Sun.

In the present work, an attempt is made to construct a model of the motion of Ca ions after detachment from the parent body at various distances from the Sun. The influence of radiation pressure, gravity, and magnetic drift in the Sun's radial magnetic field is taken into account. Radiation pressure on calcium ions in the H and K resonance lines of CaII in the solar spectrum is the dominant factor in the dynamics of the ions, which accelerate as they move away from the Sun. The dynamical model makes it possible to estimate the range of distances at which the disintegration of parent bodies and the formation of gas clouds may occur, and to assess the consistency of the proposed acceleration mechanism with the observed velocities (Gulyaev and Shcheglov 1999). Despite the crudeness of the model, one can note its good agreement with the observations.

Method and model. According to the estimates of (Shestakova et al., 1990), the lifetime of a neutral calcium atom is short. At a distance of $6R_{\square}$ from the center of the Sun, it will be ionized within 160 s. The lifetime of the first calcium ion before the next ionization at the same distance from the Sun is already three orders of magnitude larger: 1.5×10^5 s. Owing to the short lifetime of the calcium atom in the solar radiation field, we do not assume the formation of calcium ions by any other mechanisms, but take into account only ionization by solar radiation. Thus, the motion of the neutral atom is excluded from consideration, and we will immediately calculate the motion of the ion after detachment from the parent body. To construct a dynamical model of the motion of CaII ions in the observed region near the Sun, we will rely on the following assumptions:

- 1) Ca atoms are produced as a result of the disintegration of small bodies of the Solar System or of interstellar bodies moving along Keplerian orbits. The origin (interstellar or interplanetary) of the parent body from which the cloud of the observed cold gas formed is of no fundamental importance; therefore, we do not distinguish between interstellar and interplanetary origins of the parent bodies.

- 2) We neglect the influence of the corona and mixing with it, since the relaxation time of the first Ca ion with the corona at a distance of $6R_{\odot}$, according to the theory (Alfven and Feltkhammar, 1967), exceeds by more than an order of magnitude the lifetime of this ion until the next ionization by the Sun's UV radiation. As the relaxation time, we adopt the slowing-down time of calcium ions on protons, which, according to (Alfven and Feltkhammar, 1967), is determined from the relation:

$$\frac{m_i^2 v_i v_T^2}{4\pi \cdot e_i^2 e_p^2 (1 + m_i / m_p) n_p \ln \Lambda \cdot f(v / v_T)} \quad (1)$$

where $v_T = (2kT_p / m_p)^{0.5}$ and for $v = v_T$, $f(v / v_T) = 0.43$. Λ - Debye shielding radius. This quantity, according to (1), is, by our estimates, 1.5×10^5 s, by more than an order of magnitude. Thus, consideration of the processes of interaction of

calcium ions with the coronal material is meaningful only after their transition to the next ionization stage, in which emission in the CaII K line is absent.

- 3) The possible contribution of cyclotron rotation to the observed velocities is not yet taken into account. This contribution may be significant near the starting positions and decreases as the ion moves away from the Sun.

We represent the trajectory of the parent body by the classical equation of elliptical motion:

$$r = \frac{p}{1 + e \cos \varphi} \quad (2)$$

for which the conservation-of-angular-momentum equation, or the areal law $r^2 \dot{\varphi} = \sqrt{\mu \cdot p}$, is also valid. At the moment of detachment of the ion from the parent body, its radial and tangential velocities will be equal to:

$$\dot{r} = \sqrt{\frac{\mu}{p}} e \sin \varphi, \quad r \dot{\varphi} = \sqrt{\frac{\mu}{p}} (1 + e \cos \varphi) \quad (3)$$

Let an ion be separated from its parent body at some location I¹ on the line SI (see Figure 1). The force acting within the ionic radius the vector can be represented as several forces:

$$m \ddot{r} = F_K + F_H + F_g + F_M \quad (4)$$

Where \ddot{r} is the radial acceleration, m is the mass of the Ca, F_K, F_H are the radiation-pressure forces in the CaII K and H resonance lines, F_g is the gravitational force. F_M is the force due to the action of the magnetic field, considered in the first-order approximation, i.e., averaged over one Larmor period. In the Sun's radial field, the action of this force will manifest as a gradient drift toward a weaker field (magnetic mirror acceleration), i.e., away from the Sun. In this case, the entire Equation (3) will be the equation of motion of the guiding center about which the particle undergoes cyclotron rotation. A complete derivation of the general equation of motion is presented in (Shestakova et al., 2013) and the Equation (4) of motion is transformed to a form convenient for numerical solution:

$$\ddot{r} = (0.14782 I_K h_K + 0.078506 I_H h_H + 0.27394 \cdot 10^5 (J \cdot r_{st} / r - 1)) / (r / r_{sol})^2 \text{ cm/sec}^2, \quad (5)$$

where $J = (r \dot{\varphi})^2 / (r \dot{\varphi}_o)^2$ accounts for the difference between the velocity of the parent body and circular Keplerian motion, and is specified by the initial conditions. In all cases considered, for $e \leq 1$, the quantity satisfies $0 < J \leq 2$. The resulting Equation (5)

is used further for calculations of the ion motion, averaged over the Larmor period, after detachment from the parent body.

Many attempts have been made to explain the spatial distribution of dust in the vicinity of the Sun, the orbital distribution of dust particles, the temperature profile, and the boundary of the dust-free zone, where dust completely sublimates and turns into gas (Mann et al., 1996). However, most of these attempts do not have a solid observational basis, because successful observations of the dust corona, which are carried out under conditions of total solar eclipses, are very rare. The question of the location of the boundary of the dust-free zone still remains open.

We pay special attention to the choice of the starting distance of the ions, since the starting distance is not merely a model parameter, but precisely the distance at which the boundary of the dust-free zone or the orbit of the parent body may be located.

The starting distance at which melting of chondritic material begins appears natural. The difficulty is that this material is a mixture of various minerals: olivine, pyroxene, plagioclase, and various impurities. Olivine and pyroxene together constitute three quarters of the mass of meteoritic material (Krinov et al., 1955). Olivine consists of a mixture of forsterite (Mg_2SiO_4) and fayalite (Fe_2SiO_4). Pyroxene consists of a mixture of $MgSiO_3$ and $FeSiO_3$ (Mg-pyroxene and Fe-pyroxene). Plagioclase contains high concentrations of impurities of Al, Ca, and Na.

The most fusible substance among those listed above is Fe-bearing olivine and pyroxene, whose melting temperatures are approximately the same and amount to about 1370 K (Krinov et al., 1955). In the blackbody approximation, bodies at a distance of $9R_{\odot}$ from the center of the Sun will have such a temperature. The refractory mineral forsterite (Mg_2SiO_4) melts at 2160 K (Krinov et al., 1955), which coincides with the blackbody temperature at a distance of $3.6R_{\odot}$. Even closer to the Sun, particles of oxides may approach: alumina (Al_2O_3) and a variety of calcium silicate orthosilicate whose melting temperatures are close to 2300 K and 2400 K, respectively, as well as graphite, whose melting temperature is significantly higher. Near $6R_{\odot}$ particles of the aluminosilicate mullite and plagioclase, which melt at 2100 K (Krinov et al., 1955), and siliceous glass (SiO_2), whose melting temperature is close to 2000 K, also evaporate. Mg-pyroxene and pure iron melt at $5R_{\odot}$.

The distance deserves $9R_{\odot}$ special attention, since about 40% of meteoritic material melts there. This distance we will take as the basis in our calculations. One should also not forget that Ca ions can reach inner regions of the corona down to $(3-4)R_{\odot}$ together with refractory particles of Mg-olivine and plagioclase, which together can constitute up to 35% of meteoritic material. According to our estimates, at distances smaller than $3R_{\odot}$ the relaxation time of a Ca ion with the corona is shorter than the lifetime of the ion until the next ionization by the Sun's UV radiation; therefore, ions penetrating into the inner regions of the corona remain in the corona.

Results. Numerical calculations were carried out using equation (5) for various initial conditions.

- a) Start from a circular orbit, $J = 1$, $\dot{\mathbf{r}} = 0$, $R_{start} = 5R_{\odot}$

b) Start from a parabolic orbit at perihelion, $\phi = 0$, $\dot{r} = 0$, $r\dot{\phi} = 2\sqrt{\mu/p}$, где $p = 2r$, $J = 2$, $R_{start} = 9R_{\odot}$

c) Start from a parabolic orbit at quadrature, $\phi = -90^{\circ}$, $\dot{r} = -\sqrt{\mu/p}$, $r\dot{\phi} = \sqrt{\mu/p}$, $p = r$

In all three cases, variants with other starting distances were also computed. The results of the calculations for ion starts at distances of 9 and 15 solar radii are presented in Figures 1 and 2.




Figure 1. Radial velocities of calcium ions calculated for various initial distances at a heliocentric distance of $100R_{\odot}$ and the terminal velocity at infinity - R_{term} .

Figure 2 presents the results of calculations of the radial velocities at a distance of 100 solar radii and the terminal velocities at infinity.




Figure 2. Velocity of Ca-ions versus distance from the Sun after leaving the parent body. Initial orbits of parent bodies: a) Start from a circular orbit; b) start from a parabolic orbit in perihelion; c) start from a parabolic orbit in quadrature. The upper three curves - start from $9 R_{\odot}$ the lower ones- start from $15 R_{\odot}$.



Figure 3 shows the radial velocities developed by Ca ions when they start from different points of the orbits at distances of $9 R_{\odot}$ and $15 R_{\odot}$. The dynamical analysis shows that ions leaving the orbit of a parent body that penetrates into the near-solar region to distances smaller than $15 R_{\odot}$ can be accelerated to the observed velocities under the action of the forces considered. Thus, the scenario we propose does not contradict the observational results.

In (Gulyaev and Shcheglov, 2001), a second attempt at similar observations with a Fabry-Perot etalon in inclined beams, carried out during the total solar eclipse of 11 August 1999 in Bulgaria, is also described. The results of this observation differ in that strong emission was detected at the centers of the lines with zero Doppler shift. This phenomenon is explained by its temporal coincidence with the maximum of the Perseid meteor stream, which in itself may be of interest as a method for estimating the influx of meteoric material to Earth. Against such a background of powerful emissions, it was difficult to isolate fragments formed in the sublimation region. Nevertheless, three such fragments were identified near an elongation of $20 R_{\odot}$ exhibiting very high radial velocities of about 500 km/sec. The registered fragments were more compact than in the first experiment, and the corresponding radial velocities had different signs: -420 km/sec ($19 R_{\odot}$), +490 km/sec ($21 R_{\odot}$) and 830 km/sec ($24 R_{\odot}$). Thus, in the vicinity of the Sun one can observe the motion of low-ionization Ca ions with radial velocities of different signs and magnitudes close to 1000 km/sec, which does not contradict the model calculations presented in Figures 2 and 3.

The most probable scenario can be considered one in which streams of meteoroids or remnants of comets, moving on elongated orbits, penetrate into the near-solar region. The closer to the Sun the detachment of particles from the parent body occurs, the higher the velocities to which the ions can be accelerated (Figs. 1 and 2).

The best option for extending such studies would be observations outside Earth's atmosphere, which would allow the spectral investigations to be extended into the ultraviolet region.

Possibilities of space observations of resonant luminescence in the H and K lines of CaII.

1) *History of the Search.* Prior to the successful interferometric observations by Gulyaev and Shcheglov, several attempts were made to detect Ca ion resonance emission in the solar F-corona. In July 1990, Aimanova and Shestakova utilized the Nikolsky coronagraph (Almaty) in an attempt to detect CaII H and K line emissions using a spectrograph with a dispersion of approximately 0.3 \AA/mm . A series of consecutive images of the daytime sky at distances from 3 to $9 R_{\odot}$ west of the Sun along the diurnal parallel (at $0.5 R_{\odot}$ intervals) revealed a faint emission band in the CaII K line at $6 R_{\odot}$. Due to time constraints related to an expedition for the total solar eclipse of July 22, 1990, these results could not be verified and remained unpublished.

Subsequently, several more attempts were made to detect the resonance emission of the Ca ion during total solar eclipses. Observations on 22 July 1990 were carried out by A. M. Sredinin according to the program of L. I. Shestakova in Siberia, the settlement of Markovo. An interference filter placed in front of the camera was centered on the CaII

K line and had a width $\delta\lambda_{0.5} \text{ \AA} 100 \text{ \AA}$. The structural features of the outer corona were best seen in the image obtained with an exposure of 30 s, in which the sky background already had a significant density. The central part of the frame out to $6 R_{\odot}$ was strongly overexposed and was burdened by distortions associated with the displacement of the Sun during the exposure, which was made without clock drive. Outside this near-Sun region, after contact copying onto Mikrat-200 film, the sky background was practically eliminated and unusual structures were revealed in the region of the Sun's southern pole: two quasi-radial streams resembling in shape dipole field lines symmetrically curved in opposite directions from the pole. The width of the structures was $3\text{-}4 R_{\odot}$ and their extent reached $15 R_{\odot}$ from the Sun. Owing to the unreliability of the photometry of the original negative, the results also remained unpublished.

The first attempt at interferometric observations without feeding optics, using the setup of P.V. Shcheglov containing a Fabry-Perot etalon in inclined beams, was made on 11 July 1991 in Mexico by A.K. Aimanov, L.I. Shestakova. It turned out that the image with the longest exposure contained emission rings covering the entire field of view, about 30 degrees. After analysis, it was found that the emission rings belonged to mercury, which was established with complete certainty, since a mercury lamp was used as the comparison source. Apparently, at the observing site (La Paz, Mexico) advertising lighting automatically switched on as darkness fell during the total phase of the eclipse; therefore, the sky spectrum contained strong mercury lines.

Simultaneous observations in Mexico using a coronagraph (Aimanov et al., 1995) equipped with an FPE in the exit pupil allowed for radial velocity measurements of dust in the inner corona up to $6 R_{\odot}$. Velocities were estimated via CaII absorption lines; no emission was detected. The interference filter (IF) used was centered at $\lambda 3947 \text{ \AA}$ with a half-width of $\delta\lambda_{1/2} = 50 \text{ \AA}$ to encompass both H and K lines. The mercury line at $\lambda 3906 \text{ \AA}$ did not interfere as it fell outside the filter's passband.

Subsequently, observations by R.A. Gulyaev were carried out on the wide-angle interferometric setup of P.V. Shcheglov with a Fabry-Pérot etalon in inclined beams, whose sensitivity increased from eclipse to eclipse and whose observations ultimately led to success. Interferometric observations on 9 March 1997 in Chita Oblast, performed by R.A. Gulyaev on Shcheglov's setup, showed a high sensitivity level that made it possible to record the sky background at the time of the eclipse, since the entire image was covered by rings of the CaII H and K absorption lines present in the sky spectrum. However, emission was also not detected.

Only during the eclipse of 26 February 1998 in Guadeloupe did interferometric observations of the corona in the CaII H and K lines (Gulyaev and Shcheglov, 2001) show the presence of Ca-ion emission in several regions of the interferogram to the west of the Sun. Compared with 1997, the instrument sensitivity increased by a factor of 40. In the image, the sky absorption lines at the time of the eclipse were clearly visible across the entire field of view. The emission intensity in the CaII K line, observed at the center of the absorption line, turned out to be approximately equal to the intensity of the continuous spectrum in the interval between the lines. As a result, the estimate of the intensity of the continuous spectrum obtained from the image near the CaII H

and K lines is $B = 8.8 \times 10^{-3} \text{ erg}/(\text{cm}^2 \text{ s sr } \text{\AA})$. This means that, for observations with an instrumental profile of width about 1 \AA , the intensity of the observed emission falls within the range of the predicted estimates made in (Shestakova et al., 1990). According to that work, the estimated emission intensity for resonance emission in the CaII K line at a distance of $6 R_{\odot}$ is $1.6 \times (10^{-2} - 10^{-3}) \text{ erg}/(\text{cm}^2 \text{ s sr})$.

2) *Device selection.* Taking into account the results of unsuccessful and successful observational attempts, one can select the following relatively simple instrument configuration: a small-aperture camera with a filter and a wide field of view of $10^{\circ} - 20^{\circ}$, without feeding optics 10° field of view corresponds to $40 R_{\odot}$.

In front of the camera lens with an f-number from 1:1 to 1:1.5 and a diameter of $50 \pm 5 \text{ mm}$ an interference filter (IF) fully covering the light beam is mounted. In the focal plane of the lens, a detector array sensitive in the wavelength range $3900 - 4000 \text{ \AA}$ is installed.

The maximum transmission of the IF, $\lambda_{\text{max}} = 3952 \text{ \AA} \pm 2 \text{ \AA}$, is oriented toward the mean position between the CaII H and K lines, whose vacuum wavelengths are 3969.591 \AA and 3934.777 \AA , respectively. The IF half-width is $\delta\lambda_{1/2} = (52-54) \text{ \AA}$. The filter transmits both CaII H and K resonance lines, taking into account a Doppler shift of $\pm 600 \text{ km/sec}$ for $\delta\lambda_{1/2} = 52 \text{ \AA}$. A Doppler shift of 1 \AA in this region corresponds to a velocity of 76 km/sec . The shape of the filter transmission profile should be close to rectangular.

The main difficulty in conducting observations is shielding the camera from direct sunlight. The localization of the observing region is to the north, east, south, and west of the Sun. The edge of the field of view is oriented at a distance of $4 R_{\odot} = 1^{\circ}$ from the edge of the solar disk ($5 R_{\odot}$ from the center of the Sun). In one observing set, four frames are obtained from all four sides of the Sun. The north-south directions must also be monitored, since outflows from the poles (jets) are also possible.

Testing the camera and selecting the exposure can be carried out under terrestrial conditions at an observatory. The terrestrial sky at full Moon should provide an optimal signal level on the CCD array. The sky glow at full Moon is slightly weaker than the sky during a solar eclipse. The emissions observed in Gulyaev's experiment were comparable in intensity to portions of the continuous sky spectrum during eclipses. According to (Makarova et al., 1994), the brightness of the Sun in the selected spectral interval is about $B_{\odot} \approx 2.8 \times 10^7 \text{ J}/(\text{sec m}^2 \text{ m sr})$. An estimate of the sky brightness at the time of an eclipse, according to (Kimura and Mann, 1998) and (Shestakova and Demchenko, 2010), is $(1-3) \times 10^{-9} B_{\odot}$. As a result, the instrument sensitivity should be no worse than $(2 - 6) \times 10^{-2} \text{ J}/(\text{sec m}^2 \text{ m sr})$. The daytime sky at zenith will also make it possible to select the exposure, taking into account that the sky during a solar eclipse is 5000-10000 times fainter than the daytime sky, as was measured during our observations of solar eclipses.

3) *Expected Observational Results*

Within the field of view, the appearance of emission spots of different sizes is expected, indicating the presence of regions of increased concentration of calcium ions that appeared after the disintegration and dispersion of denser bodies of comet-asteroid composition.

4) Future Research Directions

Possible continuation of the research: scanning the region of the inner Solar System using a telescope with a long-slit spectrograph to measure the radial velocities of emission clouds, in order to investigate the dynamics of these objects and study the sources of their occurrence.

References

- Aimanov A.K., Aimanova G.K., & Shestakova L.I. (1995) Radial velocities in the F-corona on July 11, 1991. *Astronomy Letters*, 21(2). – P. 196-198. NASA ADS: <https://ui.adsabs.harvard.edu/abs/1995AstL...21..196A> (in Eng.).
- Gulyaev R.A., & Shcheglov P.V. (1999a) Emission formations in the F-corona [Emissionnye obrazovaniya v F-korone]. *Doklady RAN*, 366(2). – P. 199-201. (in Rus.).
- Gulyaev R.A., & Shcheglov P.V. (1999b) Interferometric observations of emission features in the F-corona. *Contributions of the Astronomical Observatory Skalnaté Pleso*, 28(3). – P. 237-242. NASA ADS: <https://ui.adsabs.harvard.edu/abs/1999CoSka..28..237G> (in Eng.).
- Gulyaev R.A., & Shcheglov P.V. (1999c) Interferometric observations of the S-corona [Interferometricheskie nablyudeniya S-korony]. *Izvestiya RAN, Seriya Fizicheskaya*, 63(11). – P. 2186-2190. (in Rus.).
- Gulyaev R.A., & Shcheglov P.V. (1999d) Emission structures in the F-corona. *Doklady Physics*, 44(5). – P. 309-311. (in Eng.).
- Gulyaev R.A., & Shcheglov P.V. (1999e) Interferometric observations of the S-corona. *Izvestiya RAN, Seriya Fizicheskaya*, 63(11). – P. 2186-2190. (in Eng.).
- Gulyaev R.A., & Shcheglov P.V. (1999f) Emission formations in the F-corona. *Reports of the Academy of Sciences*, 366(2). – P. 199-201. NASA ADS: <https://ui.adsabs.harvard.edu/abs/1999DokPh..44..309G> (in Eng.).
- Gulyaev R.A., & Shcheglov P.V. (2001) Nablyudeniya rezonansnogo svecheniya atomov v oblastiakh sublimatsii tverdogo veshchestva v blizhnem okolosolnechnom prostranstve [Observations of the resonance glow of atoms in regions of solid material sublimation in the near circumsolar space]. *Uspekhi Fizicheskikh Nauk*, 171(2). – P. 217-219. <https://doi.org/10.3367/UFNr.0171.200102f.0217> (in Rus.).
- Gulyaev R.A., & Shcheglov P.V. (2001). Observations of the resonance glow of atoms in region of solid material sublimation in the near circumsolar space. *Physics-Uspekhi*, 44(2). – P. 203-205. <https://doi.org/10.3367/UFNr.0171.200102f.0217> (in Eng.).
- Kimura H., & Mann I. (1998) Brightness of the solar F-corona. *Earth, Planets and Space*, 50. – P. 493-499. <https://doi.org/10.1186/BF03352140> (in Eng.).
- Krinov E.A. (1955) *Fundamentals of Meteoritics [Osnovy meteoritiki]*. Moscow (in Rus.).
- Makarova E.A., Kazachevskaya T.V., & Kharitonov A.V. (1994) On the variability of some characteristics of solar radiative flux. *Solar Physics*, 152. – P. 195-200. <https://doi.org/10.1007/BF01473204> (in Eng.).
- MacQueen R.M. (1968) Infrared observation of the outer solar corona. *The Astrophysical Journal*, 154. – P. 1059-1076. <https://doi.org/10.1086/149825> (in Eng.).
- Mann I., et al. (2004) Dust near the Sun. *Space Science Reviews*, 110. – P. 269-305. (in Eng.).
- Peterson A.W. (1967) Experimental detection of thermal radiation from interplanetary dust. *The Astrophysical Journal*, 148, L37. <https://doi.org/10.1086/180009> (in Eng.).
- Shestakova L.I. (1990) Is there a region of resonance emission by atoms and ions around the Sun? *Soviet Astronomy Letters*, 16. – P. 236-240. NASA ADS: <https://ui.adsabs.harvard.edu/abs/1990SvAL...16..236S> (in Eng.).
- Shestakova L.I. (2004) Cometary-meteoritic gas in the solar corona: The dynamics of Ca ions. *Solar System Research*, 38(1). – P. 76-84. doi: 10.1023/B:SOLS.0000015158.95033.45 (in Eng.).
- Shestakova L.I. (2013) Movement of CaII ions in the inner Solar system. *Astronomy & Astrophysics Transactions*, 28(2). – P. 109-120. NASA ADS: <https://ui.adsabs.harvard.edu/abs/2013A&AT...28..109S> (in Eng.).

Publication Ethics and Publication Malpractice in the journals of the Central Asian Academic Research Center LLP

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the journals of the Central Asian Academic Research Center LLP implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The Central Asian Academic Research Center LLP follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the Central Asian Academic Research Center LLP.

The Editorial Board of the Central Asian Academic Research Center LLP will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

**[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)
ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)
<http://reports-science.kz/index.php/en/archive>**

Ответственный редактор *А. Ботанқызы*
Редакторы: *Д.С. Аленов, Т. Апендиев*
Верстка на компьютере *Г.Д. Жадырановой*

Подписано в печать 16.03.2026.

Формат 60x88¹/₈.
18,0 п.л. Заказ 1.